


CONTAINS
2 MEDICINES

DIFFERIN[®] EPIDUO[™]

ACNE GEL

**ADAPALENE 0.1% and
BENZOYL PEROXIDE 2.5% GEL
ACNE TREATMENT**



Previously available
only by prescription

Once Daily
Acne Treatment*

FDA approved

*Read consumer
information
leaflet before use

NET WT 0.07 oz (2 g)
Sample
(Not for Sale)

Drug Facts	
Active ingredients Adapalene 0.1% (retinoid).....	Purpose Acne treatment
Benzoyl Peroxide 2.5%.....	Acne treatment
Use ■ For the treatment of acne	
Warnings	
For external use only	
Do not use ■ on damaged skin (cuts, abrasions, eczema, sunburn) ■ if you are allergic to adapalene or any of the ingredients in this product ■ if you have very sensitive skin ■ if you are sensitive to benzoyl peroxide	
Ask a doctor or pharmacist before use if you ■ use any other acne medications ■ have any other skin or medical conditions	
If pregnant or breast-feeding, ask a doctor before use.	
When using this product ■ limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors ■ do not wax to remove hair in areas where the product has been applied ■ avoid contact with hair and dyed fabrics, which may be bleached by this product	
■ during the early weeks of use, your acne may appear to worsen before it improves (this is normal)	
■ irritation (redness, itching, dryness, burning, peeling or possible swelling) is more likely to occur:	
■ in the first four weeks of use ■ if using more than one topical acne medication at a time	
■ if irritation occurs, it may be reduced:	
■ by using the product less frequently ■ by using a moisturizer ■ by using only one topical acne medication at a time	
■ it may take up to 3 months of once daily use to see results ■ avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water. ■ wash hands after use	
Stop use and ask a doctor if ■ you become pregnant, or are planning to become pregnant, while using the product ■ you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath) ■ irritation gets worse ■ you see no improvement after 3 months of once daily use	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions ■ Adults and children 12 years of age and older: ■ use once daily ■ clean the skin gently and pat dry before applying the product ■ cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face. ■ applying more than directed will not provide faster or better results, but may worsen skin irritation ■ if going outside, apply sunscreen after using this product. If irritation or sensitivity develops, stop use of both products and ask a doctor.	
■ Children under 12 years of age: do not use	
Other information ■ store at 20°C to 25°C (68°F to 77°F) ■ protect from freezing ■ protect from light and keep away from heat	
Inactive ingredients acrylamide/sodium acryloyldimethyltaurate copolymer, docusate sodium, edetate disodium, glycerin, isohexadecane, poloxamer 124, polysorbate 80, propylene glycol, purified water, and sorbitan oleate	
Questions? 1-866-735-4137	

All trademarks are the property of their respective owners. Distributed by: Galderma Laboratories, L.P., Fort Worth, TX 76177 USA
Made in Canada 31872-0426

DIFFERIN®
EPIDUO®
ACNE GEL

CONTAINS
2 MEDICINES

DIFFERIN®
EPIDUO®
ACNE GEL

ADAPALENE 0.1% and
BENZOYL PEROXIDE 2.5% GEL
ACNE TREATMENT

DUAL
ACTION



Combines two acne-clearing ingredients
Previously available only by prescription
Dermatologist developed and tested
Once daily topical acne treatment*

*Read consumer information
leaflet before use

NET WT 0.53 oz (15 g)



First FDA-approved
over-the-counter
dual active ingredient
acne treatment

Read carton and enclosed
consumer information leaflet
before using this product. Keep
this carton and consumer
information leaflet. They contain
important information.



ACTUAL
SIZE

P58871-0

Water-based,
oil-free gel
Fragrance free

GALDERMA

Distributed by:
Galderma Laboratories, L.P.,
Fort Worth, TX 76177 USA

All trademarks are the
property of their
respective owners.

Made in Canada
www.differin.com

NDC: 0299-4908-10



LOT:
EXP:

Drug Facts

Active ingredients	Purpose
Adapalene 0.1% (retinoid)*	Acne treatment
*read consumer information leaflet	
Benzoyl Peroxide 2.5%	Acne treatment

Use

- For the treatment of acne

Warnings

For external use only

Do not use

- on damaged skin (cuts, abrasions, eczema, sunburn)
- if you are allergic to adapalene or any of the ingredients in this product
- if you have very sensitive skin
- if you are sensitive to benzoyl peroxide

Ask a doctor or pharmacist before use if you

- use any other acne medications
- have any other skin or medical conditions

If pregnant or breast-feeding, ask a doctor before use.

When using this product

- limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors
- do not wax to remove hair in areas where the product has been applied
- avoid contact with hair and dyed fabrics, which may be bleached by this product
- during the early weeks of use, your acne may appear to worsen before it improves (this is normal)
- irritation (redness, itching, dryness, burning, peeling or possible swelling) is more likely to occur:
 - in the first four weeks of use
 - if using more than one topical acne medication at the same time
- if irritation occurs, it may be reduced:
 - by using the product less frequently
 - by using a moisturizer
 - by using only one topical acne medication at a time
- it may take up to 3 months of once daily use to see results
- avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water.
- wash hands after use

Stop use and ask a doctor if

- you become pregnant, or are planning to become pregnant, while using the product
- you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath)
- irritation gets worse
- you see no improvement after 3 months of once daily use

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Adults and children 12 years of age and older:**
 - use once daily
 - clean the skin gently and pat dry before applying the product
 - cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face.
 - applying more than directed will not provide faster or better results, but may worsen skin irritation
 - if going outside, apply sunscreen after using this product. If irritation or sensitivity develops, stop use of both products and ask a doctor.
- Children under 12 years of age:** do not use

Other information

- store at 20°C to 25°C (68°F to 77°F)
- protect from freezing
- protect from light and keep away from heat

Inactive ingredients

ascydamide/sodium acryloyldimethyltaurate copolymer, docusate sodium, edetate disodium, glycerin, isohexadecane, poloxamer 124, polysorbate 80, propylene glycol, purified water, and sorbitan oleate

Questions? 1-866-735-4137

DIFFERIN[®]
EPIDUO[®]
ACNE GEL

CONTAINS
2 MEDICINES

DIFFERIN[®]
EPIDUO[®]
ACNE GEL

ADAPALENE 0.1% and
BENZOYL PEROXIDE 2.5% GEL
ACNE TREATMENT

DUAL
ACTION



Combines two acne-clearing ingredients
Previously available only by prescription
Dermatologist developed and tested
Once daily topical acne treatment*

*Read consumer information
leaflet before use

NET WT 0.53 oz (15 g)

Tube

Water-based,
oil-free gel
Fragrance free

GALDERMA
Distributed by:
Galderma Laboratories, L.P.,
Fort Worth, TX 76177 USA

All trademarks are the
property of their
respective owners.

Made in Canada
www.differin.com

NDC: 0299-4908-15



Drug Facts

Active ingredients	Purpose
Adapalene 0.1% (retinoid)*	Acne treatment
*read consumer information leaflet	
Benzoyl Peroxide 2.5%	Acne treatment

Use

- For the treatment of acne

Warnings

For external use only

Do not use

- on damaged skin (cuts, abrasions, eczema, sunburn)
- if you are allergic to adapalene or any of the ingredients in this product
- if you have very sensitive skin
- if you are sensitive to benzoyl peroxide

Ask a doctor or pharmacist before use if you

- use any other acne medications
- have any other skin or medical conditions

If pregnant or breast-feeding, ask a doctor before use.

When using this product

- limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors
- do not wax to remove hair in areas where the product has been applied
- avoid contact with hair and dyed fabrics, which may be bleached by this product
- during the early weeks of use, your acne may appear to worsen before it improves (this is normal)
- irritation (redness, itching, dryness, burning, peeling or possible swelling) is more likely to occur:
 - in the first four weeks of use
 - if using more than one topical acne medication at the same time
- if irritation occurs, it may be reduced:
 - by using the product less frequently
 - by using a moisturizer
 - by using only one topical acne medication at a time
- it may take up to 3 months of once daily use to see results
- avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water.
- wash hands after use

Stop use and ask a doctor if

- you become pregnant, or are planning to become pregnant, while using the product
- you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath)
- irritation gets worse
- you see no improvement after 3 months of once daily use

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Adults and children 12 years of age and older:**
 - use once daily
 - clean the skin gently and pat dry before applying the product
 - cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face.
 - applying more than directed will not provide faster or better results, but may worsen skin irritation
 - if going outside, apply sunscreen after using this product. If irritation or sensitivity develops, stop use of both products and ask a doctor.
- Children under 12 years of age:** do not use

Other information

- store at 20°C to 25°C (68°F to 77°F)
- protect from freezing
- protect from light and keep away from heat

Inactive ingredients

acrylamide/sodium acryloyldimethyltaurate copolymer, docusate sodium, edetate disodium, glycerin, isohexadecane, poloxamer 124, polysorbate 80, propylene glycol, purified water, and sorbitan oleate

Questions? 1-866-735-4137

First FDA-approved
over-the-counter
dual active ingredient
acne treatment

Read carton and
enclosed consumer
information leaflet
before using this
product. Keep this
carton and
consumer
information
leaflet. They
contain
important
information.

ACTUAL
SIZE

P59122-0

LOT:
EXP:



DIFFERIN[®]
EPIDUO[®]
ACNE GEL

CONTAINS
2 MEDICINES

DIFFERIN[®]
EPIDUO[®]
ACNE GEL

ADAPALENE 0.1% and
BENZOYL PEROXIDE 2.5% GEL
ACNE TREATMENT

DUAL
ACTION



Combines two acne-clearing ingredients
Previously available only by prescription
Dermatologist developed and tested
Once daily topical acne treatment*

*Read consumer information
leaflet before use

NET WT 1.06 oz (30 g)



Water-based,
oil-free gel
Fragrance free

GALDERMA

Distributed by:
Galderma Laboratories, L.P.,
Fort Worth, TX 76177 USA

All trademarks are the
property of their
respective owners.

Made in Canada

www.differin.com

NDC: XXXX-XXXX-XX



First FDA-approved
over-the-counter
dual active ingredient
acne treatment

Read carton and
enclosed consumer
information leaflet
before using this
product. Keep this
carton and consumer
information leaflet.
They contain
important
information.

ACTUAL
SIZE

PXXXXX-X

LOT:

EXP:

Drug Facts

Active ingredients	Purpose
Adapalene 0.1% (retinoid)*	Acne treatment
*Read consumer information leaflet	
Benzoyl Peroxide 2.5%	Acne treatment

Use
■ For the treatment of acne

Warnings

For external use only

Do not use

- on damaged skin (cuts, abrasions, eczema, sunburn)
- if you are allergic to adapalene or any of the ingredients in this product
- if you have very sensitive skin
- if you are sensitive to benzoyl peroxide

Ask a doctor or pharmacist before use if you

- use any other acne medications
- have any other skin or medical conditions

If pregnant or breast-feeding, ask a doctor before use.

When using this product

- limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors
- do not wax to remove hair in areas where the product has been applied
- avoid contact with hair and dyed fabrics, which may be bleached by this product
- during the early weeks of use, your acne may appear to worsen before it improves (this is normal)
- irritation (redness, itching, dryness, burning, peeling or possible swelling) is more likely to occur:
 - in the first four weeks of use
 - if using more than one topical acne medication at the same time
- if irritation occurs, it may be reduced:
 - by using the product less frequently
 - by using a moisturizer
 - by using only one topical acne medication at a time
- it may take up to 3 months of once daily use to see results
- avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water.
- wash hands after use

Stop use and ask a doctor if

- you become pregnant, or are planning to become pregnant, while using the product
- you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath)
- irritation gets worse
- you see no improvement after 3 months of once daily use

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- **Adults and children 12 years of age and older:**
 - use once daily
 - clean the skin gently and pat dry before applying the product
 - cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face.
 - applying more than directed will not provide faster or better results, but may worsen skin irritation
 - if going outside, apply sunscreen after using this product. If irritation or sensitivity develops, stop use of both products and ask a doctor.
- **Children under 12 years of age:** do not use

Other information

- store at 20°C to 25°C (68°F to 77°F)
- protect from freezing
- protect from light and keep away from heat

Inactive ingredients

acrylamide/sodium acryloyldimethyltaurate copolymer, docosate sodium, edetate disodium, glycerin, isohexadecane, poloxamer 124, polysorbate 80, propylene glycol, purified water, and sorbitan oleate

Questions? 1-866-735-4137

DIFFERIN[®]
EPIDUO
ACNE GEL

CONTAINS
2 MEDICINES

DIFFERIN[®]
EPIDUO
ACNE GEL

ADAPALENE 0.1% and
BENZOYL PEROXIDE 2.5% GEL
ACNE TREATMENT

DUAL
ACTION



Combines two acne-clearing ingredients
Previously available only by prescription
Dermatologist developed and tested
Once daily topical acne treatment*

*Read consumer information
leaflet before use

NET WT 1.6 oz (45 g)



Water-based,
oil-free gel
Fragrance free

GALDERMA

Distributed by:
Galderma Laboratories, L.P.,
Fort Worth, TX 76177 USA

All trademarks are the
property of their
respective owners.


Made in Canada

www.differin.com

NDC: XXXX-XXXX-XX



First FDA-approved
over-the-counter
dual active ingredient
acne treatment



ACTUAL SIZE

PXXXXX-X

LOT:

EXP:

Drug Facts

Active ingredients	Purpose
Adapalene 0.1% (retinoid)*	Acne treatment
*Read consumer information leaflet	
Benzoyl Peroxide 2.5%	Acne treatment

Use

- For the treatment of acne

Warnings

For external use only

Do not use

- on damaged skin (cuts, abrasions, eczema, sunburn)
- if you are allergic to adapalene or any of the ingredients in this product
- if you have very sensitive skin
- if you are sensitive to benzoyl peroxide

Ask a doctor or pharmacist before use if you

- use any other acne medications
- have any other skin or medical conditions

If pregnant or breast-feeding, ask a doctor before use.

When using this product

- limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors
- do not wax to remove hair in areas where the product has been applied
- avoid contact with hair and dyed fabrics, which may be bleached by this product
- during the early weeks of use, your acne may appear to worsen before it improves (this is normal)
- irritation (redness, itching, dryness, burning, peeling or possible swelling) is more likely to occur:
 - in the first four weeks of use
 - if using more than one topical acne medication at the same time
- if irritation occurs, it may be reduced:
 - by using the product less frequently
 - by using a moisturizer
 - by using only one topical acne medication at a time
- it may take up to 3 months of once daily use to see results
- avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water.
- wash hands after use

Stop use and ask a doctor if

- you become pregnant, or are planning to become pregnant, while using the product
- you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath)
- irritation gets worse
- you see no improvement after 3 months of once daily use

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Adults and children 12 years of age and older:
 - use once daily
 - clean the skin gently and pat dry before applying the product
 - cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face.
 - applying more than directed will not provide faster or better results, but may worsen skin irritation
 - if going outside, apply sunscreen after using this product. If irritation or sensitivity develops, stop use of both products and ask a doctor.
- Children under 12 years of age: do not use

Other information

- store at 20°C to 25°C (68°F to 77°F)
- protect from freezing
- protect from light and keep away from heat

Inactive ingredients

acrylamide/sodium acryloyldimethyltaurate copolymer, docusate sodium, edetate disodium, glycerin, isohexadecane, poloxamer 124, polysorbate 80, propylene glycol, purified water, and sorbitan oleate

Questions? 1-866-735-4137

DIFFERIN[®]
EPIDUO
ACNE GEL

CONTAINS
2 MEDICINES

DIFFERIN[®]
EPIDUO
ACNE GEL

ADAPALENE 0.1% and
BENZOYL PEROXIDE 2.5% GEL
ACNE TREATMENT

DUAL
ACTION



Combines two acne-clearing ingredients
Previously available only by prescription
Dermatologist developed and tested
Once daily topical acne treatment*

*Read consumer information
leaflet before use

NET WT 1.6 oz (45 g)



Water-based,
oil-free gel
Fragrance free

GALDERMA

Distributed by:
Galderma Laboratories, L.P.,
Fort Worth, TX 76177 USA

All trademarks are the
property of their
respective owners.

Made in Canada

www.differin.com

NDC: XXXX-XXXX-XX



First FDA-approved
over-the-counter
dual active ingredient
acne treatment

Read carton and
enclosed consumer
information leaflet
before using this
product. Keep this
carton and consumer
information leaflet.
They contain
important
information.

ACTUAL
SIZE

PXXXXX-X

LOT:

EXP:

Drug Facts

Active ingredients	Purpose
Adapalene 0.1% (retinoid)*	Acne treatment
*Read consumer information leaflet	
Benzoyl Peroxide 2.5%	Acne treatment

Use

- For the treatment of acne

Warnings

For external use only

Do not use

- on damaged skin (cuts, abrasions, eczema, sunburn)
- if you are allergic to adapalene or any of the ingredients in this product
- if you have very sensitive skin
- if you are sensitive to benzoyl peroxide

Ask a doctor or pharmacist before use if you

- use any other acne medications
- have any other skin or medical conditions

If pregnant or breast-feeding, ask a doctor before use.

When using this product

- limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors
- do not wax to remove hair in areas where the product has been applied
- avoid contact with hair and dyed fabrics, which may be bleached by this product
- during the early weeks of use, your acne may appear to worsen before it improves (this is normal)
- irritation (redness, itching, dryness, burning, peeling or possible swelling) is more likely to occur:
 - in the first four weeks of use
 - if using more than one topical acne medication at the same time
- if irritation occurs, it may be reduced:
 - by using the product less frequently
 - by using a moisturizer
 - by using only one topical acne medication at a time
- it may take up to 3 months of once daily use to see results
- avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water.
- wash hands after use

Stop use and ask a doctor if

- you become pregnant, or are planning to become pregnant, while using the product
- you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath)
- irritation gets worse
- you see no improvement after 3 months of once daily use

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Adults and children 12 years of age and older:**
 - use once daily
 - clean the skin gently and pat dry before applying the product
 - cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face.
 - applying more than directed will not provide faster or better results, but may worsen skin irritation
 - if going outside, apply sunscreen after using this product. If irritation or sensitivity develops, stop use of both products and ask a doctor.
- Children under 12 years of age:** do not use

Other information

- store at 20°C to 25°C (68°F to 77°F)
- protect from freezing
- protect from light and keep away from heat

Inactive ingredients


acrylamide-sodium acryloyldimethyltaurate copolymer, docusate sodium, edetate disodium, glycerin, isohexadecane, poloxamer 124, polysorbate 80, propylene glycol, purified water, and sorbitan oleate

Questions? 1-866-735-4137

DIFFERIN EPIDUO ACNE GEL	
ADAPALENE 0.1% and BENZOYL PEROXIDE 2.5% GEL ACNE TREATMENT	
SAMPLE (NOT FOR SALE)	
Active ingredients: Adapalene 0.1%, Benzoyl Peroxide 2.5%	
Purpose: Acne Treatment	
Directions: Adults and children 12 years of age and older: Use once daily. Clean the skin gently and pat dry before applying the product. Cover the entire affected area with a thin layer. Children under 12 years of age: do not use.	
Warnings: For external use only. Do not use on damaged skin (cuts, abrasions, eczema, sunburn). If pregnant or breast-feeding, ask a doctor before use. Store at 20°C to 25°C (68°F to 77°F). Protect from freezing.	
Distributed by: Galderma Laboratories, L.P., Fort Worth, TX 76177 USA Made in Canada. Questions or comments? 1-800-735-4137	
www.differin.com PXXXXXX GALDERMA	

Lot:

Exp:



Directions: Adults and children 12 years of age and older: Use once daily. Clean the skin gently and pat dry before applying the product. Cover the entire affected area with a thin layer. Children under 12 years of age: do not use.

Warnings: For external use only. Do not use on damaged skin (cuts, abrasions, eczema, sunburn). If pregnant or breast-feeding, ask a doctor before use. Store at 20°C to 25°C (68°F to 77°F). Protect from freezing.

GALDERMA

Distributed by: Galderma Laboratories, L.P., Fort Worth, TX 76177 USA

All trademarks are the property of their respective owners.

Made in Canada

Questions or comments? 1-866-735-4137

www.differin.com

See outer carton for complete warnings and information.

NET WT 1.53 oz (15 g)	DIFFERIN[®] EPIDUO ACNE GEL		Lot:
	ADAPALENE 0.1% and BENZOYL PEROXIDE 2.5% GEL ACNE TREATMENT		Exp:
<p> Active ingredients: Adapalene 0.1%, Benzoyl Peroxide 2.5% Purpose: Acne Treatment Directions: Adults and children 12 years of age and older: Use once daily. Clean the skin gently and pat dry before applying the product. Cover the entire affected area with a thin layer. Children under 12 years of age: do not use. Warnings: For external use only. Do not use on damaged skin (cuts, abrasions, eczema, sunburn). If pregnant or breast-feeding, ask a doctor before use. Store at 20°C to 25°C (68°F to 77°F). Protect from freezing. Distributed by: Galderma Laboratories, L.P., Fort Worth, TX 76177 USA Made in Canada Questions or comments? 1-866-735-4137 www.differin.com </p>			
See outer carton for complete warnings and information.		GALDERMA	P5xxxx-x



DIFFERIN[®]
EPIDUO
ACNE GEL
ADAPALENE 0.1% and
BENZOYL PEROXIDE 2.5% GEL
ACNE TREATMENT


NET WT 1.06 oz (30g)

Active ingredients: Adapalene 0.1%, Benzoyl Peroxide 2.5%
Purpose: Acne Treatment
Directions: **Adults and children 12 years of age and older:** Use **once** daily. Clean the skin gently and pat dry before applying the product. Cover the entire affected area with a thin layer.
Children under 12 years of age: do not use.
Warnings: **For external use only.**
Do not use on damaged skin (cuts, abrasions, eczema, sunburn).
If pregnant or breast-feeding, ask a doctor before use.
Store at 20°C to 25°C (68°F to 77°F). Protect from freezing.
Distributed by: Galderma Laboratories, L.P., Fort Worth, TX 76177 USA Made in Canada
Questions or comments? 1-866-735-4137 www.differin.com PXXXXX-X
See outer carton for complete warnings and information.

GALDERMA

Lot:
Exp:





Lot:

Exp:

DIFFERIN[®]
EPIDUO
ACNE GEL
ADAPALENE 0.1% and
BENZOYL PEROXIDE 2.5% GEL
ACNE TREATMENT

NET WT 1.6 oz (45 g)

Active ingredients: Adapalene 0.1%, Benzoyl Peroxide 2.5%

Purpose: Acne Treatment

Directions: Adults and children 12 years of age and older: Use once daily. Clean the skin gently and pat dry before applying the product. Cover the entire affected area with a thin layer.

Children under 12 years of age: do not use.

Warnings: For external use only. Do not use on damaged skin (cuts, abrasions, eczema, sunburn). If pregnant or breast-feeding, ask a doctor before use.

Store at 20°C to 25°C (68°F to 77°F). Protect from freezing.

Distributed by: Galderma Laboratories, L.P., Fort Worth, TX 76177 USA Made in Canada

Questions or comments? 1-866-735-4137 www.differin.com PXXXXX-X

See outer carton for complete warnings and information.

GALDERMA

DIFFERIN[®] EPIDUO ACNE GEL

ADAPALENE 0.1% and BENZOYL PEROXIDE 2.5% GEL ACNE TREATMENT

Frequently Asked Questions

What is Differin Epiduo Acne Gel and what is it used for?

- Differin Epiduo Acne Gel is a topical medication used for the treatment of acne in people age 12 years and older.

What should I know before using Differin Epiduo Acne Gel?

- **If pregnant or breast-feeding**, ask a doctor before use.
- Some other retinoid drugs have been shown to cause birth defects. There is no specific evidence that Differin Epiduo Acne Gel causes birth defects in humans when used topically as directed.
- **Do not use** Differin Epiduo Acne Gel if you are allergic to adapalene or any of the ingredients in this product, if you have very sensitive skin or if you are sensitive to benzoyl peroxide.
- **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

How often do I apply Differin Epiduo Acne Gel?

- Apply Differin Epiduo Acne Gel **only one time a day** and try to apply the product at the same time each day.

How do I apply Differin Epiduo Acne Gel?

- Gently clean the affected areas using a mild (non-irritating) cleanser and pat dry.
- Apply Differin Epiduo Acne Gel as a thin layer to the affected areas of the skin **only one time a day**. For example, if you get acne on the face, clean, dry and apply the product to the entire face. Differin Epiduo Acne Gel is **not** a spot treatment and should not be used to treat a single pimple.
- Avoid contact with eyes, lips and mouth. **If contact occurs, immediately flush the area with water.**
- Do not apply to damaged skin (cuts, abrasions, eczema, or sunburn).
- Wash hands after use.

How long will it take for Differin Epiduo Acne Gel to work?

- It may take up to 3 months of daily use for results to appear.
- Do not use more than one time a day. Applying more than directed will not provide faster or better results, but may worsen skin irritation.

What do I do if I need to be in the sun?

- When possible, limit sun exposure, including light from tanning beds.
- When going outdoors, use a sunscreen as labeled. Your skin may be more sensitive while using Differin Epiduo Acne Gel. If you use Differin Epiduo Acne Gel during the day, allow it to dry before applying sunscreen.



DIFFERIN[®] EPIDUO ACNE GEL

ADAPALENE 0.1% AND BENZOYL PEROXIDE 2.5% GEL ACNE TREATMENT

Frequently Asked Questions (Continued)

Can I use a moisturizer if my skin is dry?

- Yes, use of a moisturizer may help decrease dryness and other signs of irritation.
- Avoid products that may dry or irritate your skin such as harsh soaps, astringents, cosmetics that have strong skin-drying effects and products containing high levels of alcohol.

When is my skin most likely to become irritated? And what do I do?

- Irritation (redness, itching, dryness, burning) is more likely to occur:
 - In the first few weeks of use.
 - if using abrasive skin cleansers, products with drying effects or more than one topical acne medication at a time.
You may want to delay starting Differin Epiduo Acne Gel until any irritation from using the other product(s) has gone away.
 - in extreme weather such as wind and cold.
- Moisturizers may be used.
- Irritation usually lessens after 4 weeks of continued use.

What do I do if my skin becomes severely irritated?

- If irritation gets worse, stop use and ask a doctor before using Differin Epiduo Acne Gel again.

Can I remove unwanted facial hair by waxing while using Differin Epiduo Acne Gel?

- Do not use wax to remove hair in areas where Differin Epiduo Acne Gel has been applied because it may worsen skin irritation.

What ingredients are used in Differin Epiduo Acne Gel?

- Differin Epiduo Acne Gel contains:
The active ingredients are adapalene 0.1% and benzoyl peroxide 2.5%. The other ingredients are: acrylamide/sodium acryloyldimethyltaurate copolymer, docusate sodium, edetate disodium, glycerin, isohexadecane, poloxamer 124, polysorbate 80, propylene glycol, purified water, and sorbitan oleate.

How should I store Differin Epiduo Acne Gel?

- Differin Epiduo Acne Gel should be stored at 20°C to 25°C (68°F to 77°F). Protect from freezing.
- Protect from light and keep away from heat.
- Do not use Differin Epiduo Acne Gel after the expiry date marked on the packaging.

Other Questions? Where can I get more information?

Phone: 1-866-735-4137

See outer carton for complete warnings and information.

PXXXXX-X

DIFFERIN[®] EPIDUO

ACNE GEL ADAPALENE 0.1% and BENZOYL PEROXIDE 2.5% GEL ACNE TREATMENT

Frequently Asked Questions

What is Differin Epiduo Acne Gel and what is it used for?

- Differin Epiduo Acne Gel is a topical medication used for the treatment of acne in people age 12 years and older.

What should I know before using Differin Epiduo Acne Gel?

- **If pregnant or breast-feeding**, ask a doctor before use.
- Some other retinoid drugs have been shown to cause birth defects. There is no specific evidence that Differin Epiduo Acne Gel causes birth defects in humans when used topically as directed.
- **Do not use** Differin Epiduo Acne Gel if you are allergic to adapalene or any of the ingredients in this product, if you have very sensitive skin or if are you sensitive to benzoyl peroxide.
- **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

How often do I apply Differin Epiduo Acne Gel?

- Apply Differin Epiduo Acne Gel **only one time a day** and try to apply the product at the same time each day.

How do I apply Differin Epiduo Acne Gel?

- Gently clean the affected areas using a mild (non-irritating) cleanser and pat dry.
- Apply Differin Epiduo Acne Gel as a thin layer to the affected areas of the skin **only one time a day**. For example, if you get acne on the face, clean, dry and apply the product to the entire face. Differin Epiduo Acne Gel is **not** a spot treatment and should not be used to treat a single pimple.
- Avoid contact with eyes, lips and mouth. **If contact occurs, immediately flush the area with water.**
- Do not apply to damaged skin (cuts, abrasions, eczema, or sunburn).
- Wash hands after use.

How long will it take for Differin Epiduo Acne Gel to work?

- It may take up to 3 months of daily use for results to appear.
- Do not use more than one time a day. Applying more than directed will not provide faster or better results, but may worsen skin irritation.

What do I do if I need to be in the sun?

- When possible, limit sun exposure, including light from tanning beds.
- When going outdoors, use a sunscreen as labeled. Your skin may be more sensitive while using Differin Epiduo Acne Gel. If you use Differin Epiduo Acne Gel during the day, allow it to dry before applying sunscreen.

Can I use a moisturizer if my skin is dry?

- Yes, use of a moisturizer may help decrease dryness and other signs of irritation.
- Avoid products that may dry or irritate your skin such as harsh soaps, astringents, cosmetics that have strong skin-drying effects and products containing high levels of alcohol.

When is my skin most likely to become irritated? And what do I do?

- Irritation (redness, itching, dryness, burning) is more likely to occur:
 - In the first few weeks of use.
 - if using abrasive skin cleansers, products with drying effects or more than one topical acne medication at a time. You may want to delay starting Differin Epiduo Acne Gel until any irritation from using the other product(s) has gone away.
 - in extreme weather such as wind and cold.
- Moisturizers may be used.
- Irritation usually lessens after 4 weeks of continued use.

What do I do if my skin becomes severely irritated?

- If irritation gets worse, stop use and ask a doctor before using Differin Epiduo Acne Gel again.

Can I remove unwanted facial hair by waxing while using Differin Epiduo Acne Gel?

- Do not use wax to remove hair in areas where Differin Epiduo Acne Gel has been applied because it may worsen skin irritation.

What ingredients are used in Differin Epiduo Acne Gel?


- Differin Epiduo Acne Gel contains:
The active ingredients are adapalene 0.1% and benzoyl peroxide 2.5%. The other ingredients are: acrylamide/sodium acryloyldimethyltaurate copolymer, docusate sodium, edetate disodium, glycerin, isohexadecane, poloxamer 124, polysorbate 80, propylene glycol, purified water, and sorbitan oleate.

How should I store Differin Epiduo Acne Gel?

- Differin Epiduo Acne Gel should be stored at 20°C to 25°C (68°F to 77°F). Protect from freezing.
- Protect from light and keep away from heat.
- Do not use Differin Epiduo Acne Gel after the expiry date marked on the packaging.

Other Questions? Where can I get more information?

Phone: 1-866-735-4137



First FDA-approved over-the-counter dual active ingredient acne treatment

Water-based, oil-free gel | Fragrance free

Combines two acne-clearing ingredients

Previously available only by prescription

Dermatologist developed and tested

Read enclosed Drug Facts and consumer information leaflet before using this product. Keep the Drug Facts and consumer information leaflet. They contain important information.

DIFFERIN EPIDUO ACNE GEL
ADAPALENE 0.1% AND BENZOYL PEROXIDE 2.5% GEL ACNE TREATMENT

DUAL ACTION **2 in 1** **ACNE CLEARING**
 CONTAINS 2 MEDICINES
 25 Tubes
 NET WT 0.07 OZ (2 g) Per Tube

SAMPLE (NOT FOR SALE)

Drug Facts

Active ingredients
 Adapalene 0.1% (Differin) Acne treatment
 Benzoyl Peroxide 2.5% (Epiduo) Acne treatment

Use
 For the treatment of acne

Warnings
 For external use only
 Do not use if you experience skin rash, abrasions, sores, sunburn, or if you are allergic to adapalene or any of the ingredients in this product. If you have very sensitive skin, if you are sensitive to benzoyl peroxide, or if you are pregnant, stop use of this product. Use only after other acne medications. Use any other oral or medical condition.

Drug Facts (continued)

When using this product Avoid sun exposure, including light from tanning beds, and use sunscreen when going outdoors. Do not use in areas where the product has been applied. Avoid contact with hair and eyes. Avoid contact with face and hair. Avoid contact with hair and eyes. Avoid contact with face and hair. Avoid contact with face and hair.

Directions Apply a thin layer of this product to the entire face, avoiding the eyes, mouth, and nostrils. Use only after other acne medications. Use any other oral or medical condition.

Drug Facts (continued)

Directions Apply a thin layer of this product to the entire face, avoiding the eyes, mouth, and nostrils. Use only after other acne medications. Use any other oral or medical condition.

Other information Store at 20°C to 25°C (68°F to 77°F). Product from tracking.

Inactive ingredients arylmethylethylphosphorothioic acid, octadecyl sodium, sodium dodecyl sulfate, glycerin.

Drug Facts (continued)

Directions Apply a thin layer of this product to the entire face, avoiding the eyes, mouth, and nostrils. Use only after other acne medications. Use any other oral or medical condition.

Other information Store at 20°C to 25°C (68°F to 77°F). Product from tracking.

Inactive ingredients arylmethylethylphosphorothioic acid, octadecyl sodium, sodium dodecyl sulfate, glycerin.

Questions? 1-866-795-4137

GALDERMA
 Distributed by Galderma Laboratories, L.P., Fort Worth, TX 76177 USA
 All trademarks are the property of their respective owners.
 Made in Canada
 www.differin.com PXXXXXX

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
05/22/2026 08:26:11 AM