

CENTER FOR DRUG EVALUATION AND RESEARCH

Application Number 21-330

FINAL PRINTED LABELING



Drug Facts		Drug Facts (continued)							
Active ingredient (in each lozenge)	Purpose	Directions							
Nicotine polacrilex, 2mg	Stop smoking aid	<ul style="list-style-type: none"> stop smoking completely when you begin using the lozenge if you smoke your first cigarette within 30 minutes of waking up, use 4mg nicotine lozenge if you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine lozenge according to the following 12 week schedule: 							
Use		<table border="1"> <thead> <tr> <th>Weeks 1 to 6</th> <th>Weeks 7 to 9</th> <th>Weeks 10 to 12</th> </tr> </thead> <tbody> <tr> <td>1 lozenge every 1 to 2 hours</td> <td>1 lozenge every 2 to 4 hours</td> <td>1 lozenge every 4 to 8 hours</td> </tr> </tbody> </table>		Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12	1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours
Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12							
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours							
Warnings		<ul style="list-style-type: none"> nicotine lozenge is a medicine and must be used a certain way to get the best results place the lozenge in your mouth and allow the lozenge to slowly dissolve (about 20 - 30 minutes). Minimize swallowing. Do not chew or swallow lozenge. you may feel a warm or tingling sensation occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 20-30 minutes) do not eat or drink 15 minutes before using or while the lozenge is in your mouth to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day. stop using the nicotine lozenge at the end of 12 weeks. If you still feel the need to use nicotine lozenges, talk to your doctor. 							
Do not use		Other information							
<ul style="list-style-type: none"> if you continue to smoke, chew tobacco, use snuff, or use a nicotine patch or other nicotine containing products 		<ul style="list-style-type: none"> Phenylketonurics: Contains Phenylalanine 3.4 mg per lozenge store at 20 - 25°C (68 - 77°F) protect from light 							
Ask a doctor before use if you have		Inactive ingredients							
<ul style="list-style-type: none"> heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. high blood pressure not controlled with medication. Nicotine can increase your blood pressure. stomach ulcer or diabetes 		acacia, aspartame, calcium polycarboxophil, corn syrup solids, flavors, lactose, magnesium stearate, maltodextrin, mannitol, potassium bicarbonate, sodium alginate, sodium carbonate, soy protein, triethyl citrate, xanthan gum							
Ask a doctor or pharmacist before use if you are		Questions or comments?							
<ul style="list-style-type: none"> using a non-nicotine stop smoking drug taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted. 		call weekdays 1-888-569-1743 (10:00am - 4:30pm EST)							
Stop use and ask a doctor if									
<ul style="list-style-type: none"> mouth problems occur persistent indigestion or severe sore throat occurs irregular heartbeat or palpitations occur you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat 									
Keep out of reach of children and pets.									
Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.									
Directions									
<ul style="list-style-type: none"> if you are under 18 years of age, ask a doctor before use before using this product, read the enclosed User's Guide for complete directions and other important information 									

Commit Lozenge
72ct. 2mg
Back/Top Panel

From the marketers of
Nicorette®
Commit™

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Use at least 9 lozenges of Commit per day during the first six weeks.
3. **Use Long Enough** - Use Commit for the full 12 weeks.
4. **Use With a Support Program** as directed in the enclosed User's Guide.

		
To remove the lozenge, tear off single unit.	Peel off backing starting at corner with loose edge.	Push lozenge through foil.

*The American Cancer Society supports the use of stop smoking aids and counseling as effective tools when quitting smoking but does not endorse any specific product. GlaxoSmithKline pays a fee to the American Cancer Society for the use of its logo.

COMMIT™, COMMITTED QUITTERS®, and associated logo designs and overall trade dress designs are trademarks owned and/or licensed to GlaxoSmithKline or its affiliated companies.

Distributed By:
GlaxoSmithKline Consumer Healthcare, L.P.
Pittsburgh, PA 15230, Made in the U.S.A.
©2003 GlaxoSmithKline


00000XX

From the marketers of
Nicorette®
Commit™

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

Blister packaged for your protection.
Do not use if individual seals are open or torn.


Internal theft surveillance system



0 0000-0000-00 0

Commit Lozenge
72ct. 4mg
Side Panels

2mg



GlaxoSmithKline		PACKAGING SPECIFICATION	Date: 6.24.02
ID•M Imaging & Design Management • 100 Beecham Drive, Pittsburgh, PA 15205 • (800) 456-6670 • E-mail address: idm.i.graphics@sb.com			
BRAND: Commit	UPC/SKU NO.: n/a		
PROD. CATEGORY: Smoking Cessation	E.V. NO.: n/a		
DESCRIPTION: Blister	DIMENSIONS: 80mm x 105mm		
COMPONENT: Lozenge	DIE NO.: NPL002PM		
VOLUME CONTENTS/SIZE: 72ct 2mg	MANUFACTURING SITE: Aiken		
FORM NO.: 57974XA	PRINTER: Quint		
COUNTRY: USA	PREPRESS: Quint		
PMS COLORS: Black	REPLACES FORM NO.: New		
KEYWORDS:	PRF NO.: 211-02		
Artist: Rick Tenney	Revision 1: rt-6.26.02-refit copy to die		
Computer Software: Adobe Illustrator 8.0	Revision 2: rt-6.26.02-added "peel here" copy		

PS03

©2002 GlaxoSmithKline. All Rights Reserved. All logos, designs and artwork are trademarks, trade dress and/or copyrighted material, owned by and/or licensed to GlaxoSmithKline and/or its affiliated companies. Unauthorized reproduction is strictly prohibited.

- PRINTER**
PLEASE NOTE YOUR RESPONSIBILITIES:
- Use existing die. Die line for art position only.
 - Verify all bar codes meet ANSI standards.
 - Fonts included solely for this artwork and recreating this specific file. GlaxoSmithKline requires that all fonts be destroyed upon completion of this project.
 - These files are not trapped.

PMS COLOR CHIPS:



GlaxoSmithKline		Regulatory Artwork Approval
SIGNATURE:	DATE:	
COMMENTS:		

RF 602

T
SITE, Aiken
O. Nitw
12 ds - Type correction "COMMITTED"

ss and/or copyrighted material, owned by
rized reproduction is strictly prohibited.

NOTE YOUR RESPONSIBILITIES:
isting die. Die line for art position only.
all bar codes meet ANSI standards.
cluded solely for this artwork and
ing this specific file. GlaxoSmithKline
is that all fonts be destroyed upon
ation of this project.
files are not trapped.



EV Code
for Proofing

for multi-color
foil stamp.

Includes User's Guide
**FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE
WITHIN 30 MINUTES OF WAKING UP.**
If you smoke your first cigarette MORE THAN 30 MINUTES after waking up:
use Commit 2mg Lozenge

4mg

72 LOZENGES, 4mg EACH

Pantone 349 screen overprints
Pantone 347 screen to create
gradation shown above.

Commit
Nicorette
From the marketers of

Drug Facts	Drug Facts (continued)						
<p>Active Ingredient (in each lozenge) nicotine polacrilex 4mg</p> <p>Purpose Stop smoking aid</p> <p>Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.</p> <p>Warnings If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking while using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.</p> <p>Do not use • if you continue to smoke, chew tobacco, use snuff, or use a nicotine patch or other nicotine-containing products.</p> <p>Ask a doctor before use if you have • heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. • high blood pressure not controlled with medication. Nicotine can increase your blood pressure. • stomach pain or diabetes.</p> <p>Ask a doctor or pharmacist before use if you are • using a non-nicotine stop smoking drug • taking prescription medicine for depression or asthma. Your or condition may need to be adjusted.</p> <p>Stop use and ask a doctor if • mouth problems occur • persistent indigestion or severe sore throat occurs • irregular heartbeat or palpitations occur • you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat.</p> <p>Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions • if you are under 18 years of age, ask a doctor before use • before using the product, read the enclosed User's Guide for complete directions and other important information.</p>	<p>Directions • stop smoking completely when you begin using the lozenge • if you smoke your first cigarette more than 30 minutes after waking up, use 4mg nicotine lozenge according to the following 12 week schedule.</p> <table border="1"> <thead> <tr> <th>Weeks 1 to 6</th> <th>Weeks 7 to 9</th> <th>Weeks 10 to 12</th> </tr> </thead> <tbody> <tr> <td>1 lozenge every 1 to 2 hours</td> <td>1 lozenge every 2 to 4 hours</td> <td>1 lozenge every 4 to 6 hours</td> </tr> </tbody> </table> <p>• nicotine lozenge is a medicine and must be used a certain way to get the best results • place the lozenge in your mouth and allow the lozenge to slowly dissolve (about 20 - 30 minutes). Minimize swallowing. Do not chew or swallow lozenge • you may feel a warm or tingling sensation • occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 20-30 minutes) • do not eat or drink 15 minutes before using or while the lozenge is in your mouth • to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks • do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you dizziness, numbness, nausea or other side effects • do not use more than 6 lozenges in 6 hours. Do not use more than 30 lozenges per day • stop using the nicotine lozenge at the end of 12 weeks. If you still feel the need to use nicotine lozenges, talk to your doctor.</p> <p>Other Information • (Methylxanthines) Caffeine (Theophylline) 3.4 mg per lozenge • store at 20 - 25°C (68 - 77°F) • protect from light</p> <p>Inactive ingredients aspartame, calcium polyacrylate, flavors, magnesium stearate, mannitol, phenylalanine, sucraloside, sodium alginate, sodium carbonate, sorbitol gum</p> <p>Questions or comments? call weekdays 1-800-418-4188 (10:00am - 4:30pm EST)</p>	Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12	1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 6 hours
Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12					
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 6 hours					

From the marketers of
Nicorette

Commit

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use Enough - Use at least 9 lozenges if Commit per day during the first six weeks.
3. Use Long Enough - Use Commit for the full 12 weeks.
4. Use With a Support Program as directed in the enclosed User's Guide.

To remove the lozenge, use all single use.

Place all backing staining at center with lower edge.

Press lozenge through lip.

*GlaxoSmithKline provides an annual grant to the American Cancer Society for stop smoking education and cancer-related research in return for the use of their best logo design and overall trade dress designs are trademarks owned and/or licensed to GlaxoSmithKline or its affiliate companies.

©2002 GlaxoSmithKline

0288022401

2880

From the marketers of
Nicorette

NEW

Commit

nicotine polacrilex lozenge, 4mg
**STOP SMOKING AID
LOZENGE**

Includes User's Guide
**FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE
WITHIN 30 MINUTES OF WAKING UP.**
If you smoke your first cigarette MORE THAN 30 MINUTES after waking up:
use Commit 2mg Lozenge

4mg

72 LOZENGES, 4mg EACH

From the marketers of
Nicorette


Con

not for sale to those under 18 years of age. Proof of age required. Not for sale in vending machines where proof of age cannot be provided.


R blister packaged for your convenience. Do not use if individual blister pack is damaged.

0766-33C

BEST POSSIBLE COPY



1/mg



gsk GlaxoSmithKline		PACKAGING SPECIFICATION	Date: 6.24.02
I•D•M Imaging & Design Management • 100 Beecham Drive, Pittsburgh, PA 15205 • (800) 456-6670 • E-mail address: idm.i.graphics@sb.com			
BRAND: Commit	UPC/SKU NO.: n/a		
PROD. CATEGORY: Smoking Cessation	E.V. NO.: n/a		
DESCRIPTION: Blister	DIMENSIONS: 80mm x 105mm		
COMPONENT: Lozenge	DIE NO.: NPL0002PM		
VOLUME CONTENTS/SIZE: 72ct 4mg	MANUFACTURING SITE: Aiken		
FORM NO.: 57975XA	PRINTER: Quint		
COUNTRY: USA	PREPRESS: Quint		
PMS COLORS: Black	REPLACES FORM NO.: New		
KEYWORDS:	PRF NO.: 211-02		
Artist: Rick Tenney	Revision 1: rt-6.26.02-refit copy to die		
Computer Software: Adobe Illustrator 8.0	Revision 2: rt-6.26.02-added "peel here"		

©2002 GlaxoSmithKline. All Rights Reserved. All logos, designs and artwork are trademarks, trade dress and/or copyrighted material, owned by and/or licensed to GlaxoSmithKline and/or its affiliated companies. Unauthorized reproduction is strictly prohibited.

gsk GlaxoSmithKline		Regulatory Artwork Approval
SIGNATURE:	DATE:	
COMMENTS:		

REG02

PRINTER
PLEASE NOTE YOUR RESPONSIBILITIES:

- Use existing die. Die line for art position only.
- Verify all bar codes meet ANSI standards.
- Fonts included solely for this artwork and recreating this specific file. GlaxoSmithKline requires that all fonts be destroyed upon completion of this project.
- These files are not trapped.

PMS COLOR CHIPS:



Black

WALLET CARD

WHERE TO CALL FOR HELP:

American Lung Association
1-800-586-4572

American Cancer Society
1-800-227-2345

American Heart Association
1-800-242-8721

Quitting Buddy or Friend who has Quit

©2002 GlaxoSmithKline

CommitTM

nicotine polacrilex lozenge
2mg and 4mg User's Guide

**HOW TO USE
CommitTM LOZENGES
AND TIPS TO HELP YOU
QUIT SMOKING.**

©2002 GlaxoSmithKline

59185XA

- Not for sale to those under 18 years of age.
- Proof of age required.
- Not for sale in vending machines or from any source where proof of age cannot be verified.

©2002 GlaxoSmithKline

PLANNING YOUR SUCCESS

- 1) The key to accomplishing anything important is commitment. When it comes to quitting smoking, that is especially true. **Commit™** Lozenges can help if you really want to quit. **Commit™** Lozenges help reduce withdrawal symptoms including nicotine craving associated with quitting smoking.
- 2) Your chances of staying off cigarettes are much better if you start with at least 9 **Commit™** Lozenges daily. For best results, use the lozenges on a regular schedule (as outlined in this User's Guide).
- 3) Stop smoking completely when you start using **Commit™** Lozenges. Even a single cigarette is likely to put you right back to square one.
- 4) This User's Guide outlines a 12-week plan for **Commit™** Lozenges. Even though you may feel confident about your non-smoking status after a few

weeks, it's important to stick with the plan to help you remain smoke free.

- 5) **Commit™** Lozenges work best when used together with a support plan. See insert between pages 8 and 9 for instructions on enrollment in the Committed Quitters Personalized Stop Smoking Plan.
- 6) After the first six weeks, start using fewer **Commit™** Lozenges, as directed in the Instructions, so you can become smoke-and nicotine-free by the end of the 12 week plan.
- 7) If you have questions about using **Commit™** Lozenges, call 1-800-419-4766 weekdays (10:00am - 4:30pm EST), or talk to your pharmacist or family doctor.

YES! YOU WANT TO QUIT.

Wonderful. You've made the most important decision of all, to stop smoking. And by choosing **Commit™** Lozenges to help you, you're starting on the right path. Now remember, using **Commit™** doesn't just mean taking a

©2002 GlaxoSmithKline

1

Commit™ Lozenge. It means setting and following a program like the one we suggest in this User's Guide.

Your own success depends on your effort, your level of addiction to tobacco, and your commitment to following your program.

LET'S FACE IT.

Quitting smoking isn't easy! You or someone you know may have tried unsuccessfully. That's okay. It's hard to stop smoking the first time you try. The important part is to learn from your previous attempts, consider what went wrong and keep trying to quit until you succeed.

Look to this User's Guide for support as you undergo this terrific task. The guide includes important information on how to use **Commit™** Lozenges and also gives you tips to help you stop smoking. Refer back to it often for advice, answers, and encouragement to help you stay on track.

2

©2002 GlaxoSmithKline

GET MOTIVATED. STAY MOTIVATED.

Everyone has a reason for quitting—whether you're concerned about your health, your appearance, family or peer pressure, or the effect of secondhand smoke on your loved ones—all of the above, or something else entirely. Whatever your reasons, write them down. There's a wallet card inside the back cover of this User's Guide. Write your reasons on the card and carry it with you. When you have an urge to smoke or experience a difficult moment it can help you focus on your reasons for quitting. Lots of people quit with a co-worker, spouse or friend and use them as a quitting buddy. You can help each other out by providing extra encouragement in tough moments.

There may be support groups in your area for people trying to quit. Call your local chapter of the American Lung Association, American Cancer Society or American Heart Association for further information. Toll free phone numbers are printed on the wallet card on the back cover of this User's Guide.

UNDERSTANDING THE DOUBLE-EDGED SWORD.

Smoking has two addictive components, a physical and a mental need for the nicotine in tobacco. You need to conquer both to succeed. **Commit™** Lozenges can ease your physical nicotine addiction. But your readiness and resolve are necessary to help overcome the mental side of your cigarette dependence. So once you're ready, it's time to begin. But first, read and consider the following important warnings.

IMPORTANT WARNINGS

This product is only for those who want to stop smoking.

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Do not use

- if you continue to smoke, chew tobacco, use snuff, or use a nicotine patch or other nicotine containing products.

Ask a doctor before use if you have

- heart disease, recent heart attack or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcers or diabetes.

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur

©2002 GlaxoSmithKline

3

Stop use and ask a doctor if (cont.)

- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat

Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

YOU'RE READY TO START.

Okay, you're ready. To become a non-smoker, start today. Now before you do anything else, you have a bit of planning to do. Read this User's Guide all the way through. You want to make sure you bought the right dose to start. If you typically smoke your first cigarette within 30 minutes of waking up use the 4mg Commit™ Lozenges. If you smoke your first cigarette more than 30 minutes after waking up use the 2mg Commit™ Lozenges. Next, plan your

quitting schedule. Get a calendar to follow your progress and mark the following four important dates (see the stickers in the middle of this booklet).

THE PROGRAM

STEP 1. (Weeks 1-6) Starting on your quit date it's best to use at least 9 Commit™ Lozenges each day, one every 1 - 2 hours.

First choose the day you plan to quit (make it soon). Place the Step 1 sticker on this date. That's the day you will stop smoking cigarettes completely and start using Commit™ Lozenges to calm your cravings for nicotine and help you stay smoke free. Prior to the quit date, get rid of all your cigarettes to remove temptations and make it more difficult to start smoking again.

Use a Commit™ Lozenge every 1 to 2 hours and at least 9 lozenges each day for the first 6 weeks to help prevent unexpected cravings and improve your chances of quitting. These aren't ordinary lozenges. Place the lozenge in your mouth and allow the lozenge to

slowly dissolve (about 20 - 30 minutes). Minimize swallowing. Do not chew or swallow the lozenge. You may feel a warm or tingling sensation. Occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 20 to 30 minutes). Remember to read the instructions on page 7 before you take your first Commit™ Lozenge.

STEP 2. (The next three weeks, that is weeks 7-9). At the beginning of week 7 start using fewer Commit™ Lozenges, one every 2-4 hours.

After six weeks, you should wait a little longer between lozenges, one lozenge every two to four hours. This will help you gradually use fewer Commit™ Lozenges. Put the Step 2 sticker on the first day of week 7 to help remind you when to start reducing the number of Commit™ Lozenges you take.

STEP 3. (The last three weeks, that is weeks 10-12). At the beginning of week 10, reduce Commit™ Lozenge use even further, one every 4-8 hours.

At the beginning of week 10 further decrease the number of Commit™ Lozenges you use each day to reduce the amount of nicotine you get. You should do this by using one lozenge every 4 to 8 hours. Put the Step 3 sticker on the first day of week 10 so you know when you should be starting this last step to becoming smoke and nicotine-free.

END. At the end of week 12 stop using Commit™ Lozenges to become both cigarette and nicotine-free.

Put the "EX-SMOKER" sticker on your calendar on the date 12 weeks after the day you stopped smoking and started using Commit™ Lozenges. You should not use Commit™ Lozenges beyond this date.

BE PREPARED.

Since smoking is an addiction, it is hard to quit. Even after you stop, there will be times when you WANT a cigarette, sometimes strongly. (See also section on "Challenges To Watch For"). The best

defense is to be prepared. Plan now for handling tough times so you don't give in. For example: think about situations when you usually get a craving for cigarettes or where you think you might experience strong cravings. Try to avoid these situations where you can (for example, avoid spending time with smokers, or drinking alcohol, if those things tempt you to smoke).

Change your habits. For example, take your coffee break somewhere else. Take a walk. In other words, break the association between your usual habits and cigarettes.

If you do encounter a situation where you feel a strong craving, fight it! Take a break from the situation; keep yourself busy or distracted with other activities. Remind yourself why you want to quit, and above all, remind yourself that having "just one" really will hurt your goal of quitting!

To prepare for tough situations, assemble a "survival package"—items that can

keep you distracted in case you get a craving. For example, you may include cinnamon gum or hard candy, relaxing music, and things to keep your hands busy like a smooth stone, paper clips, or a rubber ball.

Track your progress as you quit. Keep a journal. Write down how many pieces of Commit™ Lozenges you use each day. Note if and when you get a craving. If you slip and have a cigarette, don't give up. Stop smoking again and get back on your program with Commit™ Lozenges.

Establish your support network. Keep friends' and family members' phone numbers ready to get the moral support you need. Before quitting, ask friends and family to support and encourage you. Think of specific ways they can help.

Reward yourself. Set aside little gifts to yourself such as a CD or video, which you can earn by overcoming difficult hurdles.

HOW Commit™ LOZENGES WORK. Commit™ Lozenges are a form of Nicotine Replacement Therapy. They deliver nicotine to your body, temporarily relieving craving and nicotine withdrawal symptoms when you quit smoking. But unlike cigarettes, Commit™ Lozenges deliver a lower, steady level of nicotine to your blood. When used as directed, Commit™ Lozenges help you regulate, control, and gradually reduce your body's craving for nicotine.

The good news is that Commit™ Lozenges contain no tar or carbon monoxide, and therefore don't present the same medical risks as cigarettes.

However, the lozenges still deliver nicotine, the addictive ingredient in cigarettes. And for some people the nicotine in Commit™ Lozenges can occasionally cause mouth or throat irritation, headaches, nausea, hiccups, upset stomach or dizziness.

USING Commit™ LOZENGES PROPERLY. Remember, Commit™ Lozenges aren't like ordinary lozenges such as cough drops. This lozenge is designed to deliver nicotine into your system through the lining of your mouth, not in your stomach like most other medicines. It is important to minimize swallowing the dissolved medicine in these lozenges so that it can be properly absorbed in your mouth.

Do not use more than one lozenge at a time, or many lozenges one after another since this can cause hiccups, heartburn, nausea or other side effects.

Read all the following instructions before using Commit™ Lozenges. Refer to them often to make sure you're using Commit™ Lozenges correctly.

IMPORTANT: Don't worry or give up if you do not like the taste of the lozenge at first. Commit™ Lozenges are a medication, not a candy. Most people get used to the taste after a day or two. Remember, staying with the plan will help you quit. Stop smoking completely before you start using Commit™ Lozenges.

- 1) Remove the Commit™ Lozenge from the blister. Place the lozenge in your mouth and allow the lozenge to slowly dissolve (about 20 – 30 minutes). Minimize swallowing. Do not chew or swallow the lozenge. You may feel a warm or tingling sensation.
- 2) Occasionally move the lozenge from one side of your mouth to the other side until completely dissolved (about 20-30 minutes).


To reduce cravings or urges to smoke and other withdrawal symptoms, use Commit™ Lozenges according to the following dosage schedule.

Weeks 1 through 6	Weeks 7 through 9	Weeks 10 through 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

Do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day. Stop using the lozenge at the end of 12 weeks (3 months).

FOR THE BEST CHANCE OF QUITTING, use Commit™ Lozenges on a regular schedule, using at least 9 lozenges a day during the first 6 weeks. That will help your body better adjust to the lack of cigarettes and better help prevent cravings. Some people may need more lozenges to reduce their cravings. Do not exceed the recommended maximum daily dosage of 20 lozenges per day. Do not continuously use one lozenge after another, since this may cause you hiccups, heartburn, nausea or other side effects.

PLACE THESE STICKERS ON YOUR CALENDER:



STEP 1
1 lozenge every
1 to 2 hours

STEP 2
1 lozenge every
2 to 4 hours

AT BEGINNING OF WEEK #1
(QUIT DATE)

AT BEGINNING OF WEEK #7

©2002 GlaxoSmithKline



**COMMITTED
QUITTERS**

Personalized Stop Smoking Plan
brought to you by

Commit[™]
nicotine polacrilex lozenge
and
GlaxoSmithKline

COMMIT[™], COMMITTED QUITTERS[®], and associated logo designs and overall trade dress designs are trademarks owned and/or licensed to GlaxoSmithKline or its affiliated companies.

FREE Personalized
Stop Smoking Plan

What is **COMMITTED QUITTERS**

TO JOIN COMMITTED QUITTERS

Personalized Stop Smoking Plan


Enroll online
at
www.committedquitters.com
or call 1-800-770-0708

and ask for your **FREE** Personalized Stop Smoking Plan

- Provide your **PERSONAL CODE**, printed on the back of this insert.
- You will be personally interviewed by a plan specialist to understand your specific needs and to design a Stop Smoking Plan just for you.

• A **FREE**, custom-tailored plan to help you break the psychological addiction to smoking.

• Throughout your quit attempt, you will receive personalized advice on how to cope with situations that make you want to smoke.



© 2002 GlaxoSmithKline

Read and follow label directions
PERSONAL USE

Call Between 7 am and 12 midnight EST
or enroll online 24 hours a day
(ONE PLAN PER CUSTOMER)

**COMMITTED
QUITTERS**

PLACE THESE STICKERS ON YOUR CALENDER:

**STEP
3**
1 lozenge every
4 to 8 hours

EX-SMOKER

AT BEGINNING OF WEEK #10 12 WEEKS AFTER QUIT DATE

©2002 GlaxoSmithKline

Do not eat or drink 15 minutes before using or while the lozenge is in your mouth.

CUTTING BACK ON YOUR Commit™ LOZENGE USAGE.

The whole reason for using Commit™ Lozenges is to decrease and slowly eliminate your need for nicotine, while you control cravings. So, as the above schedule indicates, you should gradually reduce the amount of Commit™ Lozenges you take per day. Some people find it easier to reduce by substituting ordinary sweets or sugar free candy for some of the Commit™ Lozenges they would normally use. As time goes on, you can increase the number of pieces of candy as you further reduce your use of Commit™ Lozenges. Stop using Commit™ Lozenge at the end of week 12. If you still feel the need to use Commit™ Lozenge after week 12, talk with your doctor.

MAKE QUITTING EASIER ON YOURSELF.

Soon after your quit date, parties, bars,

celebrations, and socializing may all tempt you to smoke. Please remember these tips to help you resist those urges and stay smoke-free.

The Day You Quit Smoking:

- Look to your family and friends for support. Let them know what to do or avoid doing to help you quit.
- Throw away ALL cigarettes, ashtrays, matches, lighters. You don't need them. You don't want them and you want to make it difficult to go back.
- Keep yourself occupied. Take a walk. See a movie. See friends. Do anything to keep your mind off cigarettes.
- Calculate all the money you'll save by not buying cigarettes. Probably well over \$1,000 a year! \$1,000 a year! Think of what you can spend it on!
- Know what situations are going to make you want to smoke. Plan now how you'll avoid them or deal with them so you don't smoke.
- Keep Commit™ Lozenges next to your

©2002 GlaxoSmithKline

9

bed so you're prepared when you get up. A lot of people get cravings first thing in the morning.

- Make an appointment to see your dentist and get the tobacco stains cleaned off. While you're getting rid of the evidence of cigarettes in the house, do the same for your teeth. Have clothes or drapes that smell of smoking cleaned.
- Now that your house is smoke-free, try to spend most of your time in smoke-free environments.
- If you usually smoked with coffee or alcohol, try to keep away from them for now. Remember you are also trying to break a habit.
- Smoking is a "hands-on" habit. So use something else to occupy your hands: a rubber band or a pen.
- Now's a good time to get active. Find activities to take your mind off cigarettes and relax. Take up jogging, swimming, or walking.

10

©2002 GlaxoSmithKline

- Don't stress out about gaining weight. Dieting now may weaken your efforts to quit smoking. Eat sensibly and exercise daily; drink large quantities of water and fruit juices; this can help your chances of staying smoke-free.
- Laugh. Watch a sitcom. Read a comic book. It really helps.

REMEMBER: Urges to smoke are temporary. They'll pass, even if you don't smoke.

WHAT YOU CAN EXPECT.

As you are successful at staying smoke-free, initially you will probably notice a few of the following typical withdrawal symptoms, so don't be surprised. Use of **Commit™** Lozenges reduces these symptoms, but may not eliminate them entirely. They will go away with time. Stay focused on your goal of becoming an ex-smoker. Research shows that if you manage to avoid all smoking in the first week (that means not having a single puff), your chances of success increase dramatically.

The First Few Days. You may feel nervous or irritable or have difficulty concentrating during the first few days after you quit smoking. Your body needs time to regain balance. Initially, you might feel a little out of sorts, get headaches, feel light-headed, or have trouble sleeping. Your smoker's cough may get worse before it improves. But fear not, it's a positive sign. Coughing helps clean your lungs of the tar residue you got from smoking.

After a Couple of Weeks. Your confidence and ability to cope with urges to smoke should be getting stronger. But don't be over-confident and think you can smoke just one cigarette. Even now, having even a single puff can lead to a return to smoking cigarettes regularly. Be prepared and remember why you wanted to stop smoking.

Have you noticed that your sense of taste and smell has improved? You are probably coughing less and finding it easier to breathe. You've also probably

noticed your withdrawal symptoms are subsiding (though don't worry if they're still there; they last longer for some people). These are all positive signs that your body is getting used to your success at stopping smoking.

By The End of The first Month. You are less likely to have cravings for cigarettes as often. However sudden cravings may still happen, and when they do, be on your guard, as they can be strong and seem to come out of the blue. Be prepared for these challenging times. The key is do what you can so these unexpected cravings can't beat you. Keep focused on the ways non-smokers are more attractive than smokers. Their breath smells better. Their clothes and hair are fresher. Their teeth are cleaner and brighter. Their skin is less likely to wrinkle. Not smoking around children and your friends is also healthier for them too.

What if You Do Slip And Smoke!
"What if I relapse?" One cigarette is a slip-up, but it's not the end of the quit

©2002 GlaxoSmithKline

11

effort. Everybody slips at something. The key is this: forgive yourself and stop at that one cigarette. Don't let this slip ruin your good intentions, keep at your quit attempt. So, throw out your cigarettes and continue with your quit attempt, keeping in mind what went wrong and led to the slip.

If you do go back to smoking, certainly don't throw out your Commit™ Lozenges. Keep them for the next time you're ready to quit. In fact research says that even if you are back to smoking regularly the best thing you can do is learn and try again.

Try to understand the reason you had those cigarettes that made you slip. That's important, because now you can plan better to deal with these moments next time. It's true you stumbled, but don't think of yourself as having failed. Encourage yourself by treating the last attempt as a learning experience, even a "trial run" for the real thing.

Take a look at the usage instructions and check that you used the Commit™

Lozenges correctly and for the full 12 weeks of the program. When you try again make sure you use enough and the right way. That way you'll be best equipped to deal with the unexpected cravings.

Don't forget; quitting isn't easy and it takes practice to do anything. Stopping smoking is no different.

YOU'VE MADE IT.

Once your twelve week quitting program is over, you've taken your last Commit™ Lozenge. Now you are both cigarette and nicotine-free. Get up and give yourself a standing ovation. We mean it. Do you realize that you have just done a really difficult thing!

Now's a good time to think back on the process. Think of all your reasons for quitting smoking. Think of your goals. Think of how they're going to be a reality now.

Think of what you're going to do with your newly liberated cigarette money. The places you can now go smoke-free.

Think of the extra time you may have added to your life and what you can do with it. And although you may still experience the occasional temptation, and cigarettes still want you back, think positively. Think forward. And consider yourself a proud non-smoker.

FREQUENTLY ASKED QUESTIONS.

1. When I stop smoking and start using Commit™ Lozenges how will I feel?
Commit™ Lozenges help reduce cravings, but be prepared for some nicotine withdrawal symptoms. After you stop smoking they can begin almost at once and are normally at their strongest during the first three or four days. For some people, any of the following may occur:

- unexpected craving or urges for cigarettes
- anxiety, irritability, restlessness, mood changes, nervousness
- drowsiness
- trouble concentrating
- increased appetite and weight gain

- headaches, muscular pain, constipation, fatigue

Commit™ Lozenges are designed to reduce the craving for nicotine you used to satisfy with cigarettes. Commit™ Lozenges can also help provide relief from other withdrawal symptoms such as irritability and nervousness.

2. Are Commit™ Lozenges just swapping one type of nicotine addiction for another?
Commit™ Lozenges do contain nicotine, however there is probably less nicotine in your daily dose of lozenges than in your cigarettes. Commit™ Lozenges give you enough nicotine to help you combat the physical withdrawal symptoms so you can cope with the mental side of stopping smoking. Also, since the nicotine from the lozenges goes into your blood stream more slowly, it produces less of the effects of nicotine that people find rewarding. In fact, when used as directed in the 12 week program, Commit™ Lozenges gradually wean you off your dependence for both nicotine and

cigarettes. Remember, don't use Commit™ Lozenges together with nicotine patches or other nicotine containing products.

3. Can Commit™ Lozenges do any harm?

Some people with conditions like heart disease or people taking prescription medicine for asthma or depression should not use this product without talking to their doctor – check the IMPORTANT WARNINGS on page 3. You may also experience side effects such as hiccups, mouth or throat irritation, heartburn or other stomach problems such as nausea especially if Commit™ Lozenges are chewed or swallowed. In any case, Commit™ Lozenges do not contain the tar, carbon monoxide, and other toxins present in cigarette smoke.

4. Will I put on weight?

In the first couple of months after quitting smoking, some people do put on a few pounds. But think of it this way. Overall, you'll be healthier and look

better. You can always tackle your weight by changing your diet and increasing the amount you exercise once you have gotten through the difficult part of stopping smoking.

5. Does taking Commit™ Lozenges cost more than smoking?

If you normally smoke a pack and a half a day, your total cost of using Commit™ Lozenges during the 12-week period is about the same as smoking. But guess what? After you've finished the Commit™ Lozenge program all that money you used to spend on cigarettes is now savings. And think of the health issues you'll hopefully be able to avoid.

6. What if I have a cigarette and start smoking?

Don't panic. First, don't think badly of yourself. Throw away your cigarettes and forgive yourself. Then think about what went wrong and get back on track. In fact people who have already tried to stop smoking are more likely to be successful the next time.

CHALLENGES TO WATCH FOR.

Once you quit smoking, you are likely to experience periodic, and sometimes intense, temptations to smoke. Certain situations present special challenges. Some common ones include:

Stress and upset.

When you are feeling stressed or upset, you may think a cigarette will make everything better. It won't. Find other ways to relax and unwind.

The blues.

You may be especially vulnerable when you feel bored or blue. Remember that having a cigarette will just make you feel worse.

Smoking cues.

Seeing cigarettes or watching other people smoke can trigger temptation. Remember that you choose not to smoke anymore.

Alcohol.

Drinking and smoking seem to go together, and alcoholic beverages may weaken your resolve, making drinking dangerous to your quit effort. Avoid

drinking early in your quit effort, and try to drink with non-smokers.

Automatic slips.

Sometimes you may find yourself preparing to smoke without even realizing it. Watch out for those moments when your hand seems to 'automatically' reach for a cigarette. Watch out for these situations; they can trigger a relapse. You probably know which one (s) are most dangerous for you; plan ahead to deal with the situation effectively. Always remember that you're trying to break a habit, and the most important thing is to do something to combat the urge in these situations.

COPING AFTER QUITTING.

The key to staying smoke-free is to prepare for and cope with challenges as they occur. If you find yourself tempted to smoke, do something! Here are some things to consider.

- **Escape.** Leave the situation, even for a few minutes. Most temptations don't last long.
- **Distract yourself.** Get your mind off smoking. Think of something else or get busy with something.
- **Relax.** Don't let stress get to you. Think of pleasant relaxing things; breathe slowly and regularly. Let the stress drain out of you.
- **Talk yourself out of it.** What you say to yourself matters. So, remind yourself how important it is for you to quit; remind yourself you can't have just one; or just command yourself to **STOP.**

16

©2002 GlaxoSmithKline

59185XA

WALLET CARD

My most important reasons to quit smoking are:

©2002 GlaxoSmithKline

gsk GlaxoSmithKline		PACKAGING SPECIFICATION	Date: 07.29.02
I•D•M Imaging & Design Management • 100 Beecham Drive, Pittsburgh, PA 15205 • (800) 456-6670 • E-mail address: idm.i.graphics@gsk.com			
BRAND: Commit	UPC/SKU NO.: N/A		
PROD. CATEGORY: Smoking Cessation	E.V. NO.: 9956 (1 2 of 5)		
DESCRIPTION: Nicotine Polacrilex Lozenge	DIMENSIONS: 100mm (W) x 75mm (H) Folded		
COMPONENT: User's Guide	DIE NO.: NPL0002BR		
VOLUME CONTENTS/SIZE: 72ct and 168ct	MANUFACTURING SITE: Aiken		
FORM NO.: 59185XA	PRINTER: Color Art		
COUNTRY: U.S.A.	PREPRESS: Filmet		
PMS COLORS: 4 color process	REPLACES FORM NO.: New		
KEYWORDS:	PRF NO.: 259-02		
Artist: Dan Sommer	Revision 1:		
Computer Software: Adobe Illustrator v. 8.0	Revision 2:		

©2002 GlaxoSmithKline.

FS03

All Rights Reserved. All logos, designs and artwork are trademarks, trade dress and/or copyrighted material, owned by and/or licensed to GlaxoSmithKline and/or its affiliated companies. Unauthorized reproduction is strictly prohibited.

gsk GlaxoSmithKline		Regulatory Artwork Approval
SIGNATURE:	DATE:	
COMMENTS:		

REG02

PRINTER

PLEASE NOTE YOUR RESPONSIBILITIES:

- Use existing die. Die line for art position only.
- Verify all bar codes meet ANSI standards.
- Fonts included solely for this artwork and recreating this specific file. GlaxoSmithKline requires that all fonts be destroyed upon completion of this project.
- These files are not trapped.

PMS COLOR CHIPS:



**This is a representation of an electronic record that was signed electronically and
this page is the manifestation of the electronic signature.**

/s/

Mary Robinson
10/25/02 03:49:41 PM
INTERDISCIPLINARY

Helen Cothran
10/25/02 04:26:36 PM
INTERDISCIPLINARY

**This is a representation of an electronic record that was signed electronically and
this page is the manifestation of the electronic signature.**

/s/

Mary Robinson
10/25/02 03:49:41 PM
INTERDISCIPLINARY

Helen Cothran
10/25/02 04:26:36 PM
INTERDISCIPLINARY

Helen Cothran
10/25/02 04:47:38 PM
INTERDISCIPLINARY