

CENTER FOR DRUG EVALUATION AND RESEARCH

APPROVAL PACKAGE FOR:

APPLICATION NUMBER


21-137

Approved Labeling

DESCRIPTION

Levolet® (levothyronine sodium tablets, USP) contains synthetic crystalline L-3,3',5,5'-tetraiodo-L-thyronine sodium salt [levothyronine (T₄) sodium]. Synthetic T₄ is identical to that produced in the human thyroid gland.

Levothyronine (T₄) sodium has an empirical formula of C₁₅H₁₁I₄NaO₄ · H₂O, molecular weight of 798.95 g/mol (anhydrous), and structural formula as shown.



Inactive Ingredients: Croscarmellose sodium, magnesium stearate and microcrystalline cellulose. This product also contains 333 mcg potassium iodides per tablet (see CONTRAINDICATIONS and PRECAUTIONS, Pregnancy). The following are the coloring additives per tablet strength.

Table with 2 columns: Tablet Strength (mg) and Color additives. Lists various strengths from 0.05 to 0.3 mg and their corresponding color additives like FD&C Yellow No. 6, FD&C Blue No. 1, etc.

CLINICAL PHARMACOLOGY

Thyroid hormone synthesis and secretion is regulated by the hypothalamic-pituitary-thyroid axis. Thyrotropin-releasing hormone (TRH) released from the hypothalamus stimulates secretion of thyrotropin-stimulating hormone (TSH) from the anterior pituitary. TSH, in turn, is the physiologic stimulus for the synthesis and secretion of thyroid hormones, L-thyronine (T₃) and L-triiodothyronine (T₃), by the thyroid gland. Circulating serum T₃ and T₄ levels exert a feedback effect on both TRH and TSH secretion. When serum T₃ and T₄ levels increase, TRH and TSH secretion decrease. When thyroid hormone levels decrease, TRH and TSH secretion increase.

The mechanisms by which thyroid hormones exert their physiologic actions are not completely understood, but it is thought that their principal effects are exerted through control of DNA transcription and protein synthesis. T₃ and T₄ diffuse into the cell nucleus and bind to thyroid receptor proteins attached to DNA. This hormone nuclear receptor complex activates gene transcription and synthesis of messenger RNA and cytoplasmic proteins.

Thyroid hormones regulate multiple metabolic processes and play an essential role in normal growth and development, and normal maturation of the central nervous system and bone. The metabolic actions of thyroid hormones include augmentation of cellular respiration and thermogenesis, as well as metabolism of proteins, carbohydrates and lipids. The protein anabolic effects of thyroid hormones are essential to normal growth and development.

The physiological actions of thyroid hormones are produced predominantly by T₃, the majority of which (approximately 80%) is derived from T₄ by deiodination in peripheral tissues.

Levothyronine, at doses individualized according to patient response, is effective as replacement or supplemental therapy in hypothyroidism of any etiology, except transient hypothyroidism during the recovery phase of subacute thyroiditis.

Levothyronine is also effective in the suppression of pituitary TSH secretion in the treatment or prevention of various types of eut thyroid goiters, including thyroid nodules, Hashimoto's thyroiditis, multinodular goiter and as adjunctive therapy in the management of thyrotropin-dependent well-differentiated thyroid cancer (see INDICATIONS and USAGE, PRECAUTIONS, and DOSAGE AND ADMINISTRATION).

PHARMACOKINETICS

Absorption: Absorption of orally administered T₄ from the gastrointestinal (GI) tract ranges from 40% to 80%. The majority of the levothyronine dose is absorbed from the antrum and upper ileum. The relative bioavailability of Levolet tablets, compared to an equal normal dose of oral levothyronine sodium solution, is approximately 84.6% - 98.3%. T₄ absorption is increased by fasting, and decreased in malabsorption syndromes and by certain foods such as soybean infant formula. Dietary fiber decreases bioavailability of T₄. Absorption may also decrease with age. In addition, many drugs and foods affect T₄ absorption (see PRECAUTIONS, Drug Interactions and Drug-Food Interactions).

Distribution: Circulating thyroid hormones are greater than 99% bound to plasma proteins, including thyroxine-binding globulin (TBG), thyroxine-binding prealbumin (TBPA), and albumin (TBA), whose capacities and affinities vary for each hormone. The higher affinity of both TBG and TBPA for T₄ partially explains the higher serum levels, slower metabolic clearance, and longer half-life of T₄ compared to T₃. Protein-bound thyroid hormones exist in reverse equilibrium with small amounts of free hormone. Only unbound hormone is metabolically active. Many drugs and physiologic conditions affect the binding of thyroid hormones to serum proteins (see PRECAUTIONS, Drug Interactions and Drug-Laboratory Test Interactions). Thyroid hormones do not readily cross the placental barrier (see PRECAUTIONS, Pregnancy).

Metabolism: T₄ is slowly eliminated (see Table 1). The major pathway of thyroid hormone metabolism is through sequential deiodination. Approximately eighty-percent of circulating T₄ is derived from peripheral T₄ by mono-deiodination. The liver is the major site of degradation for both T₃ and T₄, with T₄ deiodination also occurring at a number of additional sites, including the kidney and other tissues. Approximately 80% of the daily dose of T₄ is deiodinated to yield equal amounts of T₃ and reverse T₃ (rT₃). T₃ and rT₃ are further deiodinated to diiodothyronine. Thyroid hormones are also metabolized via conjugation with glucuronides and sulfates and excreted directly into the bile and gut, where they undergo enterohepatic recirculation.

Elimination: Thyroid hormones are primarily eliminated by the kidneys. A portion of the conjugated hormone reaches the colon which is then excreted in the feces. Approximately 20% of T₄ is eliminated in the stool. Urinary excretion of T₄ decreases with age.

Table 1: Pharmacokinetic Parameters of Thyroid Hormones in Euthyroid Patients

Table with 5 columns: Hormone, Time to Peak (hr), Bioactive Potency, Half-life (days), and Area Under Curve (AUC). Rows for Levothyronine (T4) and Liothyronine (T3).

INDICATIONS AND USAGE

Levothyronine sodium is used for the following indications:

Hypothyroidism: As replacement or supplemental therapy in congenital or acquired hypothyroidism of any etiology, except transient hypothyroidism during the recovery phase of subacute thyroiditis. Specific indications include: primary (myxedema), secondary (secondary), and tertiary (hypothalamic) hypothyroidism and subclinical hypothyroidism. Primary hypothyroidism may result from functional deficiency, primary atrophy, partial or total congenital absence of the thyroid gland, or from the effects of surgery, radio-iodine, or drugs, with or without the presence of goiter.

Pituitary TSH Suppression: In the treatment or prevention of various types of eut thyroid goiters (see WARNINGS and PRECAUTIONS), including thyroid nodules (see WARNINGS and PRECAUTIONS), subacute or chronic lymphocytic thyroiditis (Hashimoto's thyroiditis), multinodular goiter (see WARNINGS and PRECAUTIONS), and as an adjunct to surgery and radioactive therapy in the management of thyrotropin-dependent well-differentiated thyroid cancer.

CONTRAINDICATIONS

Levothyronine is contraindicated in patients with untreated subclinical (suppressed serum TSH level with normal T₃ and T₄ levels) or overt thyrotoxicosis of any etiology and in patients with acute myocardial infarction. Levothyronine is contraindicated in patients with uncorrected adrenal insufficiency, since thyroid hormones may precipitate an acute adrenal crisis by increasing the metabolic clearance of glucocorticoids (see PRECAUTIONS). Levolet is contraindicated in patients with hypersensitivity to any of the inactive ingredients in Levolet tablets. (See DESCRIPTION, Inactive Ingredients.)

Pregnancy and Lactation: The extra iodine contained in Levolet may increase the risk of suppression of thyroid function in the fetus or nursing infant, particularly when the mother is already ingesting iodides from other sources. It is therefore recommended that non-iodine containing thyroid hormone preparations be used in pregnant women, women who may become pregnant, or nursing mothers. (See PRECAUTIONS, Pregnancy Category.)

WARNINGS

WARNING: Thyroid hormones, including Levolet, either alone or with other therapeutic agents, should not be used for the treatment of obesity or for weight loss. In euthyroid patients, doses within the range of daily hormonal requirements are ineffective for weight reduction. Excessive doses may

Information for Patients

Patients should be informed of the following information to aid in the safe and effective use of Levolet.

- 1. Notify your physician if you are allergic to any iodine or medicines are pregnant or intend to become pregnant, or are breast feeding, or are taking any other medications...
2. It is recommended that you substitute a non-iodine containing thyroid hormone preparation for Levolet if you plan to become pregnant...
3. Notify your physician if you experience any of the following symptoms: rapid or irregular heartbeat, chest pain, shortness of breath, leg cramps, headache, nervousness, irritability, sleeplessness, tremors, change in appetite, weight gain or loss, vomiting, diarrhea, excessive sweating, heat intolerance, fever, changes in menstrual periods, hives or skin rash, or any other unusual medical event.

Laboratory Tests

General: The diagnosis of hypothyroidism is confirmed by measuring TSH levels using a sensitive assay (second generation assay sensitivity < 0.1 mIU/L or third generation assay sensitivity < 0.01 mIU/L) and measurement of free-T₄.

The adequacy of therapy is determined by periodic assessment of appropriate laboratory tests and clinical evaluation. The choice of laboratory tests depends on various factors including the etiology of the underlying thyroid disease, the presence of concomitant medical conditions, including pregnancy, and the use of concomitant medications (see PRECAUTIONS, Drug Interactions and Drug-Laboratory Test Interactions). Persistent clinical and laboratory evidence of hypothyroidism despite an apparent adequate replacement dose of Levolet may be evidence of inadequate absorption, poor compliance, drug interactions, or decreased T₄ potency of the drug product.

Adults

In adult patients with primary (thyroidal) hypothyroidism serum TSH levels (as measured by a sensitive assay) alone may be used to monitor therapy. The frequency of TSH monitoring during levothyronine dose titration depends on the clinical situation, but it is generally recommended at 6-8 week intervals until normalization for patients who have recently initiated levothyronine therapy and whose serum TSH has normalized, or in patients who have had low dosage of brand of levothyronine changed, the serum TSH concentration should be measured after 8-12 weeks. When the optimum replacement dose has been attained, clinical (physical examination) and biochemical monitoring may be performed every 6-12 months, depending on the clinical situation, and whenever there is a change in the patient's status. It is recommended that a physical examination and a serum TSH measurement be performed at least annually in patients receiving Levolet (see WARNINGS, PRECAUTIONS, and DOSAGE AND ADMINISTRATION).

Pediatrics

In patients with congenital hypothyroidism, the adequacy of replacement therapy should be assessed by measuring both serum TSH (using a sensitive assay) and total- or free-T₄. During the first three years of life, the serum total- or free-T₄ should be maintained at all times in the upper half of the normal range. While the aim of therapy is to normalize the serum TSH level, this is not always possible in a small percentage of patients, particularly in the first few months of therapy. TSH may not normalize due to a resetting of the pituitary-thyroid feedback threshold as a result of an in utero hypothyroidism. Failure of the serum T₄ to increase into the upper half of the normal range within 2 weeks of initiation of Levolet therapy and/or of the serum TSH to decrease below 20 mIU/L within 4 weeks should alert the physician to the possibility that the child is not receiving adequate therapy. Careful inquiry should then be made regarding compliance, dose of medication administered, and method of administration, prior to raising the dose of Levolet.

The recommended frequency of monitoring of TSH and total- or free-T₄ in children is as follows: at 2 and 4 weeks after the initiation of treatment; every 1-2 months during the first year of life; every 2-3 months between 1 and 3 years of age; and every 3 to 12 months thereafter until growth is completed. More frequent intervals of monitoring may be necessary if poor compliance is suspected or abnormal values are obtained. It is recommended that TSH and T₄ levels, and a physical examination, if indicated, be performed 2 weeks after any change in Levolet dosage. Routine clinical examination, including assessment of mental and physical growth and development, and bone maturation, should be performed at regular intervals (see WARNINGS, PRECAUTIONS, Pediatric Use, and DOSAGE AND ADMINISTRATION).

Secondary (pituitary) and tertiary (hypothalamic) hypothyroidism

Adequacy of therapy should be assessed by measuring serum free-T₄ levels, which should be maintained in the upper half of the normal range in these patients.

Drug Interactions

Many drugs affect thyroid hormone pharmacokinetics and metabolism (e.g., absorption, synthesis, secretion, catabolism, protein binding, and target tissue response) and may alter the therapeutic response to Levolet. In addition, thyroid hormones and thyroid status have varied effects on the pharmacokinetics and actions of other drugs. A listing of drug-thyroid axis interactions is contained in Table 2.

The list of drug-thyroid axis interactions in Table 2 may not be comprehensive due to the introduction of new drugs that interact with the thyroid axis or the discovery of previously unknown interactions. The prescriber should be aware of this fact and should consult appropriate reference sources. (e.g., package inserts of newly approved drugs, medical literature) for additional information if a drug-drug interaction with levothyronine is suspected.

Table 2: Drug-Thyroid Axis Interactions

Table with 3 columns: Drug or Drug Class, Effect, and Details. Lists various drug classes like Dopamine/Dopamine Agonists, Glucocorticoids, etc., and their effects on thyroid hormone secretion or absorption.

Levolet NDA 21-137

Pg 1

absorption is approximately 84% - 98%. T_4 absorption is increased by fasting and decreased in malabsorption syndromes and by certain foods such as soybean infant formula. Dietary fiber decreases bioavailability of T_4 . Absorption may also decrease with age. In addition, many drugs and foods affect T_4 absorption (see PRECAUTIONS, Drug Interactions and Drug-Drug Interactions).

Distribution: Circulating thyroid hormones are greater than 99% bound to plasma proteins, including thyroxine-binding globulin (TBG), thyroxine-binding albumin (TBA), and albumin (A), whose capacities and affinities vary for each hormone. The higher affinity of both TBG and TBA for T_4 partially explains the higher serum levels of T_4 compared to T_3 . Protein-bound thyroid hormones are not available for tissue utilization. Only unbound hormone is metabolically active. Many drugs and physiologic conditions affect the binding of thyroid hormones to serum proteins (see PRECAUTIONS, Drug Interactions and Drug-Laboratory Test Interactions). Thyroid hormones do not readily cross the placental barrier (see PRECAUTIONS, Pregnancy).

Metabolism: T_4 is slowly eliminated (see Table 1). The major pathway of thyroid hormone metabolism is through sequential deiodination. Approximately 80% of circulating T_4 is derived from peripheral T_4 by monodeiodination. The liver is the major site of deiodination for both T_4 and T_3 , with T_4 deiodination also occurring at a number of additional sites, including the kidney and other tissues. Approximately 30% of the daily dose of T_4 is deiodinated to yield equal amounts of T_3 and reverse T_3 (rT_3), and rT_3 is further deiodinated to diiodotyrosine. Thyroid hormones are also metabolized via conjugation with glucuronides and sulfates and excreted chiefly into the bile and gut, where they undergo enterohepatic recirculation.

Elimination: Thyroid hormones are primarily excreted by the kidneys. A portion of the conjugated hormone reaches the colon unchanged and is eliminated in the feces. Approximately 20% of T_4 is eliminated in the stool. Urinary excretion of T_4 decreases with age.

Table 1: Pharmacokinetic Parameters of Thyroid Hormones in Euthyroid Patients

Thyroid Hormone	Rate in Thyroglobulin	Biologic Potency	$t_{1/2}$ (days)	Protein Binding (%)
Levothyroxine (T_4)	10 - 20	1	6 - 7	99.96
Liothyronine (T_3)	1	4	≤ 2	99.5

1 to 4 days in hyperthyroidism; 9 to 10 days in hypothyroidism. 1 includes TBG, TBA, and TBA.

INDICATIONS AND USAGE

Levothyroxine sodium is used for the following indications:

Hypothyroidism: As replacement or supplemental therapy in congenital or acquired hypothyroidism of any etiology, except transient hypothyroidism during the recovery phase of subacute thyroiditis. Specific indications include: primary (thyroidal), secondary (pituitary), and tertiary (hypothalamic) hypothyroidism and subclinical hypothyroidism. Primary hypothyroidism may result from functional deficiency, primary atrophy, partial or total congenital absence of the thyroid gland, or from the effects of surgery, resection, or drugs, with or without the presence of goiter.

Pituitary TSH Suppression: In the treatment or prevention of various types of euthyroid goiters (see WARNINGS and PRECAUTIONS), including thyroid nodules (see WARNINGS and PRECAUTIONS), subacute or chronic lymphocytic thyroiditis (Hashimoto's thyroiditis), multinodular goiter (see WARNINGS and PRECAUTIONS), and as an adjunct to surgery and radioiodine therapy in the management of thyrotropin-dependent well-differentiated thyroid cancer.

CONTRAINDICATIONS

Levothyroxine is contraindicated in patients with untreated subclinical (suppressed serum TSH level with normal T_3 and T_4 levels) or overt thyrotoxicosis of any etiology and in patients with acute myocardial infarction. Levothyroxine is contraindicated in patients with uncorrected adrenal insufficiency, since thyroid hormones may precipitate an acute adrenal crisis by increasing the metabolic clearance of glucocorticoids (see PRECAUTIONS, Levolet is contraindicated in patients with hypersensitivity to any of the inactive ingredients in Levolet tablets. (See DESCRIPTION, Inactive Ingredients.)

Pregnancy and Lactation: The extra iodide contained in Levolet may increase the risk of suppression of thyroid function in the fetus or nursing infant, particularly when the mother is already ingesting iodides from other sources. It is therefore recommended that non-iodine containing thyroid hormone preparations be used in pregnant women, women who may become pregnant, or nursing mothers. (See PRECAUTIONS, Pregnancy Category I)

WARNINGS

WARNINGS: Thyroid hormones, including Levolet, either alone or with other therapeutic agents, should not be used for the treatment of obesity or for weight loss. In euthyroid patients, doses within the range of daily hormonal requirements are ineffective for weight reduction. Larger doses may produce serious or even life threatening manifestations of toxicity, particularly when given in association with sympathomimetic amine such as those used for their anorectic effects.

Levothyroxine sodium should not be used in the treatment of male or female infertility unless this condition is associated with hypothyroidism.

In patients with nontoxic diffuse goiter or nodular thyroid disease, particularly the elderly or those with underlying cardiovascular disease, levothyroxine sodium therapy is contraindicated if the serum TSH level is already suppressed, due to the risk of precipitating overt thyrotoxicosis (see CONTRAINDICATIONS). If the serum TSH level is not suppressed, Levolet should be used with caution in conjunction with careful monitoring of thyroid function for evidence of hyperthyroidism and clinical monitoring for potential associated adverse cardiovascular signs and symptoms of hyperthyroidism.

In patients with nontoxic or nodular thyroid disease, particularly the elderly or those with underlying cardiovascular disease, Levolet should be used with caution in conjunction with careful monitoring of thyroid function for evidence of hyperthyroidism and clinical monitoring for potential associated adverse cardiovascular signs and symptoms of hyperthyroidism.

Levolet should also be used with caution in patients with euthyroid Graves' disease previously treated with antithyroid drugs, and with careful monitoring of thyroid function for evidence of hyperthyroidism.

PRECAUTIONS

General

Levothyroxine has a narrow therapeutic index. Regardless of the indication for use, careful dosage titration is necessary to avoid the consequences of over- or under-treatment. These consequences include, among others, effects on growth and development, cardiovascular function, bone metabolism, reproductive function, cognitive function, emotional state, gastrointestinal function, and on glucose and lipid metabolism. Many drugs interact with levothyroxine sodium, necessitating adjustments in dosing to maintain therapeutic response (see Drug Interactions).

Antibody production may increase with increased iodine intake in patients with Hashimoto's thyroiditis.

To maximize thyroid radioiodine uptake, Levolet should be temporarily withheld in patients scheduled to receive radioiodine diagnostically or therapeutically.

Patients with autonomous thyroid tissue: Exercise caution when administering Levolet to patients with autonomous thyroid tissue (e.g., nontoxic goiter or nodular thyroid disease) and to patients with euthyroid Graves' disease previously treated with antithyroid drugs in order to prevent precipitation of thyrotoxicosis (see WARNINGS).

Effects on bone mineral density in women: Long-term levothyroxine sodium therapy has been associated with increased bone resorption, thereby decreasing bone mineral density, especially in postmenopausal women or in women who are receiving subtherapeutic doses of levothyroxine sodium. The increased bone resorption may be associated with increased serum levels and urinary excretion of calcium and phosphorus, elevations in bone alkaline phosphatase and suppressed serum osteocalcin levels. Therefore, it is recommended that patients receive and maintain levothyroxine sodium at the minimum dose necessary to achieve the desired clinical and biochemical response.

Patients with underlying cardiovascular disease: Exercise caution when administering levothyroxine to patients with cardiovascular disorders and to the elderly in whom there is an increased risk of occult cardiac disease. In these patients, levothyroxine therapy should be initiated at lower doses than those recommended in younger individuals or in patients without cardiac disease (see WARNINGS, PRECAUTIONS, Geriatric Use, and DOSAGE AND ADMINISTRATION). If cardiac symptoms develop or worsen, the levothyroxine dose should be reduced or withheld for one week and then cautiously restarted at a lower dose. Over-treatment with levothyroxine sodium may have adverse cardiovascular effects such as an increase in heart rate, cardiac arrhythmias, and cardiac contractility and may precipitate angina or arrhythmias. Patients with coronary artery disease who are receiving levothyroxine therapy should be monitored closely during surgical procedures, since the possibility of precipitating cardiac arrhythmias may be greater in those treated with levothyroxine. Concurrent administration of levothyroxine and sympathomimetic agents to patients with coronary artery disease may precipitate coronary insufficiency.

Patients with nontoxic diffuse goiter or nodular thyroid disease: Exercise caution when administering levothyroxine to patients with nontoxic diffuse goiter or nodular thyroid disease in order to prevent precipitation of thyrotoxicosis (see WARNINGS), if the serum TSH is already suppressed. Levothyroxine sodium should not be administered (see CONTRAINDICATIONS).

Associated endocrine disorders

Hypoparathyroidism/hypocalcemia disorders: In patients with secondary or tertiary hypothyroidism, additional hypoparathyroidism/hypocalcemia deficiencies should be considered, and, if diagnosed, treated (see PRECAUTIONS, Autoimmune polyendocrine syndromes for adrenal insufficiency).

Autoimmune polyendocrine syndromes: Occasionally, chronic autoimmune thyroiditis may occur in association with other autoimmune disorders such as adrenal insufficiency, pernicious anemia, and insulin-dependent diabetes mellitus. Patients with autoimmune adrenal insufficiency should be treated with replacement glucocorticoids prior to initiation of treatment with levothyroxine sodium. Failure to do so may precipitate an acute adrenal crisis when thyroid hormone therapy is initiated, due to increased metabolic clearance of glucocorticoids by thyroid hormone. Patients with diabetes mellitus may require upward adjustments of their antidiabetic therapeutic regimens when treated with levothyroxine (see PRECAUTIONS, Drug Interactions).

Other associated medical conditions

Infants with congenital hypothyroidism appear to be at increased risk for other congenital anomalies, with cardiovascular anomalies (pulmonary stenosis, aortic aortic defect, and ventricular septal defect) being the most common association.

Secondary (pituitary) and tertiary (hypothalamic) hypothyroidism

Adequacy of therapy should be assessed by measuring serum free T_4 levels, which should be maintained in the upper half of the normal range in these patients.

Drug Interactions

Many drugs affect thyroid hormone pharmacokinetics and metabolism (e.g., absorption, synthesis, conversion, reabsorption, protein binding, and target tissue response) and may alter the therapeutic response to Levolet. In addition, thyroid hormone and thyroid status have varied effects on the pharmacokinetics and action of other drugs. A listing of drug-thyroid axis interactions is contained in Table 2.

The list of drug-thyroid axis interactions in Table 2 may not be comprehensive due to the introduction of new drugs that interact with the thyroid axis or the possibility of previously unknown interactions. The prescriber should be aware of this fact and should consult appropriate reference sources, (e.g., package inserts of newly approved drugs, medical literature) for additional information if a drug-drug interaction with levothyroxine is suspected.

Table 2: Drug-Thyroid Axis Interactions

Drug or Drug Class	Effect
Drugs that may reduce TSM secretion—the reduction is not sustained; therefore, hypothyroidism does not occur	
Doxazone / Doxamine Agonists Glucocorticoids Oestrogens	Use of these agents may result in a transient reduction in TSM secretion when administered at the following doses: Doxazone (2 mg/kg/day); Glucocorticoids (hydrocortisone ≥ 100 mg/day or equivalent); Oestrogens (≥ 100 mcg/day)
Drugs that alter thyroid hormone secretion	
Drugs that may decrease thyroid hormone secretion, which may result in hyperthyroidism	
Amiodarone Antituberculars Iodide (including iodine-containing radiographic contrast agents) Lithium Methimazole Propylthiouracil (PTU) Sulfonamides Toluamide	Long-term lithium therapy can result in goiter in up to 50% of patients, and either subclinical or overt hyperthyroidism, each in up to 20% of patients. The fetus, neonate, elderly and nursing patients with underlying thyroid disease (e.g., Hashimoto's thyroiditis or with Graves' disease previously treated with radioiodine or surgery) are among those individuals who are particularly susceptible to iodine-induced hyperthyroidism. Oral cholecystographic agents and amiodarone are slowly cleared, producing more prolonged hyperthyroidism than parenterally administered iodinated contrast agents. Long-term antitubercular therapy may minimally decrease T_4 and T_3 levels and increase TSM, although all values remain within normal limits in most patients.
Drugs that may increase thyroid hormone secretion, which may result in hyperthyroidism	
Amiodarone Iodide (including iodine-containing radiographic contrast agents)	Iodide and drugs that contain pharmacologic amounts of iodide may cause hyperthyroidism in euthyroid patients with Graves' disease previously treated with antithyroid drugs or in euthyroid patients with thyroid autonomy (e.g., multinodular goiter or hyperfunctioning thyroid adenoma). Hyperthyroidism may develop over several weeks and may persist for several months after therapy discontinuation. Amiodarone may induce hyperthyroidism by causing thyrotoxicosis.
Drugs that may decrease T_4 absorption, which may result in hypothyroidism	
Antacids - Aluminum & Magnesium Hydroxides - Simethicone Bile Acid Sequestrants - Cholestyramine - Colestipol Calcium Carbonate Cation Exchange Resins - Kayexalate Ferrous Sulfate Sucralfate	Concurrent use may reduce the efficacy of levothyroxine by binding and delaying or preventing absorption, potentially resulting in hypothyroidism. Calcium carbonate may form an insoluble chelate with levothyroxine, and ferrous sulfate likely forms a ternary-thyroxine complex. Administer levothyroxine at least 4 hours apart from these agents.
Drugs that may alter T_4 and T_3 serum transport—but FT_4 concentration remains normal, and, therefore, the patient remains euthyroid	
Drugs that may increase serum TBG concentration	
Colchicine Estrogen-containing oral contraceptives Estrogens (oral) Marfan / Methadone 5-Fluorouracil Mitotane Tyrosinase	Androgens / Anabolic Steroids Asparaginase Glucocorticoids Slow-Release Nicotinic Acid
Drugs that may cause protein-binding site displacement	
Furosemide (≥ 30 mg IV) Mazone Methadone Non Steroidal Anti-inflammatory Drugs - Fenamate - Phenytoin Sulicylates (≥ 2 g/day)	Administration of these agents with levothyroxine results in an initial transient increase in FT_4 . Continued administration results in a decrease in serum T_4 and normal FT_4 and TSM concentrations and, therefore, patients are clinically euthyroid. Sulicylates inhibit binding of T_4 and T_3 to TBG and transthyretin. An initial increase in serum FT_4 is followed by return of FT_4 to normal levels with sustained therapeutic serum salicylate concentrations, although total T_4 levels may decrease by as much as 30%.
Drugs that may alter T_4 and T_3 metabolism	
Drugs that may increase hepatic metabolism, which may result in hypothyroidism	
Carbamazepine Hydantoin Phenytoin Rifampin	Concomitant hepatic microsomal drug-metabolizing enzyme activity may cause increased hepatic degradation of levothyroxine resulting in increased levothyroxine requirements. Phenytoin and carbamazepine reduce serum protein binding of levothyroxine, and total- and free- T_4 may be reduced by 20% to 40%, but most patients have normal serum TSM levels and are clinically euthyroid.
Drugs that may decrease T_4 5'-deiodinase activity	
Antituberculars Bile-adsorbent resins - (e.g., Propylthiouracil ≥ 150 mg/day) Glucocorticoids - (e.g., Dexamethasone 2 mg/day) Propylthiouracil (PTU)	Administration of these enzyme inducers decreases the peripheral conversion of T_4 to T_3 , leading to decreased T_3 levels. However, serum T_4 levels are usually normal but may occasionally be slightly increased. In patients treated with large doses of propylthiouracil (> 100 mg/day), T_3 and T_4 levels change slightly, TSM levels remain normal, and patients are clinically euthyroid. It should be noted that actions of antitubercular agents may be impaired when the hypothyroid patient is converted to the euthyroid state. Short-term administration of large doses of glucocorticoids may decrease serum T_4 concentrations by 30% with minimal change in serum T_3 levels. However, long-term glucocorticoid therapy may result in slightly decreased T_3 and T_4 levels due to decreased TSG production (see above).
Miscellaneous	
Anticoagulants (oral) - Coumatin Derivatives - Indandione Derivatives	Thyroid hormones appear to increase the catabolism of vitamin K-dependent clotting factors, thereby increasing the anticoagulant activity of oral anticoagulants. Concomitant use of these agents impairs the compensatory increases in clotting factor synthesis. Prothrombin time should be carefully monitored in patients taking levothyroxine and oral anticoagulants, and the dose of anticoagulant therapy adjusted accordingly.

LEVOLET®
(LEVOTHYROXINE
SODIUM TABLETS,
USP)
Rx only

Antidepressants • Tricyclics (e.g., Amitriptyline) • Anticholinergics (e.g., Atropine) • Selective Serotonin Reuptake Inhibitors (SSRIs; e.g., Sertraline)	Concurrent use of tricyclic antidepressants and levothyroxine may increase the therapeutic and toxic effects of both drugs possibly due to increased receptor sensitivity to catecholamines. Toxic effects may include increased risk of cardiac arrhythmias and CNS stimulation, onset of action of tricyclics may be accelerated. Administration of sertraline in patients stabilized on levothyroxine may result in increased levothyroxine requirements.
Antidiabetic Agents • Biguanides • Sulfonylureas • Thiazolidinediones • Insulin	Addition of levothyroxine to antidiabetic or insulin therapy may result in increased antidiabetic agent or insulin requirements. Careful monitoring of glucose control is recommended, especially when thyroid therapy is started, changed, or discontinued.
Cardiac Glycosides	Serum digoxin glycoside levels may be reduced in hyperthyroidism or when the hyperthyroid patient is converted to the euthyroid state. Therapeutic effect of cardiac glycosides may be reduced.
Cytokines • Interferon- α • Interleukin-2	Therapy with interferon- α has been associated with the development of antilevothyroxine antibodies in 20% of patients and some have transient hypothyroidism, hyperthyroidism, or both. Patients who have antilevothyroxine antibodies before treatment are at higher risk for thyroid dysfunction during treatment. Interleukin-2 has been associated with transient hypothyroidism in 22% of patients. Interferon- β and - γ have not been reported to cause thyroid dysfunction.
Growth Hormones • Somatotropin • Somatropin	Excessive use of thyroid hormones with growth hormones may accelerate epiphyseal closure. However, untreated hypothyroidism may interfere with growth response to growth hormone.
Levamisole	Concurrent use may produce marked hyperthyroidism and tachycardia; cautious administration to patients receiving thyroid hormone therapy is recommended.
Methylnamine Bronchodilators (e.g., Theophylline)	Decreased methylnamine clearance may occur in hypothyroid patients; clearance returns to normal when the euthyroid state is achieved.
Radioactive Agents	Thyroid hormones may reduce the uptake of ^{123}I , ^{131}I , and $^{99\text{m}}\text{Tc}$.
Sympathomimetics	Concurrent use may increase the effects of sympathomimetics or thyroid hormone. Thyroid hormones may increase the risk of coronary insufficiency when sympathomimetic agents are administered to patients with coronary artery disease.
Chloral Hydrate Diazepam Ethinamate Levamisole Levamisole Metoclopramide N-Mercaptopurine Nitroprusside Para-aminosalicylic acid sodium Perphenazine Rasorcol (excessive topical use) Thiazide Diuretics	These agents have been associated with thyroid hormone and/or TSH level alterations by various mechanisms.

Drug-Drug Interactions: Levothyroxine increases the response to oral anticoagulant therapy. Therefore, a decrease in the dose of anticoagulant may be warranted with correction of the hypothyroid state or when the Levolet dose is increased. Prothrombin time should be closely monitored to permit appropriate and timely dosage adjustments (see Table 2).

Digoxin Interactions: The therapeutic effects of digoxin glycosides may be reduced by levothyroxine. Serum digoxin glycoside levels may be decreased when a hypothyroid patient becomes euthyroid, necessitating an increase in the dose of digoxin glycosides (see Table 2).

Drug-Food Interactions: Consumption of certain foods may affect levothyroxine absorption, thereby necessitating adjustments in dosing. Soybean flour (infant formula), cotton seed meal, walnuts, and dietary fiber may bind and decrease the absorption of levothyroxine sodium from the GI tract.

Drug-Laboratory Test Interactions: Changes in TBG concentration must be considered when interpreting T_4 and T_3 values, which necessitates measurement and evaluation of unbound (free) hormone and/or determination of the free- T_4 index (FT $_4$). Pregnancy, infectious hepatitis, estrogens, estrogen-containing oral contraceptives, and acute intermittent porphyria increase TBG concentrations. Decreases in TBG concentrations are observed in nephrosis, severe hypoproteinemia, severe liver disease, scleroderma, and after androgen or corticosteroid therapy (see also Table 2). Fetal hyper- or hypothyroidism-binding globulins have been described, with the incidence of TBG deficiency approximating 1 in 3000.

Carcinogenesis, Mutagenesis, and Impairment of Fertility: Animal studies have not been performed to evaluate the carcinogenic potential, mutagenic potential or effects on fertility of levothyroxine. The synthetic T_4 in Levolet is identical to that produced naturally by the human thyroid gland. Although there has been a reported association between prolonged thyroid hormone therapy and breast cancer, this has not been confirmed. Patients receiving Levolet for appropriate clinical indications should be treated to the lowest effective replacement dose.

Pregnancy Category X. See CONTRAINDICATIONS: Levolet may cause fetal harm when administered to a pregnant woman. Non-iodine containing thyroid hormone preparations are recommended for use in pregnant women, women who may become pregnant, or nursing mothers. Published literature indicates that potassium iodide (25 mg/kg/day [150 mg/day]) in the pregnant rat results in fetal malformations (ventricular septal defects, high aortic arch, aberrant subclavian arteries, incomplete lung, amphibole and growth retardation), representing a TBG multiple of the potassium iodide exposure in Levolet based on surface area comparison (mg/m 2). No known no effect level has been established. Higher doses (100 mg/kg/day; 600 mg/m 2 - 3000 mg human exposure) resulted in fetolethality (mortality). If the patient becomes pregnant while taking this drug, the patient should be switched to a non-iodine containing thyroid preparation and be apprised of the potential hazard to the fetus.

Hypothyroidism during pregnancy is associated with a higher rate of complications, including spontaneous abortion, pre-eclampsia, edema and premature delivery. Maternal hypothyroidism may have an adverse effect on fetal and childhood growth and development. Hypothyroid pregnant patients should be treated only with non-iodine containing thyroid hormone products. Serum T $_4$ levels should be monitored every trimester and 6-8 weeks postpartum.

Thyroid hormones cross the placental barrier as evidenced by levels in cord blood of diuretic fetuses being approximately one-third maternal level. Transfer of thyroid hormone from the mother to the fetus, however, may not be adequate to prevent a clear hypothyroidism.

Lactation and Nursing Mothers: There are insufficient data from the published literature to establish the safety of Levolet in nursing mothers. The potassium iodide component in Levolet may increase the risk of thyroid dysfunction suppression in nursing infants, particularly when the nursing mother is ingesting iodine from other sources. It is recommended that a non-iodine containing thyroid hormone preparation be used (see WARNINGS).

Pediatric Use

General

The goal of treatment in pediatric patients with hypothyroidism is to achieve and maintain normal intellectual and physical growth and development. The initial dose of levothyroxine varies with age and body weight (see DOSAGE AND ADMINISTRATION, Table 3). Dosing adjustments are based on an assessment of the individual patient's clinical and laboratory parameters (see WARNINGS and PRECAUTIONS, Laboratory Tests).

In children in whom a diagnosis of permanent hypothyroidism has not been established, it is recommended that levothyroxine administration be discontinued for a 30-day trial period. Only after the child is at least 3 years of age, serum T_4 and TSH levels should then be obtained. If the T_4 is low and the TSH high, the diagnosis of permanent hypothyroidism is established, and levothyroxine therapy should be reinstituted. If the T_4 and TSH levels are normal, euthyroidism may be assumed and, therefore, the hypothyroidism can be considered to have been transient. In this instance, the physician should carefully monitor the child and repeat the thyroid function tests if any signs of symptoms of hypothyroidism develop. In this setting, the clinician should have a high index of suspicion of relapse. If the results of the levothyroxine withdrawal test are inconclusive, careful follow-up and subsequent testing will be necessary.

Since some more severely affected children may become clinically hypothyroid when treatment is discontinued for 30 days, an alternate approach is to reduce the replacement dose of levothyroxine by half during the 30-day trial period. If, after 30 days, the serum TSH is elevated above 20 mIU/L, the diagnosis of permanent hypothyroidism is confirmed, and full replacement therapy should be resumed. However, if the serum TSH has not risen to greater than 20 mIU/L, levothyroxine treatment should be discontinued for another 30-day trial period followed by repeat serum T_4 and TSH testing.

The presence of concomitant medical conditions should be considered in certain clinical circumstances and, if present, appropriately treated (see PRECAUTIONS).

Congenital Hypothyroidism (see WARNINGS, PRECAUTIONS, Laboratory Tests, and DOSAGE AND ADMINISTRATION)

Rapid restoration of normal serum T_4 concentrations is essential for preventing the adverse effects of congenital hypothyroidism on intellectual development.

DOSAGE AND ADMINISTRATION

General Principles

The goal of replacement therapy is to achieve and maintain a clinical and biochemical euthyroid state. The goal of suppressive therapy is to inhibit growth and/or function of abnormal thyroid tissue. The dose of Levolet that is adequate to achieve these goals depends on a variety of factors including the patient's age, body weight, cardiovascular status, concomitant medical conditions, concomitant medications, and the specific nature of the condition being treated (see WARNINGS and PRECAUTIONS). Hence, the following recommendations serve only as starting guidelines. Dosing must be individualized and adjustments must be based on periodic assessment of the patient's clinical response and laboratory parameters (see PRECAUTIONS, Laboratory Tests).

Levolet should be taken in the morning on an empty stomach, at least one-half hour before any food is eaten. Levolet should be taken at least 4 hours apart from drugs that are known to interfere with its absorption (see PRECAUTIONS, Drug Interactions).

Due to the long half-life of levothyroxine, the peak therapeutic effect of a given dose of levothyroxine sodium may not be attained for 4-8 weeks.

Caution should be exercised when administering Levolet to patients with underlying cardiovascular disease, to the elderly, and to those with concomitant agents insufficiency (see PRECAUTIONS).

Specific Patient Populations

Hyperthyroidism: Levolet may be used in children in whom Growth and Puberty are Complete (see WARNINGS and PRECAUTIONS, Laboratory Tests). Therapy may begin at full replacement doses in otherwise healthy individuals less than 50 years of age and in those older than 50 years who have recently treated for hyperthyroidism or who have had hyperthyroidism for only a short time (such as a few months). The average full replacement dose of levothyroxine sodium is approximately 1.7 mcg/kg/day (e.g., 100-125 mcg/day for a 70 kg adult). Older patients may require less than 1 mcg/kg/day. Levothyroxine sodium doses greater than 200 mcg/day are seldom required. An inadequate response to daily doses \geq 300 mcg/day is rare and may indicate poor compliance, malabsorption, and/or drug interactions.

For most patients older than 50 years or for patients under 50 years of age with underlying cardiac disease, an initial starting dose of 25-50 mcg/day of levothyroxine sodium is recommended, with gradual increments in dose at 6-8 week intervals, as needed. The recommended starting dose of levothyroxine sodium in elderly patients with cardiac disease is 12.5-25 mcg/day, with gradual dose increments at 4-6 week intervals. The levothyroxine sodium dose is generally adjusted in 12.5-25 mcg increments until the patient with primary hypothyroidism is clinically euthyroid and the serum TSH has normalized.

In patients with severe hypothyroidism, the recommended initial levothyroxine sodium dose is 12.5-25 mcg/day with increases of 25 mcg/day every 2-4 weeks, accompanied by clinical and laboratory assessment, until the TSH level is normalized.

In patients with secondary (pituitary) or tertiary (hypothalamic) hypothyroidism, the levothyroxine sodium dose should be titrated until the patient is clinically euthyroid and the serum free- T_4 level is restored to the upper half of the normal range.

Pediatric Dosage: Congenital or Acquired Hypothyroidism (see WARNINGS and PRECAUTIONS, Laboratory Tests)

General Principles

In general, levothyroxine therapy should be instituted at full replacement doses as soon as possible. Delays in diagnosis and institution of therapy may have deleterious effects on the child's intellectual and physical growth and development.

Under-treatment and over-treatment should be avoided (see WARNINGS and PRECAUTIONS, Pediatric Use).

Levolet may be administered to infants and children who cannot swallow intact tablets by crushing the tablet and suspending the freshly crushed tablet in a small amount (5-10 mL) of 1:2 teaspoonful of water. This suspension can be administered by spoon or dropper. DO NOT STORE THE SUSPENSION. Foods that decrease absorption of levothyroxine, such as soybean infant formula, should not be used for administering levothyroxine sodium tablets (see PRECAUTIONS, Drug-Food Interactions).

Newborns

The recommended starting dose of levothyroxine sodium in newborn infants is 10-15 mcg/kg/day. A lower starting dose (e.g., 25 mcg/day) should be considered in infants at risk for cardiac failure, and the dose should be increased in 4-6 weeks as needed, based on clinical and laboratory response to treatment. In infants with very low (e.g., 5 mcg/dL) or undetectable serum T_4 concentrations, the recommended initial starting dose is 50 mcg/kg/day of levothyroxine sodium.

Infants and Children

Levothyroxine therapy is usually initiated at full replacement doses, with the recommended dose per body weight decreasing with age (see Table 3). However, in children with chronic or severe hypothyroidism, an initial dose of 25 mcg/day of levothyroxine sodium is recommended, with increments of 25 mcg every 2-4 weeks until the desired effect is achieved.

Hypothyroidism in an older child can be minimized if the starting dose is one-fourth of the recommended full replacement dose, and the dose is then increased on a weekly basis by an amount equal to one-fourth the full-recommended replacement dose until the full recommended replacement dose is reached.

Table 3. Levothyroxine Sodium Dosing Guidelines for Pediatric Hypothyroidism

AGE	Daily Dose Per Kg Body Weight
0-3 months	10-15 mcg/kg/day
3-6 months	8-10 mcg/kg/day
6-12 months	6-8 mcg/kg/day
1-5 years	5-6 mcg/kg/day
6-12 years	4-5 mcg/kg/day
\geq 12 years but growth and puberty incomplete	2-3 mcg/kg/day
Growth and puberty complete	1.7 mcg/kg/day

* The dose should be adjusted based on clinical response and laboratory parameters (see PRECAUTIONS, Laboratory Tests and Pediatric Use)

Pregnancy: Levolet should not be used in women who are pregnant, plan to become pregnant, or are nursing. Levolet should be substituted with a non-iodine containing thyroid hormone preparation (see CONTRAINDICATIONS and PRECAUTIONS).

Subclinical Hypothyroidism: If this condition is treated, a lower levothyroxine sodium dose (e.g., 1 mcg/kg/day) than that used for full replacement may be adequate to normalize the serum TSH level. Patients who are not treated should be monitored yearly for changes in clinical status and thyroid laboratory parameters.

TSH Suppression in Well-differentiated Thyroid Cancer and Thyroid Nodules: The target level for TSH suppression in these conditions has not been established with controlled studies in action on the efficacy of TSH suppression for benign nodular disease in children. Therefore, the dose of Levolet used for TSH suppression should be individualized based on the specific disease and the patient being treated. In the treatment of well-differentiated (papillary and follicular) thyroid cancer, levothyroxine is used as an adjunct to surgery and radioactive iodine therapy. Generally, TSH is suppressed to $<$ 0.1 mIU/L, and this usually requires a levothyroxine sodium dose of greater than 2 mcg/kg/day. However, in patients with high-risk tumors, the target level for TSH suppression may be $<$ 0.01 mIU/L. In the treatment of benign nodules and nodules of uncertain nature, TSH is generally suppressed to a higher target (e.g., 0.1 mIU/L to lower 0.5 or 1.0 mIU/L) than that used for the treatment of thyroid cancer. Levothyroxine sodium is contraindicated if the serum TSH is not suppressed due to the risk of precipitating overt hyperthyroidism (see CONTRAINDICATIONS, WARNINGS and PRECAUTIONS).

Myxedema Coma: Myxedema coma is a life-threatening emergency characterized by poor circulation and hypometabolism, and may result in unpredictable absorption of levothyroxine sodium from the gastrointestinal tract. Therefore, oral thyroid hormone drug products are not recommended to treat this condition. Thyroid hormone products formulated for intravenous administration should be utilized.

HOW SUPPLIED

LEVOLET[®] (levothyroxine sodium tablets, USP) are round color-coded tablet embossed with a V and 10 number.

STRENGTH (mcg)	COLOR	EMBOSING	SIZE	SIZE
25 mcg	White	3111/V	Bottles of 100	Bottles of 1000
50 mcg	White	3112/V	Bottles of 100	Bottles of 1000
75 mcg	White/Purple	3113/V	Bottles of 100	Bottles of 1000
100 mcg	White	3114/V	Bottles of 100	Bottles of 1000
125 mcg	Yellow	3115/V	Bottles of 100	Bottles of 1000
150 mcg	Orange	3116/V	Bottles of 100	Bottles of 1000
175 mcg	Brown	3117/V	Bottles of 100	Bottles of 1000
200 mcg	Dark Blue	3122/V	Bottles of 100	Bottles of 1000
225 mcg	Light Blue	3123/V	Bottles of 100	Bottles of 1000
250 mcg	Pink	3124/V	Bottles of 100	Bottles of 1000
300 mcg	Green	3127/V	Bottles of 100	Bottles of 1000

Page 2

and evaluation of unbound (free) hormone and/or determination of the free T_4 index (FTI). Pregnancy infections, hepatitis, estrogens, estrogen-containing oral contraceptives, and acute intermittent porphyria increase T₄ concentrations. Decreases in T₄ concentrations are observed in nephrosis, severe hypoparathyroidism, severe liver disease, acromegaly, and after ablation of corticosteroid therapy (see also Table 2). Paradoxical hypothyroidism or hypothyroidism during glucocorticoid therapy has been described, with the incidence of T₄ deficiency approximating 1 in 1000.

Carcinogenesis, Mutagenesis, and Impairment of Fertility: Animal studies have not been performed to evaluate the carcinogenic potential, mutagenic potential or effects on fertility of levothyroxine. The synthetic T_4 in Levolet is identical to that produced naturally by the human thyroid gland. Although there has been a reported association between prolonged thyroid hormone therapy and breast cancer, this has not been confirmed. Patients receiving Levolet for appropriate clinical indications should be treated in the lowest effective replacement dose.

Pregnancy Category X: See **CONTRAINDICATIONS:** Levolet may cause fetal harm when administered to a pregnant woman. Non-iodine containing thyroid hormone preparations are recommended for use in pregnant women, women who may become pregnant, or nursing mothers. Published literature indicates that potassium iodide 25 mg/day (25 mg/m²) in the pregnant rat results in fetal malformations (ventricular septal defects, cleft palate arch, abnormal skeletal ossification, incomplete lung, omphalocele and growth retardation), representing a TSD of 200 mg/m² of the potassium iodide dose in Levolet based on surface area comparison (mg/m²). No known no effect level has been established. Higher doses (100 mg/kg/day, 600 mg/m², 3000 mg human equivalent) resulted in fetal/maternal mortality. If the patient becomes pregnant while taking this drug, the patient should be switched to a non-iodine containing thyroid preparation and be apprised of the potential hazard to the fetus.

Hypothyroidism during pregnancy: is associated with a higher rate of complications, including spontaneous abortion, pre-eclampsia, stillbirth and premature delivery. Maternal hypothyroidism may have an adverse effect on fetal and childhood growth and development. Hypothyroid pregnant patients should be treated only with non-iodine containing thyroid hormone products. Serum TSH levels should be monitored every trimester and 6-8 weeks postpartum.

Thyroid hormones cross the placental barrier to some extent as evidenced by levels in cord blood of atrophic fetuses being approximately one-third maternal levels. Transfer of thyroid hormone from the mother to the fetus, however, may not be adequate to prevent in utero hypothyroidism.

Lactation and Nursing Mothers: There are insufficient data from the published literature to establish the safety of Levolet in nursing mothers. The potassium iodide contained in Levolet may increase the risk of thyroid function suppression in nursing infants, particularly when the nursing mother is ingesting iodides from other sources. It is recommended that a non-iodine containing thyroid hormone preparation be used (see **WARNINGS**).

Pediatric Use

INDICATIONS

The goal of treatment in pediatric patients with hypothyroidism is to achieve and maintain normal intellectual and physical growth and development.

The initial dose of levothyroxine varies with age and body weight (see **DOSE AND ADMINISTRATION, Table 3**). Dosing adjustments are based on an assessment of the individual patient's clinical and laboratory parameters (see **WARNINGS** and **PRECAUTIONS, Laboratory Tests**).

In children in whom a diagnosis of permanent hypothyroidism has not been established, it is recommended that levothyroxine administration be discontinued for a 30-day trial period, but only after the child is at least 3 years of age. Serum T_4 and TSH levels should then be obtained. If the T_4 is low and the TSH high, the diagnosis of permanent hypothyroidism is established, and levothyroxine therapy should be reinitiated. If the T_4 and TSH levels are normal, euthyroidism may be assumed and, therefore, the hypothyroidism can be considered to have been transient. In this instance, however, the physician should carefully monitor the child and repeat the thyroid function tests if any signs or symptoms of hypothyroidism develop. In this setting, the clinician should have a high index of suspicion of relapse. If the results of the levothyroxine withdrawal test are inconclusive, careful follow-up and subsequent testing will be necessary.

Since some more severely affected children may become clinically hypothyroid when treatment is discontinued for 30 days, an alternate approach to reduce the replacement dose of levothyroxine by half during the 30-day trial period. If, after 30 days, the serum TSH is elevated above 20 mU/L, the diagnosis of permanent hypothyroidism is confirmed, and full replacement therapy should be resumed. However, if the serum TSH has not risen to greater than 20 mU/L, levothyroxine treatment should be discontinued for another 30-day trial period followed by repeat serum T_4 and TSH testing.

The presence of concomitant medical conditions should be considered in certain clinical circumstances and, if present, appropriately treated (see **PRECAUTIONS**).

Congenital Hypothyroidism (see **WARNINGS, PRECAUTIONS, Laboratory Tests, and DOSE AND ADMINISTRATION)**

Rapid restoration of normal serum T_4 concentrations is essential for preventing the adverse effects of congenital hypothyroidism on intellectual development as well as on overall physical growth and maturation. Therefore, Levolet therapy should be initiated immediately upon diagnosis and is generally continued for life.

During the first 2 weeks of Levolet therapy, infants should be closely monitored for cardiac overload, arrhythmias, and aspiration from avid suckling.

The patient should be monitored closely to avoid undertreatment or overtreatment. Undertreatment may have deleterious effects on intellectual development and linear growth. Overtreatment has been associated with craniosynostosis in infants, and may adversely affect the tempo of brain maturation and accelerate the bone age, with resultant premature closure of the epiphyses and compromised adult stature.

Acquired Hypothyroidism in Pediatric Patients

The patient should be monitored closely to avoid undertreatment and overtreatment. Undertreatment may result in poor school performance due to impaired concentration and slowed maturation, and in reduced adult height. Overtreatment may accelerate the bone age and result in premature epiphyseal closure and compromised adult stature.

Treated children may manifest a period of catch-up growth, which may be adequate in some cases to normalize adult height in children with severe or prolonged hypothyroidism; catch-up growth may not be adequate to normalize adult height.

Geriatric Use

Because of the increased prevalence of cardiovascular disease among the elderly, levothyroxine therapy should not be initiated at the full replacement dose (see **WARNINGS, PRECAUTIONS, and DOSE AND ADMINISTRATION**).

ADVERSE REACTIONS

Adverse reactions associated with levothyroxine therapy are primarily those of hyperthyroidism due to therapeutic overdosage (see **PRECAUTIONS** and **OVERDOSAGE**). They include the following:

- General:** fatigue, increased appetite, weight loss, heat intolerance, fever, excessive sweating;
- Central nervous system:** headache, hyperactivity, nervousness, anxiety, irritability, emotional lability, insomnia;
- Musculoskeletal:** tremors, muscle weakness;
- Cardiovascular:** palpitations, tachycardia, arrhythmias, increased pulse and blood pressure, heart failure, angina, myocardial infarction, cardiac arrest;
- Respiratory:** dyspnea;
- Gastrointestinal:** diarrhea, vomiting, abdominal cramps and nausea in liver function tests;
- Dermatologic:** hair loss, flushing;
- Endocrine:** decreased bone mineral density;
- Reproductive:** menstrual irregularities, impaired fertility.

Pseudotumor cerebri and slipped capital femoral epiphysis have been reported in children receiving levothyroxine therapy. Overtreatment may result in craniosynostosis in infants and premature closure of the epiphyses in children, with resultant compromised height.

Seizures have been reported rarely with the institution of levothyroxine therapy.

Infants on levothyroxine dosage will produce or fail to ameliorate the signs and symptoms of hypothyroidism.

Hypersensitivity reactions to inactive ingredients have occurred in patients treated with thyroid hormone products. These include urticaria, pruritus, skin rash, flushing, angioedema, various GI symptoms (abdominal pain, nausea, vomiting and diarrhea), fever, a stridor, serum sickness and wheezing. Hypersensitivity to levothyroxine itself is not known to occur.

OVERDOSAGE

The signs and symptoms of overdosage are those of hyperthyroidism (see **PRECAUTIONS** and **ADVERSE REACTIONS**). In addition, confusion and disorientation may occur. Cerebral embolism, shock, coma, and death have been reported. Seizures have occurred in a child ingesting 18 mg of levothyroxine. Symptoms may not necessarily be evident or may not appear until several days after ingestion of levothyroxine sodium.

Treatment of Overdosage

Levothyroxine sodium should be reduced in dose or temporarily discontinued if signs or symptoms of overdosage occur.

Acute Massive Overdosage: This may be a life-threatening emergency; therefore, symptomatic and supportive therapy should be instituted immediately. If not contraindicated (e.g., by sodium, coma, or loss of the gag reflex), the stomach should be emptied by means of gastric lavage to decrease gastrointestinal absorption. Activated charcoal or cholestyramine may also be used to decrease absorption. Central and peripheral increased sympathetic activity may be treated by administering β -receptor antagonists, e.g., propranolol, provided there are no medical contraindications to their use. Provide respiratory support as needed; control congestive heart failure and arrhythmias; control fever, hyperglycemia, and fluid loss as necessary. Large doses of antithyroid drugs (e.g., methimazole or propylthiouracil) followed in one to two hours by large doses of iodine may be given to inhibit synthesis and release of thyroid hormones. Glucocorticoids may be given to inhibit the conversion of T_4 to T_3 . Plasmapheresis, charcoal hemoperfusion and exchange transfusion have been reserved for cases in which continued clinical deterioration occurs despite conventional therapy. Because T_4 is highly protein bound, very little drug will be removed by dialysis.

3-6 months	8-8 mcg/kg/day
6-12 months	8-8 mcg/kg/day
1-5 years	5-6 mcg/kg/day
6-12 years	4-5 mcg/kg/day
>12 years but growth and puberty incomplete	2-3 mcg/kg/day
Adolescents and adults	1-1.7 mcg/kg/day

The dose should be adjusted based on clinical response and laboratory parameters (see **PRECAUTIONS, Laboratory Tests** and **Pediatric Use**).

Warnings: Levolet should not be used in women who are pregnant, plan to become pregnant, or are nursing. Levolet should be substituted with a non-iodine containing thyroid hormone preparation (see **CONTRAINDICATIONS** and **PRECAUTIONS**).

Subclinical Hypothyroidism: If this condition is treated, a lower levothyroxine sodium dose (e.g., 1 mcg/kg/day) than that used for full replacement may be adequate to normalize the serum TSH level. Patients who are not treated should be monitored yearly for changes in clinical status and thyroid laboratory parameters.

TSH Suppression in Well-differentiated Thyroid Cancer and Thyroid Nodules: The target level for TSH suppression in these conditions has not been established with controlled studies. In addition, the efficacy of TSH suppression for benign nodular disease is controversial. Therefore, the dose of Levolet used for TSH suppression should be individualized based on the specific disease and the patient being treated. In the treatment of well-differentiated (papillary and follicular) thyroid cancer, levothyroxine is used as an adjunct to surgery and radioactive iodine therapy. Generally, TSH is suppressed to <0.1 mU/L, and this usually requires a levothyroxine sodium dose of greater than 2 mcg/kg/day. However, in patients with high-risk tumors, the target level for TSH suppression may be <0.01 mU/L. In the treatment of benign nodules and nontoxic multinodular goiter, TSH is generally suppressed to a higher target (e.g., 0.1 mU/L to either 0.5 or 1.0 mU/L) than that used for the treatment of thyroid cancer. Levothyroxine sodium is contraindicated if the serum TSH is already suppressed due to the risk of precipitating overt thyrotoxicosis (see **CONTRAINDICATIONS, WARNINGS** and **PRECAUTIONS**).

Myxedema Coma: Myxedema coma is a life-threatening emergency characterized by poor circulation and hypometabolism, and may result in unpredictable absorption of levothyroxine sodium from the gastrointestinal tract. Therefore, oral thyroid hormone drug products are not recommended to treat this condition. Thyroid hormone products formulated for intravenous administration should be utilized.

HOW SUPPLIED

LEVOLET® (levothyroxine sodium tablets, USP) are round color-coded tablet embossed with a V and ID number.

STRENGTH (mcg)	COLOR	EMBOSSING	SIZE	SIZE
25 mcg	Peach	3911/V	Bottles of 100	Bottles of 1000
50 mcg	White	3912/V	Bottles of 100	Bottles of 1000
75 mcg	Blue-White	3913/V	Bottles of 100	Bottles of 1000
88 mcg	Olive	3912/V	Bottles of 100	Bottles of 1000
100 mcg	Yellow	3914/V	Bottles of 100	Bottles of 1000
112 mcg	Pink	3915/V	Bottles of 100	Bottles of 1000
125 mcg	Brown	3916/V	Bottles of 100	Bottles of 1000
137 mcg	Dark Blue	3922/V	Bottles of 100	Bottles of 1000
150 mcg	Purple	3915/V	Bottles of 100	Bottles of 1000
175 mcg	Light Blue	3920/V	Bottles of 100	Bottles of 1000
200 mcg	Pink	3918/V	Bottles of 100	Bottles of 1000
300 mcg	Green	3917/V	Bottles of 100	Bottles of 1000

STORAGE CONDITIONS

Store between 20° and 25°C (68°-77°F) with excursions permitted from 15° to 30°C (59°-86°F).

Rx only

Manufactured by:
VINTAGE PHARMACEUTICALS, INC.
Charlotte, NC 28206

IN-181/180577
Rev 12/02
RI

Pg 4