

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

208313Orig1s000

LABELING

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use INFUGEM safely and effectively. See full prescribing information for INFUGEM.

INFUGEM (gemcitabine in sodium chloride injection), for intravenous use
Initial U.S. Approval: 1996

INDICATIONS AND USAGE

INFUGEM is a nucleoside metabolic inhibitor indicated:

- in combination with carboplatin, for the treatment of advanced ovarian cancer that has relapsed at least 6 months after completion of platinum-based therapy. (1.1)
- in combination with paclitaxel, for first-line treatment of metastatic breast cancer after failure of prior anthracycline-containing adjuvant chemotherapy, unless anthracyclines were clinically contraindicated. (1.2)
- in combination with cisplatin for the treatment of non-small cell lung cancer. (1.3)
- as a single agent for the treatment of pancreatic cancer. (1.4)

DOSAGE AND ADMINISTRATION

INFUGEM is for intravenous infusion only.

- Ovarian Cancer: 1000 mg/m² over 30 minutes on Days 1 and 8 of each 21-day cycle. (2.1)
- Breast Cancer: 1250 mg/m² over 30 minutes on Days 1 and 8 of each 21-day cycle. (2.2)
- Non-Small Cell Lung Cancer: 1000 mg/m² over 30 minutes on Days 1, 8, and 15 of each 28-day cycle or 1250 mg/m² over 30 minutes on Days 1 and 8 of each 21-day cycle. (2.3)
- Pancreatic Cancer: 1000 mg/m² over 30 minutes once weekly for the first 7 weeks, then one week rest, then once weekly for 3 weeks of each 28-day cycle. (2.4)

DOSAGE FORMS AND STRENGTHS

Injection: single-dose premixed infusion bags containing 10 mg/mL of gemcitabine in 0.9% sodium chloride:

- 1200 mg in 120 mL (3)
- 1300 mg in 130 mL (3)
- 1400 mg in 140 mL (3)
- 1500 mg in 150 mL (3)
- 1600 mg in 160 mL (3)
- 1700 mg in 170 mL (3)
- 1800 mg in 180 mL (3)
- 1900 mg in 190 mL (3)

- 2000 mg in 200 mL (3)
- 2200 mg in 220 mL (3)

CONTRAINDICATIONS

Patients with a known hypersensitivity to gemcitabine (4)

WARNINGS AND PRECAUTIONS

- Schedule-Dependent Toxicity: Increased toxicity with infusion time greater than 60 minutes or dosing more frequently than once weekly. (5.1)
- Myelosuppression: Monitor for myelosuppression prior to each cycle and reduce or withhold dose for severe myelosuppression. (5.2, 5.7)
- Pulmonary Toxicity and Respiratory Failure: Discontinue INFUGEM immediately for unexplained new or worsening dyspnea or evidence of severe pulmonary toxicity. (5.3)
- Hemolytic-Uremic Syndrome (HUS): Monitor renal function prior to initiation and during therapy. Discontinue INFUGEM for HUS or severe renal impairment. (5.4)
- Hepatic Toxicity: Monitor hepatic function prior to initiation and during therapy. Discontinue INFUGEM for severe hepatic toxicity. (5.5)
- Embryo-Fetal Toxicity: Can cause fetal harm. Advise females and males of reproductive potential to use effective contraception. (5.6, 8.1)
- Exacerbation of Radiation Therapy Toxicity: May cause severe and life-threatening toxicity when administered during or within 7 days of radiation therapy. (5.7)
- Capillary Leak Syndrome: Discontinue INFUGEM. (5.8)
- Posterior Reversible Encephalopathy Syndrome (PRES): Discontinue INFUGEM. (5.9)

ADVERSE REACTIONS

The most common adverse reactions for the single agent (≥20%) are nausea/vomiting, anemia, hepatic transaminitis, neutropenia, increased alkaline phosphatase, proteinuria, fever, hematuria, rash, thrombocytopenia, dyspnea, and peripheral edema. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Sun Pharmaceutical Industries, Inc. at 1-800-818-4555 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

Lactation: Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION

Revised: 07/2018

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 1.1 Ovarian Cancer
- 1.2 Breast Cancer
- 1.3 Non-Small Cell Lung Cancer
- 1.4 Pancreatic Cancer

2 DOSAGE AND ADMINISTRATION

- 2.1 Ovarian Cancer
- 2.2 Breast Cancer
- 2.3 Non-Small Cell Lung Cancer
- 2.4 Pancreatic Cancer
- 2.5 Dose Modifications for Non-Hematologic Adverse Reactions
- 2.6 Infusion Bag Selection and Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Schedule-Dependent Toxicity
- 5.2 Myelosuppression
- 5.3 Pulmonary Toxicity and Respiratory Failure
- 5.4 Hemolytic Uremic Syndrome
- 5.5 Hepatic Toxicity
- 5.6 Embryo-Fetal Toxicity
- 5.7 Exacerbation of Radiation Therapy Toxicity
- 5.8 Capillary Leak Syndrome
- 5.9 Posterior Reversible Encephalopathy Syndrome

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

- 6.2 Postmarketing Experience

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Gender

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Ovarian Cancer
- 14.2 Breast Cancer
- 14.3 Non-Small Cell Lung Cancer (NSCLC)
- 14.4 Pancreatic Cancer

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Ovarian Cancer

INFUGEM in combination with carboplatin is indicated for the treatment of patients with advanced ovarian cancer that has relapsed at least 6 months after completion of platinum-based therapy.

1.2 Breast Cancer

INFUGEM in combination with paclitaxel is indicated for the first-line treatment of patients with metastatic breast cancer after failure of prior anthracycline-containing adjuvant chemotherapy, unless anthracyclines were clinically contraindicated.

1.3 Non-Small Cell Lung Cancer

INFUGEM is indicated in combination with cisplatin for the first-line treatment of patients with inoperable, locally advanced (Stage IIIA or IIIB), or metastatic (Stage IV) non-small cell lung cancer.

1.4 Pancreatic Cancer

INFUGEM is indicated as first-line treatment for patients with locally advanced (nonresectable Stage II or Stage III) or metastatic (Stage IV) adenocarcinoma of the pancreas. INFUGEM is indicated for patients previously treated with fluorouracil.

2 DOSAGE AND ADMINISTRATION

2.1 Ovarian Cancer

Recommended Dose and Schedule

The recommended dose of INFUGEM is 1000 mg/m² as an intravenous infusion over 30 minutes on Days 1 and 8 of each 21-day cycle, in combination with carboplatin AUC 4 intravenously after INFUGEM administration on Day 1 of each 21-day cycle. Select the INFUGEM premixed bag(s) that allow for a variance of up to 5% of the BSA-calculated dose as described in Table 5 [*see Dosage and Administration (2.6)*].

Refer to the carboplatin prescribing information for additional information.

Dosage Modifications

Recommended INFUGEM dosage modifications for myelosuppression are described in Table 1 and Table 2 [*see Warnings and Precautions (5.2)*]. Refer to the dosage modifications recommendations for non-hematologic adverse reactions [*see Dosage and Administration (2.5)*].

Table 1: Recommended Dosage Modifications for INFUGEM for Myelosuppression on Day of Treatment in Ovarian Cancer

Treatment Day	Absolute Neutrophil Count (x 10 ⁶ /L)		Platelet Count (x 10 ⁶ /L)	Dosage Modification
Day 1	Greater than or equal to 1500	and	Greater than or equal to 100,000	None
	Less than 1500	or	Less than 100,000	Delay treatment cycle
Day 8	Greater than or equal to 1500	and	Greater than or equal to 100,000	None
	1000 to 1499	or	75,000 to 99,999	50% of full dose
	Less than 1000	or	Less than 75,000	Hold

Table 2: Recommended Dosage Modification for INFUGEM for Myelosuppression in Previous Cycle in Ovarian Cancer

Occurrence	Myelosuppression During Treatment Cycle	Dosage Modification
Initial Occurrence	<ul style="list-style-type: none"> Absolute neutrophil count less than 500 x 10⁶/L for more than 5 days or Absolute neutrophil count less than 100 x 10⁶/L for more than 3 days or Febrile neutropenia or Platelets less than 25,000x10⁶/L or Cycle delay of more than one week due to toxicity 	Permanently reduce INFUGEM to 800 mg/m ² on Days 1 and 8
Subsequent Occurrence	If any of the above toxicities occur after the initial dose reduction	Permanently reduce INFUGEM dose to 800 mg/m ² on Day 1 only

2.2 Breast Cancer

Recommended Dose and Schedule

The recommended dose of INFUGEM is 1250 mg/m² intravenously over 30 minutes on Days 1 and 8 of each 21-day cycle that includes paclitaxel. Paclitaxel should be administered at 175 mg/m² on Day 1 as a 3-hour intravenous infusion before INFUGEM administration. Select the INFUGEM premixed bag(s) that allow for a variance of up to 5% of the BSA-calculated dose as described in Table 6 [see *Dosage and Administration (2.6)*].

Refer to the paclitaxel prescribing information for additional information.

Dosage Modifications

Recommended INFUGEM dosage modifications for myelosuppression are described in Table 3 [see *Warnings and Precautions (5.2)*]. Refer to the dosage modification recommendations for non-hematologic adverse reactions [see *Dosage and Administration (2.5)*].

Table 3: Recommended Dosage Modifications for INFUGEM for Myelosuppression on Day of Treatment in Breast Cancer

Treatment Day	Absolute Neutrophil Count (x 10 ⁶ /L)		Platelet Count (x 10 ⁶ /L)	Dosage Modification
Day 1	Greater than or equal to 1500	and	Greater than or equal to 100,000	None
	Less than 1500	or	Less than 100,000	Hold
Day 8	Greater than or equal to 1200	and	Greater than 75,000	None
	1000 to 1199	or	50,000 to 75,000	75% of full dose
	700 to 999	and	Greater than or equal to 50,000	50% of full dose
	Less than 700	or	Less than 50,000	Hold

2.3 Non-Small Cell Lung Cancer

Recommended Dose and Schedule

Every 4-week schedule

The recommended dose of INFUGEM is 1000 mg/m² intravenously over 30 minutes on Days 1, 8, and 15 in combination with cisplatin therapy. Administer cisplatin intravenously at 100 mg/m² on Day 1 after the infusion of INFUGEM. Select the INFUGEM premixed bag(s) that allow for a variance of up to 5% of the BSA-calculated dose as described in Table 5 [see *Dosage and Administration (2.6)*].

Every 3-week schedule

The recommended dose of INFUGEM is 1250 mg/m² intravenously over 30 minutes on Days 1 and 8 in combination with cisplatin therapy. Administer cisplatin intravenously at 100 mg/m² on Day 1 after the infusion of INFUGEM. Select the INFUGEM premixed bag(s) that allow for a variance of up to 5% of the BSA-calculated dose as described in Table 6 [see *Dosage and Administration (2.6)*].

Refer to the cisplatin prescribing information for additional information.

Dosage Modifications

Recommended INFUGEM dosage modifications for myelosuppression are described in Table 4 [see *Dosage and Administration (2.4)*, *Warnings and Precautions (5.2)*]. Refer to the dosage modifications recommendations for non-hematologic adverse reactions [see *Dosage and Administration (2.5)*].

2.4 Pancreatic Cancer

Recommended Dose and Schedule

The recommended dose of INFUGEM is 1000 mg/m² over 30 minutes intravenously. The recommended treatment schedule is as follows:

- Weeks 1 to 8: weekly dosing for the first 7 weeks followed by one week rest.
- After week 8: weekly dosing on Days 1, 8, and 15 of 28-day cycles.

Select the INFUGEM premixed bag(s) that allow for a variance of up to 5% of the BSA-calculated dose as described in Table 5 [see *Dosage and Administration (2.6)*].

Dosage Modifications

Recommended dosage modifications for INFUGEM for myelosuppression are described in Table 4 [see *Warnings and Precautions (5.2)*]. Refer to the dosage modifications recommendations for non-hematologic adverse reactions [see *Dosage and Administration (2.5)*].

Table 4: Recommended INFUGEM Dosage Modifications for Myelosuppression in Pancreatic Cancer and Non-Small Cell Lung Cancer

Absolute Neutrophil Count (x 10⁶/L)		Platelet Count (x 10⁶/L)	Dosage Modification
Greater than or equal to 1000	and	Greater than or equal to 100,000	None
500 to 999	or	50,000 to 99,999	75% of full dose
Less than 500	or	Less than 50,000	Hold

2.5 Dosage Modifications for Non-Hematologic Adverse Reactions

Permanently discontinue INFUGEM for any of the following:

- Unexplained dyspnea or other evidence of severe pulmonary toxicity [see *Warnings and Precautions (5.3)*]
- Hemolytic uremic syndrome [see *Warnings and Precautions (5.4)*]
- Severe hepatic toxicity [see *Warnings and Precautions (5.5)*]
- Capillary leak syndrome [see *Warnings and Precautions (5.8)*]
- Posterior reversible encephalopathy syndrome [see *Warnings and Precautions (5.9)*]

Withhold INFUGEM or reduce dose by 50% for other severe (Grade 3 or 4) non-hematological toxicity until resolved.

2.6 Infusion Bag Selection and Administration

See the INFUGEM Instructions for Use for additional information concerning premixed infusion bag(s) selection and spiking the infusion bag instructions.

Infusion Bag Selection

INFUGEM is provided in premixed bags that are ready for infusion and do not require any further preparation prior to use. Do not dilute prior to use. Do not remove or add medication.

Select the INFUGEM premixed bag(s) for infusion based on the patient's BSA range as outlined below in Table 5 for 1000 mg/m² (ovarian cancer, non-small cell lung cancer, and pancreatic cancer) and Table 6 for 1250 mg/m² (breast cancer, non-small cell lung cancer). The INFUGEM administered dose may vary from the BSA-calculated dose by no more than 5%.

Use another formulation of gemcitabine for patients who require a dose that is less than those listed in the Table 5 or Table 6 below (i.e., <1150 mg).

Table 5: INFUGEM Infusion Bag(s) Selection for Gemcitabine Doses of 1000 mg/m² (Non-Small Cell Lung Cancer, Ovarian Cancer, Pancreatic Cancer)

BSA Range (m²)	INFUGEM Infusion Bag(s)
1.16 to 1.25	1200 mg
1.26 to 1.35	1300 mg
1.36 to 1.45	1400 mg
1.46 to 1.55	1500 mg
1.56 to 1.65	1600 mg
1.66 to 1.75	1700 mg
1.76 to 1.85	1800 mg
1.86 to 1.95	1900 mg
1.96 to 2.10	2000 mg
2.11 to 2.30	2200 mg
2.31 to 2.45	2400 mg (1200 mg and 1200 mg)
2.46 to 2.55	2500 mg (1200 mg and 1300 mg)
2.56 to 2.64	2600 mg (1300 mg and 1300 mg) ^a

^a Suggested combination. Other possible combinations can be used to reach the appropriate dose.

Table 6: INFUGEM Infusion Bag(s) Selection for Gemcitabine Doses of 1250 mg/m² (Breast Cancer Non-Small Cell Lung Cancer)

BSA (m²)	INFUGEM Infusion Bag(s)
1.16 to 1.24	1500 mg
1.25 to 1.32	1600 mg
1.33 to 1.40	1700 mg
1.41 to 1.47	1800 mg
1.48 to 1.56	1900 mg
1.57 to 1.68	2000 mg
1.69 to 1.84	2200 mg
1.85 to 1.96	2400 mg (1200 mg and 1200 mg)
1.97 to 2.04	2500 mg (1300 mg and 1200 mg)
2.05 to 2.12	2600 mg (1300 mg and 1300 mg) ^a
2.13 to 2.20	2700 mg (1200 mg and 1500 mg) ^a
2.21 to 2.28	2800 mg (1400 mg and 1400 mg) ^a
2.29 to 2.36	2900 mg (1200 mg and 1700 mg) ^a
2.37 to 2.44	3000 mg (1500 mg and 1500 mg) ^a
2.45 to 2.52	3100 mg (1200 mg and 1900 mg) ^a
2.53 to 2.60	3200 mg (1600 mg and 1600 mg) ^a
2.61 to 2.64	3300 mg (1600 mg and 1700 mg) ^a

^a Combinations represented above are suggested combinations. Other possible combinations of bags can be used to reach the appropriate dose.

Administration

Infuse all doses of INFUGEM over 30 minutes. If two premixed infusion bags are required to achieve the

prescribed dose, infuse the total volume of both bags over 30 minutes.

After removing the overwrap check for leaks by squeezing the inner bag firmly. If leaks are found, discard the bag.

INFUGEM injection is a clear, colorless solution. Visually inspect for any particulate matter or discoloration prior to use. Discard if particulate matter or discoloration is found.

INFUGEM is a cytotoxic drug. Follow applicable special handling and disposal procedures.¹

Exercise caution and wear gloves when handling INFUGEM. Immediately wash the skin thoroughly or rinse the mucosa with copious amounts of water if INFUGEM contacts the skin or mucus membranes. Death has occurred in animal studies due to dermal absorption.

3 DOSAGE FORMS AND STRENGTHS

Injection: 10 mg/mL of gemcitabine, a clear, colorless, sterile solution in sodium chloride available in the following single-dose, premixed intravenous infusion bags:

- 1200 mg gemcitabine in 0.9% sodium chloride injection (1200 mg/120 mL)
- 1300 mg gemcitabine in 0.9% sodium chloride injection (1300 mg/130 mL)
- 1400 mg gemcitabine in 0.9% sodium chloride injection (1400 mg/140 mL)
- 1500 mg gemcitabine in 0.9% sodium chloride injection (1500 mg/150 mL)
- 1600 mg gemcitabine in 0.9% sodium chloride injection (1600 mg/160 mL)
- 1700 mg gemcitabine in 0.9% sodium chloride injection (1700 mg/170 mL)
- 1800 mg gemcitabine in 0.9% sodium chloride injection (1800 mg/180 mL)
- 1900 mg gemcitabine in 0.9% sodium chloride injection (1900 mg/190 mL)
- 2000 mg gemcitabine in 0.9% sodium chloride injection (2000 mg/200 mL)
- 2200 mg gemcitabine in 0.9% sodium chloride injection (2200 mg/220 mL)

4 CONTRAINDICATIONS

INFUGEM is contraindicated in patients with a known hypersensitivity to gemcitabine [*see Adverse Reactions (6.1)*].

5 WARNINGS AND PRECAUTIONS

5.1 Schedule-Dependent Toxicity

In clinical trials evaluating the maximum tolerated dose of gemcitabine, prolongation of the infusion time beyond 60 minutes or more frequent than weekly dosing resulted in an increased incidence of clinically significant hypotension, severe flu-like symptoms, myelosuppression, and asthenia. The half-life of gemcitabine is influenced by the length of the infusion [*see Clinical Pharmacology (12.3)*]. Refer to the recommended INFUGEM dosing schedule [*see Dosage and Administration (2.1, 2.2, 2.3, 2.4)*].

5.2 Myelosuppression

Myelosuppression manifested by neutropenia, thrombocytopenia, and anemia occurs with INFUGEM as a single agent, and the risks are increased when INFUGEM is combined with other cytotoxic drugs. In clinical trials, Grade 3-4 neutropenia, anemia, and thrombocytopenia occurred in 25%, 8%, and 5%, respectively of patients

receiving single-agent gemcitabine. The frequencies of Grade 3-4 neutropenia, anemia, and thrombocytopenia varied from 48% to 71%, 8 to 28%, and 5 to 55%, respectively, in patients receiving gemcitabine in combination with another drug [see *Adverse Reactions (6.1)*].

Monitor patients receiving INFUGEM prior to each dose with a complete blood count (CBC), including differential and platelet count, and modify the dosage as recommended [see *Dosage and Administration (2.1, 2.2, 2.3, and 2.4)*].

5.3 Pulmonary Toxicity and Respiratory Failure

Pulmonary toxicity, including interstitial pneumonitis, pulmonary fibrosis, pulmonary edema, and adult respiratory distress syndrome (ARDS), has been reported. In some cases, these pulmonary events can lead to fatal respiratory failure despite discontinuation of therapy. The onset of pulmonary symptoms may occur up to 2 weeks after the last dose of INFUGEM.

Permanently discontinue INFUGEM in patients who develop unexplained dyspnea, with or without bronchospasm, or have any evidence of pulmonary toxicity [see *Adverse Reactions (6.1 and 6.2)*].

5.4 Hemolytic Uremic Syndrome

Hemolytic uremic syndrome (HUS), including fatalities from renal failure or the requirement for dialysis, can occur in patients treated with INFUGEM. In clinical trials, HUS was reported in 6 of 2429 patients (0.25%). Most fatal cases of renal failure were due to HUS [see *Adverse Reactions (6.1 and 6.2)*].

Assess renal function prior to initiation of INFUGEM and periodically during treatment. Consider the diagnosis of HUS in patients who develop anemia with evidence of microangiopathic hemolysis, elevation of bilirubin or LDH, or reticulocytosis; severe thrombocytopenia; or evidence of renal failure (elevation of serum creatinine or BUN). Permanently discontinue INFUGEM in patients with HUS or severe renal impairment. Renal failure may not be reversible even with discontinuation of therapy.

5.5 Hepatic Toxicity

Drug-induced liver injury, including liver failure and death, has been reported in patients receiving gemcitabine alone or in combination with other potentially hepatotoxic drugs [see *Adverse Reactions (6.1 and 6.2)*]. Administration of INFUGEM in patients with concurrent liver metastases or a preexisting medical history of hepatitis, alcoholism, or liver cirrhosis can lead to exacerbation of the underlying hepatic insufficiency.

Assess hepatic function prior to initiation of INFUGEM and periodically during treatment. Permanently discontinue INFUGEM in patients that develop severe liver injury.

5.6 Embryo-Fetal Toxicity

INFUGEM can cause fetal harm when administered to a pregnant woman, based on its mechanism of action. Gemcitabine was teratogenic, embryotoxic, and fetotoxic in mice and rabbits.

Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with INFUGEM and for 6 months after the final dose. Advise male patients with female partners of reproductive potential to use effective contraception during and for 3 months following the final dose of INFUGEM [see *Use in Specific Populations (8.1, 8.3)*].

5.7 Exacerbation of Radiation Therapy Toxicity

INFUGEM is not recommended for use in combination with radiation therapy.

Concurrent (given together or ≤ 7 days apart) — Life-threatening mucositis, especially esophagitis and pneumonitis occurred in a trial in which gemcitabine was administered at a dose of 1000 mg/m² to patients with non-small cell lung cancer for up to 6 consecutive weeks concurrently with thoracic radiation.

Non-concurrent (given >7 days apart) — Excessive toxicity has not been observed when gemcitabine is administered more than 7 days before or after radiation. Radiation recall has been reported in patients who receive gemcitabine after prior radiation.

5.8 Capillary Leak Syndrome

Capillary leak syndrome (CLS) with severe consequences has been reported in patients receiving gemcitabine as a single agent or in combination with other chemotherapeutic agents. Permanently discontinue INFUGEM if CLS develops during therapy.

5.9 Posterior Reversible Encephalopathy Syndrome

Posterior reversible encephalopathy syndrome (PRES) has been reported in patients receiving gemcitabine as a single agent or in combination with other chemotherapeutic agents. PRES can present with headache, seizure, lethargy, hypertension, confusion, blindness, and other visual and neurologic disturbances.

Confirm the diagnosis of PRES with magnetic resonance imaging (MRI) and permanently discontinue INFUGEM if PRES develops during therapy.

6 ADVERSE REACTIONS

The following serious adverse reactions are discussed in greater detail in another section of the label

- Hypersensitivity [*see Contraindications (4)*]
- Schedule-Dependent Toxicity [*see Warnings and Precautions (5.1)*]
- Myelosuppression [*see Warnings and Precautions (5.2)*]
- Pulmonary Toxicity and Respiratory Failure [*see Warnings and Precautions (5.3)*]
- Hemolytic Uremic Syndrome [*see Warnings and Precautions (5.4)*]
- Hepatic Toxicity [*see Warnings and Precautions (5.5)*]
- Capillary Leak Syndrome [*see Warnings and Precautions (5.8)*]
- Posterior Reversible Encephalopathy Syndrome [*see Warnings and Precautions (5.9)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Single-Agent Use:

The data described below reflect exposure to gemcitabine as a single agent administered at doses between 800 mg/m² to 1250 mg/m² over 30 minutes intravenously, once weekly, in 979 patients with a variety of malignancies. The most common ($\geq 20\%$) adverse reactions of single-agent gemcitabine are nausea/vomiting, anemia, increased ALT, increased AST, neutropenia, increased alkaline phosphatase, proteinuria, fever, hematuria, rash,

thrombocytopenia, dyspnea, and edema. The most common ($\geq 5\%$) Grade 3 or 4 adverse reactions were neutropenia, nausea/vomiting, increased ALT, increase alkaline phosphatase, anemia, increased AST, and thrombocytopenia. Approximately 10% of the 979 patients discontinued gemcitabine due to adverse reactions. Adverse reactions resulting in discontinuation of gemcitabine in 2% of 979 patients were cardiovascular adverse reactions (myocardial infarction, cerebrovascular accident, arrhythmia, and hypertension) and adverse reactions resulting in discontinuation of gemcitabine in less than 1% of the 979 patients were anemia, thrombocytopenia, hepatic dysfunction, renal dysfunction, nausea/vomiting, fever, rash, dyspnea, hemorrhage, infection, stomatitis, somnolence, flu-like syndrome, and edema.

Tables 7 and 8 present the incidence of adverse reactions and laboratory abnormalities, respectively, reported in 979 patients with various malignancies receiving single-agent gemcitabine across 5 clinical trials.

Table 7: Selected Adverse Reactions Occurring in $\geq 10\%$ Patients Receiving Single-Agent Gemcitabine ^a

Adverse Reaction ^{c,d}	All Patients ^b		
	All Grades (%)	Grade 3 (%)	Grade 4 (%)
Nausea and Vomiting	69	13	1
Fever	41	2	0
Rash	30	<1	0
Dyspnea	23	3	<1
Diarrhea	19	1	0
Hemorrhage	17	<1	<1
Infection	16	1	<1
Alopecia	15	<1	0
Stomatitis	11	<1	0
Somnolence	11	<1	<1
Paresthesias	10	<1	0

^a Grade based on criteria from the World Health Organization (WHO)

^b N=699-974

^c Regardless of causality

^d For approximately 60% of patients, non-laboratory adverse reactions were graded only if assessed to be possibly drug-related.

Table 8: Selected Laboratory Abnormalities Occurring in Patients Receiving Single-Agent Gemcitabine^a

Laboratory Abnormality ^c	All Patients ^b		
	All Grades (%)	Grade 3 (%)	Grade 4 (%)
Hematologic			
Anemia	68	7	1
Neutropenia	63	19	6
Thrombocytopenia	24	4	1
Hepatic			
Increased ALT	68	8	2
Increased AST	67	6	2
Increased Alkaline Phosphatase	55	7	2
Hyperbilirubinemia	13	2	<1
Renal			
Proteinuria	45	<1	0

	All Patients ^b		
	All Grades (%)	Grade 3 (%)	Grade 4 (%)
Hematuria	35	<1	0
Increased BUN	16	0	0
Increased Creatinine	8	<1	0

^a Grade based on criteria from the World Health Organization (WHO)

^b N=699-974

^c Regardless of causality

Additional adverse reactions include the following:

- Transfusion requirements — Red blood cell transfusions (19%); platelet transfusions (<1%)
- Fever — Fever occurred in the absence of clinical infection and frequently in combination with other flu-like symptoms.
- Pulmonary — Dyspnea unrelated to underlying disease and sometimes accompanied by bronchospasm
- Edema — Edema (13%), peripheral edema (20%), and generalized edema (<1%)
- Flu-like Symptoms — Fever, asthenia, anorexia, headache, cough, chills, myalgia, asthenia insomnia, rhinitis, sweating, and/or malaise (19%)
- Infection — Sepsis (<1%)
- Extravasation — Injection-site reactions (4%)
- Allergic — Bronchospasm (<2%); anaphylactoid reactions

Non-Small Cell Lung Cancer:

Tables 9 and 10 present the incidence of selected adverse reactions and laboratory abnormalities respectively, occurring in $\geq 10\%$ of gemcitabine-treated patients and at a higher incidence in the gemcitabine with cisplatin arm, reported in a randomized trial (Study 3) of gemcitabine with cisplatin (n=262) administered in 28-day cycles as compared to cisplatin alone (n=260) in patients receiving first-line treatment for locally advanced or metastatic non-small cell lung cancer (NSCLC) [see *Clinical Studies (14.3)*].

Patients randomized to gemcitabine with cisplatin received a median of 4 cycles of treatment and those randomized to cisplatin received a median of 2 cycles of treatment. In this trial, the requirement for dose adjustments (>90% versus 16%), discontinuation of treatment for adverse reactions (15% versus 8%), and the proportion of patients hospitalized (36% versus 23%) were all higher for patients receiving gemcitabine with cisplatin compared to those receiving cisplatin alone. The incidence of febrile neutropenia (9/262 versus 2/260), sepsis (4% versus 1%), Grade 3 cardiac dysrhythmias (3% versus <1%) were all higher in the gemcitabine with cisplatin arm compared to the cisplatin alone arm. The two-drug combination was more myelosuppressive with 4 (1.5%) possible treatment-related deaths, including 3 cases resulting from myelosuppression with infection and one case of renal failure associated with pancytopenia and infection. No deaths due to treatment were reported on the cisplatin arm.

Table 9: Selected Adverse Reactions Occurring in $\geq 10\%$ of Patients Receiving Gemcitabine with Cisplatin and at a Higher Incidence than Patients Receiving Single-Agent Cisplatin [Between Arm Difference of $\geq 5\%$ (All Grades) or $\geq 2\%$ (Grades 3-4)] in Study 3^a

Adverse Reaction ^d	Gemcitabine with Cisplatin ^b			Cisplatin ^c		
	All Grades (%)	Grade 3 (%)	Grade 4 (%)	All Grades (%)	Grade 3 (%)	Grade 4 (%)
Nausea	93	25	2	87	20	<1

	Gemcitabine with Cisplatin ^b (%)			Cisplatin ^c (%)		
	All Grades	Grade 3	Grade 4	All Grades	Grade 3	Grade 4
Adverse Reaction^d						
Vomiting	78	11	12	71	10	9
Alopecia	53	1	0	33	0	0
Neuro Motor	35	12	0	15	3	0
Diarrhea	24	2	2	13	0	0
Neuro Sensory	23	1	0	18	1	0
Infection	18	3	2	12	1	0
Fever	16	0	0	5	0	0
Neuro Cortical	16	3	1	9	1	0
Neuro Mood	16	1	0	10	1	0
Local	15	0	0	6	0	0
Neuro Headache	14	0	0	7	0	0
Stomatitis	14	1	0	5	0	0
Hemorrhage	14	1	0	4	0	0
Hypotension	12	1	0	7	1	0
Rash	11	0	0	3	0	0

^a National Cancer Institute Common Toxicity Criteria (CTC) for severity grading

^b N=217-253; all gemcitabine with cisplatin patients with laboratory or non-laboratory data gemcitabine at 1000 mg/m² on Days 1, 8, and 15 and cisplatin at 100 mg/m² on Day 1 every 28 days

^c N=213-248; all cisplatin patients with laboratory or non-laboratory data; cisplatin at 100 mg/m² on Day 1 every 28 days

^d Adverse Reactions were graded only if assessed to be possibly drug-related.

Table 10: Selected Laboratory Abnormalities Occurring in ≥10% of Patients Receiving Gemcitabine with Cisplatin and at a Higher Incidence than Patients Receiving Single-Agent Cisplatin [Between Arm Difference of ≥5% (All Grades) or ≥2% (Grades 3-4)] in Study 3^a

	Gemcitabine with Cisplatin ^b (%)			Cisplatin ^c (%)		
	All Grades	Grade 3	Grade 4	All Grades	Grade 3	Grade 4
Laboratory Abnormality^d						
Hematologic						
Anemia	89	22	3	67	6	1
Thrombocytopenia	85	25	25	13	3	1
Neutropenia	79	22	35	20	3	1
Lymphopenia	75	25	18	51	12	5
RBC Transfusion ^e	39			13		
Platelet Transfusions ^e	21			<1		
Hepatic						
Increased Transaminases	22	2	1	10	1	0
Increased Alkaline Phosphatase	19	1	0	13	0	0
Renal						
Elevated creatinine	38	4	<1	31	2	<1
Proteinuria	23	0	0	18	0	0
Hematuria	15	0	0	13	0	0

	Gemcitabine with Cisplatin ^b			Cisplatin ^c		
	(%)			(%)		
	All Grades	Grade 3	Grade 4	All Grades	Grade 3	Grade 4
Laboratory Abnormality^d						
Other Laboratory						
Hyperglycemia	30	4	0	23	3	0
Hypomagnesemia	30	4	3	17	2	0
Hypocalcemia	18	2	0	7	0	<1

^a National Cancer Institute Common Toxicity Criteria (CTC) for severity grading

^b N=217-253; all gemcitabine with cisplatin patients with laboratory data gemcitabine at 1000 mg/m² on Days 1, 8, and 15 and cisplatin at 100 mg/m² on Day 1 every 28 days

^c N=213-248; all cisplatin patients with laboratory data; cisplatin at 100 mg/m² on Day 1 every 28 days

^d Regardless of causality

^e Percent of patients receiving transfusions

Tables 11 and 12 present the incidence of selected adverse reactions and laboratory abnormalities, respectively, occurring in $\geq 10\%$ of gemcitabine-treated patients and at a higher incidence in the gemcitabine with cisplatin arm, reported in a randomized trial (Study 4) of gemcitabine with cisplatin (n=69) administered in 21-day cycles as compared to etoposide with cisplatin alone (n=66) in patients receiving first-line treatment for locally advanced or metastatic non-small cell lung cancer (NSCLC) [see *Clinical Studies (14.3)*].

Patients in the gemcitabine cisplatin (GC) arm received a median of 5 cycles and those in the etoposide/cisplatin (EC) arm received a median of 4 cycles. The majority of patients receiving more than one cycle of treatment required dose adjustments; 81% in the GC arm and 68% in the EC arm. The incidence of hospitalizations for treatment-related adverse reactions was 22% in the GC arm and 27% in the EC arm. The proportion of discontinuation of treatment for treatment-related adverse reactions was higher for patients in the GC arm (14% versus 8%). The proportion of patients hospitalized for febrile neutropenia was lower in the GC arm (7% versus 12%). There was one death attributed to treatment, a patient with febrile neutropenia and renal failure, which occurred in the GC arm.

Table 11: Selected Adverse Reactions Occurring in Patients Receiving Gemcitabine with Cisplatin in Study 4^a

Adverse Reaction ^d	Gemcitabine with Cisplatin ^b			Etoposide with Cisplatin ^c		
	(%)			(%)		
	All Grades	Grade 3	Grade 4	All Grades	Grade 3	Grade 4
Nausea and Vomiting	96	35	4	86	19	7
Alopecia	77	13	0	92	51	0
Paresthesias	38	0	0	16	2	0
Infection	28	3	1	21	8	0
Stomatitis	20	4	0	18	2	0
Diarrhea	14	1	1	13	0	2
Edema ^e	12	-	-	2	-	-
Rash	10	0	0	3	0	0
Hemorrhage	9	0	3	3	0	3
Fever	6	0	0	3	0	0
Flu-like Syndrome ^e	3	-	-	0	-	-
Somnolence	3	0	0	3	2	0
Dyspnea	1	0	1	3	0	0

^a Grade based on criteria from the World Health Organization (WHO)

^b N=67-69; all gemcitabine with cisplatin patients with laboratory or non-laboratory data gemcitabine at 1250 mg/m² on Days 1 and 8 and cisplatin at 100 mg/m² on Day 1 every 21 days

^c N=57-63; all cisplatin with etoposide patients with laboratory or non-laboratory data; cisplatin at 100 mg/m² on Day 1 and intravenous etoposide at 100 mg/m² on Days 1, 2, and 3 every 21 days

^d Non-laboratory events were graded only if assessed to be possibly drug-related. Pain data were not collected.

^e Flu-like syndrome and edema were not graded.

Table 12: Selected Laboratory Abnormalities Occurring in Patients Receiving Gemcitabine with Cisplatin in Study 4^a

Laboratory Abnormality ^d	Gemcitabine with Cisplatin ^b (%)			Etoposide with Cisplatin ^c (%)		
	All Grades	Grade 3	Grade 4	All Grades	Grade 3	Grade 4
Hematologic						
Anemia	88	22	0	77	13	2
Neutropenia	88	36	28	87	20	56
Thrombocytopenia	81	39	16	45	8	5
RBC Transfusions ^e	29	-	-	21	-	-
Platelet Transfusions ^e	3	-	-	8	-	-
Hepatic						
Increased Alkaline Phosphatase	16	0	0	11	0	0
Increased ALT	6	0	0	12	0	0
Increased AST	3	0	0	11	0	0
Bilirubin	0	0	0	0	0	0
Renal						
Hematuria	22	0	0	10	0	0
Proteinuria	12	0	0	5	0	0
BUN	6	0	0	4	0	0
Creatinine	2	0	0	2	0	0

^a Grade based on criteria from the World Health Organization (WHO)

^b N=67-69; all gemcitabine with cisplatin patients with non-laboratory data; gemcitabine at 1250 mg/m² on Days 1 and 8 and cisplatin at 100 mg/m² on Day 1 every 21 days

^c N=57-63; all cisplatin with etoposide patients with non-laboratory data; cisplatin at 100 mg/m² on Day 1 and intravenous etoposide at 100 mg/m² on Days 1, 2, and 3 every 21 days

^d Regardless of causality

^e WHO grading scale not applicable to proportion of patients with transfusions

Breast Cancer

Tables 13 and 14 present the incidence of selected adverse reactions and laboratory abnormalities respectively, occurring in ≥10% of gemcitabine-treated patients and at a higher incidence in the gemcitabine with paclitaxel arm, reported in a randomized trial (Study 2) of gemcitabine with paclitaxel (n=262) compared to paclitaxel alone (n=259) for the first-line treatment of metastatic breast cancer (MBC) in women who received anthracycline-containing chemotherapy in the adjuvant/neo-adjuvant setting or for whom anthracyclines were contraindicated [see *Clinical Studies (14.2)*].

The requirement for dose reduction of paclitaxel was higher for patients in the gemcitabine/paclitaxel arm (5% versus 2%). The number of paclitaxel doses omitted (<1%), the proportion of patients discontinuing treatment for treatment-related adverse reactions (7% versus 5%), and the number of treatment-related deaths (1 patient in each

arm) were similar between the two arms.

Table 13: Selected Adverse Reactions Occurring in Patients Receiving Gemcitabine with Paclitaxel and at a Higher Incidence than Patients Receiving Single-Agent Paclitaxel [Between Arm Difference of $\geq 5\%$ (All Grades) or $\geq 2\%$ (Grades 3-4)] in Study 2^a

	Gemcitabine with Paclitaxel (N=262) (%)			Paclitaxel (N=259) (%)		
	All Grades	Grade 3	Grade 4	All Grades	Grade 3	Grade 4
Adverse Reaction^b						
Alopecia	90	14	4	92	19	3
Neuropathy-sensory	64	5	<1	58	3	0
Nausea	50	1	0	31	2	0
Fatigue	40	6	<1	28	1	<1
Vomiting	29	2	0	15	2	0
Diarrhea	20	3	0	13	2	0
Anorexia	17	0	0	12	<1	0
Neuropathy-motor	15	2	<1	10	<1	0
Stomatitis/pharyngitis	13	1	<1	8	<1	0
Fever	13	<1	0	3	0	0
Rash/desquamation	11	<1	<1	5	0	0
Febrile neutropenia	6	5	<1	2	1	0

^a Severity grade based on National Cancer Institute Common Toxicity Criteria (CTC) Version 2.0

^b Regardless of causality

^c Non-laboratory events were graded only if assessed to be possibly drug-related.

Table 14: Selected Laboratory Abnormalities Occurring in $\geq 10\%$ of Patients Receiving Gemcitabine with Paclitaxel and at a Higher Incidence than Patients Receiving Single-Agent Paclitaxel [Between Arm Difference of $\geq 5\%$ (All Grades) or $\geq 2\%$ (Grades 3-4)] in Study 2^a

	Gemcitabine Paclitaxel (N=262) (%)			Paclitaxel (N=259) (%)		
	All Grades	Grade 3	Grade 4	All Grades	Grade 3	Grade 4
Laboratory Abnormality^b						
Hematologic						
Anemia	69	6	1	51	3	<1
Neutropenia	69	31	17	31	4	7
Thrombocytopenia	26	5	<1	7	<1	<1
Hepatobiliary						
Increased ALT	18	5	<1	6	<1	0
Increased AST	16	2	0	5	<1	0

^a Severity grade based on National Cancer Institute Common Toxicity Criteria (CTC) Version 2.0

^b Regardless of causality

Clinically relevant Grade 3 or 4 dyspnea occurred with a higher incidence in the gemcitabine with paclitaxel arm compared with the paclitaxel arm (1.9% versus 0).

Ovarian Cancer

Tables 15 and 16 present the incidence of selected adverse reactions and laboratory abnormalities respectively, occurring in $\geq 10\%$ of gemcitabine-treated patients and at a higher incidence in the gemcitabine with carboplatin arm, reported in a randomized trial (Study 1) of gemcitabine with carboplatin (n=175) compared to carboplatin alone (n=174) for the second-line treatment of ovarian cancer in women with disease that had relapsed more than 6 months following first-line platinum-based chemotherapy [see *Clinical Studies (14.1)*]. Additional clinically significant adverse reactions, occurring in less than 10% of patients, are provided following Table 16.

The proportion of patients with dose adjustments for carboplatin (1.8% versus 3.8%), doses of carboplatin omitted (0.2% versus 0), and discontinuing treatment for treatment-related adverse reactions (10.9% versus 9.8%), were similar between arms. Dose adjustment for gemcitabine occurred in 10.4% of patients and gemcitabine dose was omitted in 13.7% of patients in the gemcitabine/carboplatin arm.

Table 15: Adverse Reactions Occurring in $\geq 10\%$ of Patients Receiving in Gemcitabine with Carboplatin and at a Higher Incidence than Patients Receiving Single-Agent Carboplatin [Between Arm Difference of $\geq 5\%$ (All Grades) or $\geq 2\%$ (Grades 3-4)] in Study 1^a

	Gemcitabine with Carboplatin (N=175) (%)			Carboplatin (N=174) (%)		
	All Grades	Grade 3	Grade 4	All Grades	Grade 3	Grade 4
Adverse Reaction^b						
Nausea	69	6	0	61	3	0
Alopecia	49	0	0	17	0	0
Vomiting	46	6	0	36	2	<1
Constipation	42	6	1	37	3	0
Fatigue	40	3	<1	32	5	0
Diarrhea	25	3	0	14	<1	0
Stomatitis/pharyngitis	22	<1	0	13	0	0

^a Grade based on Common Toxicity Criteria (CTC) Version 2.0

^b Regardless of causality

Table 16: Laboratory Abnormalities Occurring in Patients Receiving Gemcitabine with Carboplatin and at a Higher Incidence than Patients Receiving Single-Agent Carboplatin [Between Arm Difference of $\geq 5\%$ (All Grades) or $\geq 2\%$ (Grades 3-4)] in Study 1^a

	Gemcitabine plus Carboplatin (N=175) (%)			Carboplatin (N=174) (%)		
	All Grades	Grade 3	Grade 4	All Grades	Grade 3	Grade 4
Laboratory Abnormality^b						
Hematologic						
Neutropenia	90	42	29	58	11	1
Anemia	86	22	6	75	9	2
Thrombocytopenia	78	30	5	57	10	1
RBC Transfusions ^c	38			15		
Platelet Transfusions ^c	9			3		

^a Grade based on National Cancer Institute Common Toxicity Criteria (NCI CTC) Version 2.0

^b Regardless of causality

^cPercent of patients receiving transfusions. Transfusions are not CTC-graded events. Blood transfusions included both packed red blood cells and whole blood.

Hematopoietic growth factors were administered more frequently in the gemcitabine-containing arm: granulocyte growth factors (23.6% and 10.1%) and erythropoietic agents (7.3% and 3.9%).

The following clinically relevant, Grade 3 and 4 adverse reactions occurred more frequently in the gemcitabine with carboplatin arm: dyspnea (3.4% versus 2.9%), febrile neutropenia (1.1% versus 0), hemorrhagic event (2.3% versus 1.1%), motor neuropathy (1.1% versus 0.6%), and rash/desquamation (0.6% versus 0).

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of gemcitabine. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Cardiovascular - Congestive heart failure, myocardial infarction, arrhythmias, and supraventricular arrhythmias

Vascular Disorders - Peripheral vasculitis, gangrene, and capillary leak syndrome

Skin - Cellulitis, pseudocellulitis, severe skin reactions, including desquamation and bullous skin eruptions

Hepatic - Hepatic failure, hepatic veno-occlusive disease

Pulmonary - Interstitial pneumonitis, pulmonary fibrosis, pulmonary edema, and adult respiratory distress syndrome (ARDS)

Nervous System - Posterior reversible encephalopathy syndrome (PRES)

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on animal data and its mechanism of action, INFUGEM can cause fetal harm when administered to a pregnant woman. There are no available data on the use of gemcitabine in pregnant women. In animal reproductive studies, gemcitabine was teratogenic, embryotoxic, and fetotoxic in mice and rabbits (*see Data*). Advise pregnant women of the potential risk to a fetus.

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

Data

Animal Data

Gemcitabine is embryotoxic in mice. Daily dosing of gemcitabine to pregnant mice increased the incidence of fetal malformations (cleft palate, incomplete ossification) at doses of 1.5 mg/kg/day (approximately 0.005 times the 1000 mg/m² clinical dose on body surface area (BSA)). Gemcitabine was embryotoxic and fetotoxic in rabbits. Daily dosing of gemcitabine to pregnant rabbits resulted in fetotoxicity (decreased fetal viability, reduced litter sizes, and developmental delays) and increased the incidence of fetal malformations (fused pulmonary artery, absence of gall bladder) at doses of 0.1 mg/kg/day (approximately 0.002 times the 1000 mg/m² clinical dose based on BSA).

8.2 Lactation

Risk Summary

There is no information regarding the presence of gemcitabine or its metabolites in human milk, or their effects on the breastfed infant or milk production. Due to the potential for serious adverse reactions in nursing infants from INFUGEM, advise woman not to breastfeed during treatment with INFUGEM and for at least one week after the last dose.

8.3 Females and Males of Reproductive Potential

Pregnancy Testing

Verify the pregnancy status of females of reproductive potential prior to initiating INFUGEM [see *Use in Specific Populations (8.1)*].

Contraception

INFUGEM can cause fetal harm when administered to a pregnant woman [see *Use in Specific Populations (8.1)*].

Females

Advise females of reproductive potential to use effective contraception during treatment with INFUGEM and for 6 months after the last dose [see *Use in Specific Populations (8.1)*].

Males

Advise male patients with female partners of reproductive potential to use effective contraception during and for 3 months following the last dose of INFUGEM [see *Nonclinical Toxicology (13.1)*].

Infertility

Males

Based on animal studies, INFUGEM may impair fertility in males of reproductive potential [see *Nonclinical Toxicology (13.1)*].

8.4 Pediatric Use

The safety and effectiveness of INFUGEM have not been established in pediatric patients. The safety and pharmacokinetics of gemcitabine were evaluated in a trial in pediatric patients with refractory leukemia. The maximum tolerated dose was 10 mg/m²/min for 360 minutes weekly for three weeks followed by a one-week rest period. The safety and activity of gemcitabine were evaluated in a trial of pediatric patients with relapsed acute lymphoblastic leukemia (22 patients) and acute myelogenous leukemia (10 patients) at a dose of 10 mg/m²/min administered over 360 minutes weekly for three weeks followed by a one-week rest period. Patients with M1 or M2 bone marrow on Day 28 who did not experience unacceptable toxicity were eligible to receive a maximum of one additional four-week course. Toxicities observed included bone marrow suppression, febrile neutropenia, elevation of serum transaminases, nausea, and rash/desquamation. No meaningful clinical activity was observed in this trial.

8.5 Geriatric Use

In clinical studies of gemcitabine enrolling 979 patients with various cancers who received gemcitabine as a single agent, no overall differences in safety were observed between patients aged 65 and older and younger patients, with the exception of a higher rate of Grade 3-4 thrombocytopenia in older patients as compared to younger patients. In a randomized trial in women with ovarian cancer, 175 women received gemcitabine with carboplatin, of which 29% were age 65 years or older. Similar effectiveness was observed between older and younger women.

There was significantly higher Grade 3/4 neutropenia in women 65 years of age or older [see *Dosage and Administration (2.1-2.4)*].

8.6 Gender


Gemcitabine clearance is decreased in females [see *Clinical Pharmacology (12.3)*]. In single-agent studies of gemcitabine, women, especially older women, were more likely not to proceed to a subsequent cycle and to experience Grade 3/4 neutropenia and thrombocytopenia [see *Dosage and Administration (2.1-2.4)*].

10 OVERDOSAGE

There is no known antidote for overdoses of gemcitabine. Myelosuppression, paresthesias, and severe rash were the principal toxicities seen when a single dose as high as 5700 mg/m² was administered by intravenous infusion over 30 minutes every 2 weeks to several patients in a dose-escalation study. In the event of suspected overdose, the patient should be monitored with appropriate blood counts and should receive supportive therapy, as necessary.

11 DESCRIPTION

Gemcitabine is a nucleoside metabolic inhibitor. Gemcitabine hydrochloride is 2'-deoxy-2',2'-difluorocytidine monohydrochloride (β -isomer) with the following structural formula:



Gemcitabine hydrochloride is a white to off-white crystalline powder. The empirical formula for gemcitabine hydrochloride is C₉H₁₁F₂N₃O₄ • HCl. It has a molecular weight of 299.66 g/mol. Gemcitabine hydrochloride is soluble in water, slightly soluble in methanol, and practically insoluble in ethanol and polar organic solvents.

INFUGEM (gemcitabine in 0.9% sodium chloride injection) is a clear, colorless, sterile solution that is provided as a single-dose, premixed intravenous infusion bag (10 mg/mL) for intravenous use and does not require any further preparation.

Each 100 mL contains 1000 mg of gemcitabine (equivalent to 1138 mg of gemcitabine hydrochloride, USP), 900 mg of sodium chloride, and water for injection. Hydrochloric acid and/or sodium hydroxide may have been added for pH adjustment.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Gemcitabine kills cells undergoing DNA synthesis and blocks the progression of cells through the G1/S-phase boundary. Gemcitabine is metabolized by nucleoside kinases to diphosphate (dFdCDP) and triphosphate (dFdCTP) nucleosides. Gemcitabine diphosphate inhibits ribonucleotide reductase, an enzyme responsible for catalyzing the reactions that generate deoxynucleoside triphosphates for DNA synthesis, resulting in reductions in deoxynucleotide concentrations, including dCTP. Gemcitabine triphosphate competes with dCTP for incorporation into DNA. The reduction in the intracellular concentration of dCTP by the action of the diphosphate enhances the incorporation of gemcitabine triphosphate into DNA (self-potential). After the gemcitabine nucleotide is incorporated into DNA, only one additional nucleotide is added to the growing DNA strands, which eventually results in the initiation of apoptotic cell death.

12.3 Pharmacokinetics

The pharmacokinetics of gemcitabine were examined in 353 patients, with various solid tumors. Pharmacokinetic parameters were derived using data from patients treated for varying durations of therapy given weekly with periodic rest weeks and using both short infusions (<70 minutes) and long infusions (70 to 285 minutes). The total gemcitabine dose varied from 500 to 3600 mg/m².

Distribution

The volume of distribution was increased with infusion length. Volume of distribution of gemcitabine was 50 L/m² following infusions lasting <70 minutes. For long infusions, the volume of distribution rose to 370 L/m².

Gemcitabine pharmacokinetics are linear and are described by a 2-compartment model. Population pharmacokinetic analyses of combined single and multiple dose studies showed that the volume of distribution of gemcitabine was significantly influenced by duration of infusion and gender. Gemcitabine plasma protein binding is negligible.

Elimination

Metabolism

The active metabolite, gemcitabine triphosphate, can be extracted from peripheral blood mononuclear cells. The half-life of the terminal phase for gemcitabine triphosphate from mononuclear cells ranges from 1.7 to 19.4 hours.

Excretion

Gemcitabine disposition was studied in 5 patients who received a single 1000 mg/m²/30 minute infusion of radiolabeled drug. Within one (1) week, 92% to 98% of the dose was recovered, almost entirely in the urine. Gemcitabine (<10%) and the inactive uracil metabolite, 2'-deoxy-2',2'-difluorouridine (dFdU), accounted for 99% of the excreted dose. The metabolite dFdU is also found in plasma.

Specific Populations

Geriatric Patients

Clearance of gemcitabine was affected by age. The lower clearance in the elderly results in higher concentrations of gemcitabine for any given dose. Differences in either clearance or volume of distribution based on patient characteristics or the duration of infusion result in changes in half-life and plasma concentrations. Table 17 shows plasma clearance and half-life of gemcitabine following short infusions for typical patients by age and gender.

Table 17: Gemcitabine Clearance and Half-Life for the “Typical” Patient

Age	Clearance Men (L/hr/m ²)	Clearance Women (L/hr/m ²)	Half-Life ^a Men (min)	Half-Life ^a Women (min)
29	92.2	69.4	42	49
45	75.7	57	48	57
65	55.1	41.5	61	73
79	40.7	30.7	79	94

^a Half-life for patients receiving <70 minute infusion.

Gemcitabine half-life for short infusions ranged from 42 to 94 minutes, and the value for long infusions varied from 245 to 638 minutes, depending on age and gender, reflecting a greatly increased volume of distribution with longer infusions.

Male and Female Patients

Clearance of gemcitabine was affected by gender. Female patients have lower clearance and longer half-lives than male patients as described in Table 17.

Patients with Renal Impairment

No clinical studies have been conducted with gemcitabine in patients with decreased renal function.

Patients with Hepatic Impairment

No clinical studies have been conducted with gemcitabine in patients with decreased hepatic function.

Drug Interactions

When gemcitabine (1250 mg/m² on Days 1 and 8) and cisplatin (75 mg/m² on Day 1) were administered in NSCLC patients, the clearance of gemcitabine on Day 1 was 128 L/hr/m² and on Day 8 was 107 L/hr/m². Analysis of data from metastatic breast cancer patients shows that, on average, gemcitabine has little or no effect on the pharmacokinetics (clearance and half-life) of paclitaxel and paclitaxel has little or no effect on the pharmacokinetics of gemcitabine. Data from NSCLC patients demonstrate that gemcitabine and carboplatin given in combination does not alter the pharmacokinetics of gemcitabine or carboplatin compared to administration of either single agent. However, due to wide confidence intervals and small sample size, interpatient variability may be observed.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term animal studies to evaluate the carcinogenic potential of gemcitabine have not been conducted. Gemcitabine was mutagenic in an in vitro mouse lymphoma (L5178Y) assay and was clastogenic in an in vivo mouse micronucleus assay. Gemcitabine intraperitoneal doses of 0.5 mg/kg/day (about 1/700 the 1000 mg/m² clinical dose based on BSA) in male mice resulted in moderate to severe hypospermatogenesis, decreased fertility, and decreased implantations. In female mice, fertility was not affected but maternal toxicities were observed at 1.5 mg/kg/day administered intravenously (about 1/200 the 1000 mg/m² clinical dose based on BSA) and fetotoxicity or embryoletality was observed at 0.25 mg/kg/day administered intravenously (about 1/1300 the 1000 mg/m² clinical dose based on BSA).

14 CLINICAL STUDIES

14.1 Ovarian Cancer

The safety and efficacy of gemcitabine were studied in a randomized trial (Study 1) of 356 women with advanced ovarian cancer that had relapsed at least 6 months after first-line platinum-based therapy. Patients were randomized to receive either gemcitabine hydrochloride 1000 mg/m² on Days 1 and 8 of a 21-day cycle and carboplatin AUC 4 administered after gemcitabine hydrochloride infusion on Day 1 of each cycle (n=178) or to carboplatin AUC 5 administered on Day 1 of each 21-day cycle (n=178). The primary efficacy outcome measure was progression free survival (PFS).

Baseline demographics and disease characteristics in gemcitabine plus carboplatin arm were: median age of 59 (range: 36 to 78), 94% ECOG PS 0-1. 8% had evaluable disease and 92% had bidimensionally measurable disease. 40% had 6 to 12 months of platinum free interval, 59% had greater than 12 months platinum free interval; and as first-line therapy, 70% had platinum-taxane combination, 29% had platinum-non-taxane combination and 1% had platinum monotherapy.

Baseline demographics and disease characteristics in the carboplatin arm were: median age of 58 (range 21 to 81), 95% had an Eastern Cooperative Oncology Group (ECOG) performance status of 0-1; 3% had evaluable disease and 96% had bidimensionally measurable disease; 40% had a 6 to 12 month platinum-free interval and 60% had a greater than 12 month platinum-free interval; and as first-line therapy, 71% had a platinum-taxane combination, 28% had platinum-non-taxane combination, and 1% had platinum monotherapy.

The addition of gemcitabine to carboplatin resulted in statistically significant improvements in PFS and overall response rate as shown in Table 18 and Figure 1. Approximately 75% of patients in each arm received additional chemotherapy for disease progression; 13 of 120 patients in the carboplatin alone arm received gemcitabine for treatment of disease progression. There was no significant difference in overall survival between the treatment arms.

Table 18: Efficacy Results for Study 1

	Gemcitabine/Carboplatin (N=178)	Carboplatin (N=178)
Progression-free Survival Median (95% CI) ^a months	8.6 (8, 9.7)	5.8 (5.2, 7.1)
Hazard Ratio (95% CI)	0.72 (0.57, 0.9)	
p-value ^b	p=0.0038	
Overall Survival Median (95% CI) months	18 (16.2, 20.3)	17.3 (15.2, 19.3)
Hazard Ratio (95% CI)	0.98 (0.78, 1.24)	
p-value ^b	p=0.8977	
Investigator Assessed Overall Response Rate	47.2%	30.9%
p-value ^c	p=0.0016	
CR ^d	14.6%	6.2%
PR plus PRNM ^e	32.6%	24.7%
Independently Assessed Overall Response Rate ^f	46.3%	35.6%
p-value ^c	p=0.11	
CR ^d	9.1%	4%

	Gemcitabine/Carboplatin (N=178)	Carboplatin (N=178)
PR plus PRNM ^e	37.2%	31.7%

^a CI=confidence interval

^b Log rank, unadjusted


^c Chi square

^d CR=Complete response

^e PR plus PRNM=Partial response plus partial response, non-measurable disease

^f Independently reviewed cohort – gemcitabine hydrochloride/carboplatin (n=121), carboplatin (n=101); independent reviewers were unable to measure disease detected by sonography or physical exam.

Figure 1: Kaplan-Meier Curve of Progression Free Survival for Study 1



14.2 Breast Cancer

The safety and efficacy of gemcitabine were evaluated in a randomized, open-label trial (Study 2) conducted in women receiving initial treatment for metastatic breast cancer and in women who had received prior adjuvant/neoadjuvant anthracycline chemotherapy unless clinically contraindicated. Patients were randomized to receive gemcitabine 1250 mg/m² on Days 1 and 8 of a 21-day cycle and paclitaxel 175 mg/m² administered prior to gemcitabine on Day 1 of each cycle (n=267) or to receive paclitaxel 175 mg/m² on Day 1 of each 21-day cycle (n=262). The primary efficacy outcome measure was time to documented disease progression.

A total of 529 patients were enrolled. Baseline demographics and disease characteristics in the gemcitabine with paclitaxel arm were: median age of 53 (range 26 to 83); 97% had metastatic disease; 70% had baseline Karnofsky Performance Status (KPS) greater than or equal to 90%; 57% had 1 to 2 tumor sites and 43% had 3 or more tumor sites; 73% had visceral disease and 97% had prior anthracycline.

Efficacy results are presented in Table 19 and Figure 2. The addition of gemcitabine to paclitaxel resulted in a statistically significant improvement in time to documented disease progression and overall response rate compared to paclitaxel alone. There was no significant difference in overall survival.

Table 19: Efficacy Results for Study 2


	Gemcitabine /Paclitaxel (N=267)	Paclitaxel (N=262)
Efficacy Outcomes		
Time to Documented Disease Progression ^b		
Median in months	5.2	2.9
(95% CI)	(4.2, 5.6)	(2.6, 3.7)
Hazard Ratio (95% CI)	0.65 (0.524, 0.805)	
p-value	p<0.0001	
Overall Survival ^c		
Median Survival in months	18.6	15.8
(95% CI)	(16.5, 20.7)	(14.1, 17.3)
Hazard Ratio (95% CI)	0.86 (0.71, 1.04)	
p-value	Not Significant	
Overall Response Rate	40.8%	22.1%
(95% CI)	(34.9, 46.7)	(17.1, 27.2)
p-value	p<0.0001	

^a Karnofsky Performance Status

^b These represent reconciliation of investigator and Independent Review Committee assessments according to a predefined algorithm.

^c Based on the ITT population

Figure 2: Kaplan-Meier Curve of Time to Documented Disease Progression for Study 2



14.3 Non-Small Cell Lung Cancer (NSCLC)

The safety and efficacy of gemcitabine were evaluated in two randomized, multicenter trials.

Study 3: 28-Day Schedule

A randomized trial (Study 3) compared gemcitabine with cisplatin to cisplatin alone in the treatment of patients

with inoperable Stage IIIA, IIIB, or IV NSCLC who had not received prior chemotherapy. Patients were randomized to receive gemcitabine 1000 mg/m² on Days 1, 8, and 15 of a 28-day cycle with cisplatin 100 mg/m² administered on Day 1 of each cycle or to receive cisplatin 100 mg/m² on Day 1 of each 28-day cycle. The primary efficacy outcome measure was overall survival. A total of 522 patients were enrolled. Patient demographics and baseline characteristics were similar between arms with the exception of histologic subtype of NSCLC, with 48% of patients on the cisplatin arm and 37% of patients on the gemcitabine with cisplatin arm having adenocarcinoma. Efficacy results are presented in Table 20 and Figure 3 for overall survival.

Study 4: 21-Day Schedule

A randomized (1:1), multicenter trial (Study 4) was conducted in 135 patients with Stage IIIB or IV NSCLC. Patients were randomized to receive gemcitabine 1250 mg/m² on Days 1 and 8, and cisplatin 100 mg/m² on Day 1 of a 21-day cycle or to receive etoposide 100 mg/m² intravenously on Days 1, 2, and 3 and cisplatin 100 mg/m² on Day 1 of a 21-day cycle. A total of 135 patients were enrolled. Patient demographics and baseline characteristics in gemcitabine with cisplatin arm were: 93% males, median age 58 years (range 33 to 76); 48% in stage IIIB and 52% stage IV, 45 %; Baseline KPS 70 to 80, 55% Baseline KPS 90 to 100. Patient demographics and baseline characteristics in cisplatin arm were: 92% males, median age 60 years (range 35 to 75), 52% in stage IIIB and 49% stage IV NSCLC, 52% Baseline KPS 70 to 80, 49% Baseline KPS 90 to 100.

There was no significant difference in survival between the two treatment arms (Log rank p=0.18, two-sided, see Table 20). The median survival was 8.7 months for the gemcitabine with cisplatin arm versus 7 months for the etoposide with cisplatin arm. Median time to disease progression for the gemcitabine with cisplatin arm was 5 months compared to 4.1 months on the etoposide with cisplatin arm (Log rank p=0.015, two-sided). The objective response rate for the gemcitabine with cisplatin arm was 33% compared to 14% on the etoposide with cisplatin arm (Fisher’s Exact p=0.01, two-sided).

Figure 3: Kaplan-Meier Survival Curve in Gemcitabine with Cisplatin versus Cisplatin for Study 4).




Table 20: Randomized Trials of Gemcitabine with Cisplatin in Patients with NSCLC Efficacy Results for Studies 3 and 4

Trial	28-day Schedule (Study 3) ^a		21-day Schedule (Study 4) ^b	
	Gemcitabine with Cisplatin (N=260)	Cisplatin (N=262)	Gemcitabine with Cisplatin (N=69)	Etoposide with Cisplatin (N=66)
Efficacy Outcomes				
Survival				
Median in months	9	7.6	8.7	7
(95% CI ^e) months	8.2, 11	6.6, 8.8	7.8, 10.1	6, 9.7
p-value ^f	p=0.008		p=0.18	
Time to Disease Progression				
Median in months	5.2	3.7	5	4.1
(95% CI ^e) months	4.2, 5.7	3, 4.3	4.2, 6.4	2.4, 4.5
p-value ^f	p=0.009		p=0.015	
Tumor Response	26%	10%	33%	14%
p-value ^f	p<0.0001		p=0.01	

^a 28-day schedule – gemcitabine with cisplatin: gemcitabine 1000 mg/m² on Days 1, 8, and 15 and cisplatin 100 mg/m² on Day 1 every 28 days; Single-agent cisplatin: cisplatin 100 mg/m² on Day 1 every 28 days

^b 21-day schedule – gemcitabine with cisplatin: gemcitabine 1250 mg/m² on Days 1 and 8 and cisplatin 100 mg/m² on Day 1 every 21 days; etoposide with cisplatin: cisplatin 100 mg/m² on Day 1 and intravenous etoposide 100 mg/m² on Days 1, 2, and 3 every 21 days

^c N/A Not applicable

^d Karnofsky Performance Status

^e CI=confidence intervals

^f p-value two-sided Fisher's Exact test for difference in binomial proportions; log rank test for time-to-event analyses

14.4 Pancreatic Cancer

The safety and efficacy of gemcitabine was evaluated in two trials (Studies 5 and 6), a randomized, single-blind, two-arm, active-controlled trial (Study 5) conducted in patients with locally advanced or metastatic pancreatic cancer who had received no prior chemotherapy and a single-arm, open-label, multicenter trial (Study 6) conducted in patients with locally advanced or metastatic pancreatic cancer previously treated with fluorouracil or a fluorouracil-containing regimen. Study 5 randomized patients to receive gemcitabine 1000 mg/m² intravenously over 30 minutes once weekly for 7 weeks followed by a one-week rest, then once weekly dosing for 3 consecutive weeks every 28 days in subsequent cycles (n=63) or to fluorouracil 600 mg/m² intravenously over 30 minutes once weekly (n=63). In Study 6, all patients received gemcitabine 1000 mg/m² intravenously over 30 minutes once weekly for 7 weeks followed by a one-week rest, then once weekly dosing for 3 consecutive weeks every 28 days in subsequent cycles.

The primary efficacy outcome measure in both trials was "clinical benefit response". A patient was considered to have had a clinical benefit response if either of the following occurred:

- The patient achieved a ≥50% reduction in pain intensity (Memorial Pain Assessment Card) or analgesic consumption, or a 20-point or greater improvement in performance status (Karnofsky Performance Status) for a period of at least 4 consecutive weeks, without showing any sustained worsening in any of the other parameters. Sustained worsening was defined as 4 consecutive weeks with either any increase in pain intensity

or analgesic consumption or a 20-point decrease in performance status occurring during the first 12 weeks of therapy.

OR

- The patient was stable on all of the aforementioned parameters, and showed a marked, sustained weight gain ($\geq 7\%$ increase maintained for ≥ 4 weeks) not due to fluid accumulation.

Study 5 enrolled 126 patients. The demographic and entry characteristics were similar between the arms. The efficacy outcome results are shown in Table 21 and for overall survival in Figure 4. Patients treated with gemcitabine had statistically significant increases in clinical benefit response, survival, and time to disease progression compared to those randomized to receive fluorouracil. No confirmed objective tumor responses were observed in either treatment arm.


Table 21: Efficacy Results for Study 5

	Gemcitabine (N=63)	Fluorouracil (N=63)
Efficacy Outcomes		
Clinical benefit response	22.2%	4.8%
p-value ^b	p=0.004	
Survival		
Median	5.7 months	4.2 months
(95% CI)	(4.7, 6.9)	(3.1, 5.1)
p-value ^b	p=0.0009	
Time to Disease Progression		
Median	2.1 months	0.9 months
(95% CI)	(1.9, 3.4)	(0.9, 1.1)
p-value ^b	p=0.0013	

^a Karnofsky Performance Status

^b p-value for clinical benefit response calculated using the two-sided test for difference in binomial proportions. All other p-values are calculated using log rank test.

Figure 4: Kaplan-Meier Survival Curve for Study 5



15 REFERENCES

1. "OSHA Hazardous Drugs." OSHA. <http://www.osha.gov/SLTC/hazardousdrugs/index.html>

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

INFUGEM (gemcitabine in 0.9% sodium chloride injection) is a clear, colorless, sterile solution in a single-dose, premixed intravenous infusion bag with an aluminum overwrap. The container closure is not made with natural rubber latex and is tamper evident. It is available in presentations as described in Table 22.

Table 22: INFUGEM Available Presentations

Strength	Package	NDC number
1200 mg in 120 mL	1 single-dose bag per carton	62756-073-60
1300 mg in 130 mL	1 single-dose bag per carton	62756-008-60
1400 mg in 140 mL	1 single-dose bag per carton	62756-102-60
1500 mg in 150 mL	1 single-dose bag per carton	62756-219-60
1600 mg in 160 mL	1 single-dose bag per carton	62756-321-60
1700 mg in 170 mL	1 single-dose bag per carton	62756-438-60
1800 mg in 180 mL	1 single-dose bag per carton	62756-533-60
1900 mg in 190 mL	1 single-dose bag per carton	62756-614-60
2000 mg in 200 mL	1 single-dose bag per carton	62756-746-60
2200 mg in 220 mL	1 single-dose bag per carton	62756-974-60

16.2 Storage and Handling

INFUGEM is a cytotoxic drug. Follow applicable special handling and disposal procedures.¹

Unopened infusion bags of INFUGEM are stable until the expiration date indicated on the package when stored at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). [see USP Controlled Room Temperature].

Do not freeze as crystallization can occur.

17 PATIENT COUNSELING INFORMATION

Myelosuppression

Advise patients of the risks of myelosuppression and instruct them to immediately contact their healthcare provider should any signs of infection develop, including fever or if bleeding or symptoms of anemia occur [see *Warnings and Precautions* (5.2)].

Pulmonary Toxicity

Advise patients of the risks of pulmonary toxicity including respiratory failure and death. Instruct patients to immediately contact their healthcare provider for development of shortness of breath, wheezing, or cough [see *Warnings and Precautions* (5.3)].

Hemolytic-Uremic Syndrome and Renal Failure

Advise patients of the risks of hemolytic-uremic syndrome and associated renal failure. Instruct patients to

immediately contact their healthcare provider for changes in the color or volume of urine output or for increased bruising or bleeding [*see Warnings and Precautions (5.4)*].

Hepatic Toxicity

Advise patients of the risks of hepatic toxicity including liver failure and death. Instruct patients to immediately contact their healthcare provider for signs of jaundice or for pain/tenderness in the right upper abdominal quadrant [*see Warnings and Precautions (5.5)*].

Embryo-Fetal Toxicity

Advise females and males of reproductive potential that INFUGEM can cause fetal harm when administered to a pregnant woman. Advise females of reproductive potential to use effective contraception during treatment with and for 6 months after the final dose of INFUGEM. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with and for 3 months after the final dose of INFUGEM [*see Warnings and Precaution (5.6), Use in Specific Populations (8.1), (8.3)*].

Lactation

Advise women not to breastfeed during treatment with and for at least one week after the last dose of INFUGEM [*see Use in Specific Populations (8.2)*].

Fertility Effects

Advise males of reproductive potential of the potential for reduced fertility with INFUGEM use [*see Use in Specific Populations (8.3)*].

Distributed by:

Sun Pharmaceutical Industries, Inc.

Cranbury, NJ 08512

Manufactured by:

Sun Pharmaceutical Ind. Ltd.

Halol-Baroda Highway,

Halol-389 350, Gujarat, India



INSTRUCTIONS FOR USE:

INFUGEM (gemcitabine in 0.9% sodium chloride injection), 10 mg/mL

These instructions contain information on how to:

- Select the Correct INFUGEM Bag(s)
- Spike the Infusion Bag

INFUGEM for intravenous use is a clear, colorless, pre-mixed sterile solution that is available in single-dose ready to infuse bags. Do NOT remove or add medication. The bag closures are tamper evident and do not allow contamination.

Understanding the INFUGEM Dose Ranges

INFUGEM (gemcitabine) in 0.9% Sodium Chloride Injection (10 mg/mL) provides predefined dose ranges (in premixed bags) that correspond to BSA-calculated and prescribed doses. Select the INFUGEM premixed bag(s) that allow for a variance of up to 5% of the BSA-calculated dose as described in Table 1 and Table 2.

⚠ Use another formulation of gemcitabine for patients that require a dose that is less than those listed in the tables below (i.e. <1150 mg).

Selecting the Correct INFUGEM INFUSION Bag(s)

1. Use Table 1 for Gemcitabine doses of 1000 mg/m² (non-small cell lung cancer, ovarian cancer, pancreatic cancer).
Use Table 2 for Gemcitabine doses of 1250 mg/m² (breast cancer and non-small cell lung cancer).

2. Identify the appropriate INFUGEM Infusion Bag(s)* based on the BSA-calculated dose range.

* Combinations of bags listed below are suggested combinations. Other possible combinations of bags can be used to reach the appropriate dose.

⚠ Make sure to select the correct target dose table in order to determine the appropriate INFUGEM infusion bag(s).

⚠ Based on the calculated dose, one or two INFUGEM bags may be needed.

How to Spike the INFUGEM Infusion Bag(s)

To achieve the prescribed dose two bags can be administered sequentially.

- Inspect the integrity of the overwrap before removing the overwrap pouch. Do not use the product if the overwrap has been previously opened or damaged.
- Remove the overwrap pouch by tearing at the notch at the bottom of the overwrap and pulling across the infusion bag. (Figure A, B and C)




FIGURE A FIGURE B FIGURE C

- Check for leaks by squeezing the bag firmly. Inspect bag and contents for damage, discoloration or particulate matter.
- Break the tamper evident cap on the infusion port by applying pressure on one side with thumb. (Figure E) matter. (Figure D)




FIGURE D FIGURE E

- Using aseptic technique, remove the cover from the spike of the infusion set. (Figure F)
- Holding the bag with the port side up, insert the spike straight down into the infusion port. (Figure G)




FIGURE F FIGURE G

- Twist and push the spike through the diaphragm. (Figure H)
- Please follow your institution's procedures for administration and disposal. (Figure H)

NOTE: Follow the instructions for use for the infusion set.




FIGURE H

Distributed by:
Sun Pharmaceutical Industries, Inc.
Cranbury, NJ 08512

Manufactured by:
Sun Pharmaceutical Industries Ltd.
Halol-Baroda Highway,
Halol-389 350, Gujarat, India.

For more information, call 1-800-818-4555.
This Instructions for Use has been approved by the U.S. Food and Drug Administration.

RJPI0541
Revised: 07/2018


Use Table 1 to select the appropriate INFUGEM bag(s) for a patient who has been prescribed a target dose of 1000 mg/m².

Table 1: Gemcitabine Dose = 1000 mg/m²
(Non-Small Cell Lung Cancer, Ovarian Cancer, Pancreatic Cancer)

BSA Range (m ²)	Calculated Dose Range	Use this INFUGEM (gemcitabine) Infusion Bag(s)	BSA Range (m ²)	Calculated Dose Range	Use this INFUGEM (gemcitabine) Infusion Bag(s)
1.15 to 1.25	1150 mg to 1254 mg	1200 mg 1200 mg per 120 mL (10 mg/mL)	2.31 to 2.45	2305 mg to 2454 mg	2400 mg (1200 mg bag + 1200 mg per 120 mL (10 mg/mL))
1.26 to 1.35	1255 mg to 1354 mg	1300 mg 1300 mg per 130 mL (10 mg/mL)	2.46 to 2.55	2455 mg to 2554 mg	2500 mg (1300 mg bag + 1200 mg per 120 mL (10 mg/mL))
1.36 to 1.45	1355 mg to 1454 mg	1400 mg 1400 mg per 140 mL (10 mg/mL)	2.56 to 2.64	2555 mg to 2644 mg	2600 mg (1300 mg bag + 1300 mg per 130 mL (10 mg/mL))
1.46 to 1.55	1455 mg to 1554 mg	1500 mg 1500 mg per 150 mL (10 mg/mL)	2.65 to 2.74	2645 mg to 2734 mg	2700 mg (1500 mg bag + 1200 mg per 120 mL (10 mg/mL))
1.56 to 1.65	1555 mg to 1654 mg	1600 mg 1600 mg per 160 mL (10 mg/mL)	2.75 to 2.84	2735 mg to 2824 mg	2800 mg (1600 mg bag + 1200 mg per 120 mL (10 mg/mL))
1.66 to 1.75	1655 mg to 1754 mg	1700 mg 1700 mg per 170 mL (10 mg/mL)	2.85 to 2.94	2825 mg to 2914 mg	2900 mg (1700 mg bag + 1200 mg per 120 mL (10 mg/mL))
1.76 to 1.85	1755 mg to 1854 mg	1800 mg 1800 mg per 180 mL (10 mg/mL)	2.95 to 3.04	2915 mg to 3004 mg	3000 mg (1800 mg bag + 1200 mg per 120 mL (10 mg/mL))
1.86 to 1.95	1855 mg to 1954 mg	1900 mg 1900 mg per 190 mL (10 mg/mL)	3.05 to 3.14	3005 mg to 3094 mg	3100 mg (1900 mg bag + 1200 mg per 120 mL (10 mg/mL))
1.96 to 2.05	1955 mg to 2054 mg	2000 mg 2000 mg per 200 mL (10 mg/mL)	3.15 to 3.24	3095 mg to 3184 mg	3200 mg (2000 mg bag + 1200 mg per 120 mL (10 mg/mL))
2.06 to 2.15	2055 mg to 2154 mg	2100 mg 2100 mg per 210 mL (10 mg/mL)	3.25 to 3.34	3185 mg to 3274 mg	3300 mg (2100 mg bag + 1200 mg per 120 mL (10 mg/mL))
2.16 to 2.25	2155 mg to 2254 mg	2200 mg 2200 mg per 220 mL (10 mg/mL)	3.35 to 3.44	3275 mg to 3364 mg	3400 mg (2200 mg bag + 1200 mg per 120 mL (10 mg/mL))
2.26 to 2.35	2255 mg to 2354 mg	2300 mg 2300 mg per 230 mL (10 mg/mL)	3.45 to 3.54	3365 mg to 3454 mg	3500 mg (2300 mg bag + 1200 mg per 120 mL (10 mg/mL))
2.36 to 2.45	2355 mg to 2454 mg	2400 mg 2400 mg per 240 mL (10 mg/mL)	3.55 to 3.64	3455 mg to 3544 mg	3600 mg (2400 mg bag + 1200 mg per 120 mL (10 mg/mL))
2.46 to 2.55	2455 mg to 2554 mg	2500 mg 2500 mg per 250 mL (10 mg/mL)	3.65 to 3.74	3545 mg to 3634 mg	3700 mg (2500 mg bag + 1200 mg per 120 mL (10 mg/mL))
2.56 to 2.65	2555 mg to 2654 mg	2600 mg 2600 mg per 260 mL (10 mg/mL)	3.75 to 3.84	3635 mg to 3724 mg	3800 mg (2600 mg bag + 1200 mg per 120 mL (10 mg/mL))
2.66 to 2.75	2655 mg to 2754 mg	2700 mg 2700 mg per 270 mL (10 mg/mL)	3.85 to 3.94	3725 mg to 3814 mg	3900 mg (2700 mg bag + 1200 mg per 120 mL (10 mg/mL))
2.76 to 2.85	2755 mg to 2854 mg	2800 mg 2800 mg per 280 mL (10 mg/mL)	3.95 to 4.04	3815 mg to 3904 mg	4000 mg (2800 mg bag + 1200 mg per 120 mL (10 mg/mL))
2.86 to 2.95	2855 mg to 2954 mg	2900 mg 2900 mg per 290 mL (10 mg/mL)	4.05 to 4.14	3905 mg to 3994 mg	4100 mg (2900 mg bag + 1200 mg per 120 mL (10 mg/mL))
2.96 to 3.05	2955 mg to 3054 mg	3000 mg 3000 mg per 300 mL (10 mg/mL)	4.15 to 4.24	3995 mg to 4084 mg	4200 mg (3000 mg bag + 1200 mg per 120 mL (10 mg/mL))
3.06 to 3.15	3055 mg to 3154 mg	3100 mg 3100 mg per 310 mL (10 mg/mL)	4.25 to 4.34	4085 mg to 4174 mg	4300 mg (3100 mg bag + 1200 mg per 120 mL (10 mg/mL))
3.16 to 3.25	3155 mg to 3254 mg	3200 mg 3200 mg per 320 mL (10 mg/mL)	4.35 to 4.44	4175 mg to 4264 mg	4400 mg (3200 mg bag + 1200 mg per 120 mL (10 mg/mL))
3.26 to 3.35	3255 mg to 3354 mg	3300 mg 3300 mg per 330 mL (10 mg/mL)	4.45 to 4.54	4265 mg to 4354 mg	4500 mg (3300 mg bag + 1200 mg per 120 mL (10 mg/mL))
3.36 to 3.45	3355 mg to 3454 mg	3400 mg 3400 mg per 340 mL (10 mg/mL)	4.55 to 4.64	4355 mg to 4444 mg	4600 mg (3400 mg bag + 1200 mg per 120 mL (10 mg/mL))
3.46 to 3.55	3455 mg to 3554 mg	3500 mg 3500 mg per 350 mL (10 mg/mL)	4.65 to 4.74	4445 mg to 4534 mg	4700 mg (3500 mg bag + 1200 mg per 120 mL (10 mg/mL))
3.56 to 3.65	3555 mg to 3654 mg	3600 mg 3600 mg per 360 mL (10 mg/mL)	4.75 to 4.84	4535 mg to 4624 mg	4800 mg (3600 mg bag + 1200 mg per 120 mL (10 mg/mL))
3.66 to 3.75	3655 mg to 3754 mg	3700 mg 3700 mg per 370 mL (10 mg/mL)	4.85 to 4.94	4625 mg to 4714 mg	4900 mg (3700 mg bag + 1200 mg per 120 mL (10 mg/mL))
3.76 to 3.85	3755 mg to 3854 mg	3800 mg 3800 mg per 380 mL (10 mg/mL)	4.95 to 5.04	4715 mg to 4804 mg	5000 mg (3800 mg bag + 1200 mg per 120 mL (10 mg/mL))
3.86 to 3.95	3855 mg to 3954 mg	3900 mg 3900 mg per 390 mL (10 mg/mL)	5.05 to 5.14	4805 mg to 4894 mg	5100 mg (3900 mg bag + 1200 mg per 120 mL (10 mg/mL))
3.96 to 4.05	3955 mg to 4054 mg	4000 mg 4000 mg per 400 mL (10 mg/mL)	5.15 to 5.24	4895 mg to 4984 mg	5200 mg (4000 mg bag + 1200 mg per 120 mL (10 mg/mL))
4.06 to 4.15	4055 mg to 4154 mg	4100 mg 4100 mg per 410 mL (10 mg/mL)	5.25 to 5.34	4985 mg to 5074 mg	5300 mg (4100 mg bag + 1200 mg per 120 mL (10 mg/mL))
4.16 to 4.25	4155 mg to 4254 mg	4200 mg 4200 mg per 420 mL (10 mg/mL)	5.35 to 5.44	5075 mg to 5164 mg	5400 mg (4200 mg bag + 1200 mg per 120 mL (10 mg/mL))
4.26 to 4.35	4255 mg to 4354 mg	4300 mg 4300 mg per 430 mL (10 mg/mL)	5.45 to 5.54	5165 mg to 5254 mg	5500 mg (4300 mg bag + 1200 mg per 120 mL (10 mg/mL))
4.36 to 4.45	4355 mg to 4454 mg	4400 mg 4400 mg per 440 mL (10 mg/mL)	5.55 to 5.64	5255 mg to 5344 mg	5600 mg (4400 mg bag + 1200 mg per 120 mL (10 mg/mL))
4.46 to 4.55	4455 mg to 4554 mg	4500 mg 4500 mg per 450 mL (10 mg/mL)	5.65 to 5.74	5345 mg to 5434 mg	5700 mg (4500 mg bag + 1200 mg per 120 mL (10 mg/mL))
4.56 to 4.65	4555 mg to 4654 mg	4600 mg 4600 mg per 460 mL (10 mg/mL)	5.75 to 5.84	5435 mg to 5524 mg	5800 mg (4600 mg bag + 1200 mg per 120 mL (10 mg/mL))
4.66 to 4.75	4655 mg to 4754 mg	4700 mg 4700 mg per 470 mL (10 mg/mL)	5.85 to 5.94	5525 mg to 5614 mg	5900 mg (4700 mg bag + 1200 mg per 120 mL (10 mg/mL))
4.76 to 4.85	4755 mg to 4854 mg	4800 mg 4800 mg per 480 mL (10 mg/mL)	5.95 to 6.04	5615 mg to 5704 mg	6000 mg (4800 mg bag + 1200 mg per 120 mL (10 mg/mL))
4.86 to 4.95	4855 mg to 4954 mg	4900 mg 4900 mg per 490 mL (10 mg/mL)	6.05 to 6.14	5705 mg to 5794 mg	6100 mg (4900 mg bag + 1200 mg per 120 mL (10 mg/mL))
4.96 to 5.05	4955 mg to 5054 mg	5000 mg 5000 mg per 500 mL (10 mg/mL)	6.15 to 6.24	5795 mg to 5884 mg	6200 mg (5000 mg bag + 1200 mg per 120 mL (10 mg/mL))
5.06 to 5.15	5055 mg to 5154 mg	5100 mg 5100 mg per 510 mL (10 mg/mL)	6.25 to 6.34	5885 mg to 5974 mg	6300 mg (5100 mg bag + 1200 mg per 120 mL (10 mg/mL))
5.16 to 5.25	5155 mg to 5254 mg	5200 mg 5200 mg per 520 mL (10 mg/mL)	6.35 to 6.44	5975 mg to 6064 mg	6400 mg (5200 mg bag + 1200 mg per 120 mL (10 mg/mL))
5.26 to 5.35	5255 mg to 5354 mg	5300 mg 5300 mg per 530 mL (10 mg/mL)	6.45 to 6.54	6065 mg to 6154 mg	6500 mg (5300 mg bag + 1200 mg per 120 mL (10 mg/mL))
5.36 to 5.45	5355 mg to 5454 mg	5400 mg 5400 mg per 540 mL (10 mg/mL)	6.55 to 6.64	6155 mg to 6244 mg	6600 mg (5400 mg bag + 1200 mg per 120 mL (10 mg/mL))
5.46 to 5.55	5455 mg to 5554 mg	5500 mg 5500 mg per 550 mL (10 mg/mL)	6.65 to 6.74	6245 mg to 6334 mg	6700 mg (5500 mg bag + 1200 mg per 120 mL (10 mg/mL))
5.56 to 5.65	5555 mg to 5654 mg	5600 mg 5600 mg per 560 mL (10 mg/mL)	6.75 to 6.84	6335 mg to 6424 mg	6800 mg (5600 mg bag + 1200 mg per 120 mL (10 mg/mL))
5.66 to 5.75	5655 mg to 5754 mg	5700 mg 5700 mg per 570 mL (10 mg/mL)	6.85 to 6.94	6425 mg to 6514 mg	6900 mg (5700 mg bag + 1200 mg per 120 mL (10 mg/mL))
5.76 to 5.85	5755 mg to 5854 mg	5800 mg 5800 mg per 580 mL (10 mg/mL)	6.95 to 7.04	6515 mg to 6604 mg	7000 mg (5800 mg bag + 1200 mg per 120 mL (10 mg/mL))
5.86 to 5.95	5855 mg to 5954 mg	5900 mg 5900 mg per 590 mL (10 mg/mL)	7.05 to 7.14	6605 mg to 6694 mg	7100 mg (5900 mg bag + 1200 mg per 120 mL (10 mg/mL))
5.96 to 6.05	5955 mg to 6054 mg	6000 mg 6000 mg per 600 mL (10 mg/mL)	7.15 to 7.24	6695 mg to 6784 mg	7200 mg (6000 mg bag + 1200 mg per 120 mL (10 mg/mL))
6.06 to 6.15	6055 mg to 6154 mg	6100 mg 6100 mg per 610 mL (10 mg/mL)	7.25 to 7.34	6785 mg to 6874 mg	7300 mg (6100 mg bag + 1200 mg per 120 mL (10 mg/mL))
6.16 to 6.25	6155 mg to 6254 mg	6200 mg 6200 mg per 620 mL (10 mg/mL)	7.35 to 7.44	6875 mg to 6964 mg	7400 mg (6200 mg bag + 1200 mg per 120 mL (10 mg/mL))
6.26 to 6.35	6255 mg to 6354 mg	6300 mg 6300 mg per 630 mL (10 mg/mL)	7.45 to 7.54	6965 mg to 7054 mg	7500 mg (6300 mg bag + 1200 mg per 120 mL (10 mg/mL))
6.36 to 6.45	6355 mg to 6454 mg	6400 mg 6400 mg per 640 mL (10 mg/mL)	7.55 to 7.64	7055 mg to 7144 mg	7600 mg (6400 mg bag + 1200 mg per 120 mL (10 mg/mL))
6.46 to 6.55	6455 mg to 6554 mg	6500 mg 6500 mg per 650 mL (10 mg/mL)	7.65 to 7.74	7145 mg to 7234 mg	7700 mg (6500 mg bag + 1200 mg per 120 mL (10 mg/mL))
6.56 to 6.65	6555 mg to 6654 mg	6600 mg 6600 mg per 660 mL (10 mg/mL)	7.75 to 7.84	7235 mg to 7324 mg	7800 mg (6600 mg bag + 1200 mg per 120 mL (10 mg/mL))
6.66 to 6.75	6655 mg to 6754 mg	6700 mg 6700 mg per 670 mL (10 mg/mL)	7.85 to 7.94	7325 mg to 7414 mg	7900 mg (6700 mg bag + 1200 mg per 120 mL (10 mg/mL))
6.76 to 6.85	6755 mg to 6854 mg	6800 mg 6800 mg per 680 mL (10 mg/mL)	7.95 to 8.04	7415 mg to 7504 mg	8000 mg (6800 mg bag + 1200 mg per 120 mL (10 mg/mL))
6.86 to 6.95	6855 mg to 6954 mg	6900 mg 6900 mg per 690 mL (10 mg/mL)	8.05 to 8.14	7505 mg to 7594 mg	8100 mg (6900 mg bag + 1200 mg per 120 mL (10 mg/mL))
6.96 to 7.05	6955 mg to 7054 mg	7000 mg 7000 mg per 700 mL (10 mg/mL)	8.15 to 8.24	7595 mg to 7684 mg	8200 mg (7000 mg bag + 1200 mg per 120 mL (10 mg/mL))
7.06 to 7.15	7055 mg to 7154 mg	7100 mg 7100 mg per 710 mL (10 mg/mL)	8.25 to 8.34	7685 mg to 7774 mg	8300 mg (7100 mg bag + 1200 mg per 120 mL (10 mg/mL))
7.16 to 7.25	7155 mg to 7254 mg	7200 mg 7200 mg per 720 mL (10 mg/mL)	8.35 to 8.44	7775 mg to 7864 mg	8400 mg (7200 mg bag + 1200 mg per 120 mL (10 mg/mL))
7.26 to 7.35	7255 mg to 7354 mg	7300 mg 7300 mg per 730 mL (10 mg/mL)	8.45 to 8.54	7865 mg to 7954 mg	8500 mg (7300 mg bag + 1200 mg per 120 mL (10 mg/mL))
7.36 to 7.45	7355 mg to 7454 mg	7400 mg 7400 mg per 740 mL (10 mg/mL)	8.55 to 8.64	7955 mg to 8044 mg	8600 mg (7400 mg bag + 1200 mg per 120 mL (10 mg/mL))
7.46 to 7.55	7455 mg to 7554 mg	7500 mg 7500 mg per 750 mL (10 mg/mL)	8.65 to 8.74	8045 mg to 8134 mg	8700 mg (7500 mg bag + 1200 mg per 120 mL (10 mg/mL))
7.56 to 7.65	7555 mg to 7654 mg	7600 mg 7600 mg per 760 mL (10 mg/mL)	8.75 to 8.84	8135 mg to 8224 mg	8800 mg (7600 mg bag + 1200 mg per 120 mL (10 mg/mL))
7.66 to 7.75	7655 mg to 7754 mg	7700 mg 7700 mg per 770 mL (10 mg/mL)	8.85 to 8.94	8225 mg to 8314 mg	8900 mg (7700 mg bag + 1200 mg per 120 mL (10 mg/mL))
7.76 to 7.85	7755 mg to 7854 mg	7800 mg 7800 mg per 780 mL (10 mg/mL)	8.95 to 9.04	8315 mg to 8404 mg	9000 mg (7800 mg bag + 1200 mg per 120 mL (10 mg/mL))
7.86 to 7.95	7855 mg to 7954 mg	7900 mg 7900 mg per 790 mL (10 mg/mL)	9.05 to 9.14	8405 mg to 8494 mg	9100 mg (7900 mg bag + 1200 mg per 120 mL (10 mg/mL))
7.96 to 8.05	7955 mg to 8054 mg	8000 mg 8000 mg per 800 mL (10 mg/mL)	9.15 to 9.24	8495 mg to 8584 mg	9200 mg (8000 mg bag + 1200 mg per 120 mL (10 mg/mL))
8.06 to 8.15	8055 mg to 8154 mg	8100 mg 8			

Table of Content

Final Labels	2
Bag Labels.....	2
1,200 mg/120 mL.....	2
1,300 mg/130 mL.....	3
1,400 mg/140 mL.....	4
1,500 mg/150 mL.....	5
1,600 mg/160 mL.....	6
1,700 mg/170 mL.....	7
1,800 mg/180 mL.....	8
1,900 mg/190 mL.....	9
2,000 mg/200 mL.....	10
2,200 mg/220 mL.....	11
Pouch Labels.....	12
1,200 mg/120 mL.....	12
1,300 mg/130 mL.....	14
1,400 mg/140 mL.....	16
1,500 mg/150 mL.....	18
1,600 mg/160 mL.....	20
1,700 mg/170 mL.....	22
1,800 mg/180 mL.....	24
1,900 mg/190 mL.....	26
2,000 mg/200 mL.....	28
2,200 mg/220 mL.....	30
Carton Labels	32
1,200 mg/120 mL.....	32
1,300 mg/130 mL.....	33
1,400 mg/140 mL.....	34
1,500 mg/150 mL.....	35
1,600 mg/160 mL.....	36
1,700 mg/170 mL.....	37
1,800 mg/180 mL.....	38
1,900 mg/190 mL.....	39
2,000 mg/200 mL.....	40
2,200 mg/220 mL.....	41




Size : 85x227 mm




(b) (4)



Size : 85x227 mm








(b) (4)





(b) (4)



227 mm

7.5mm

7.5mm

43 mm



122 mm

62 mm

70 mm

85 mm

Size : 85x227 mm



Size : 85x227 mm




Size : 85x227 mm






Drawing No: 630 - V1 Size : 115x200 mm

Sealed Bottom



* This will be printed on pouch without any background



Sealed Bottom

75 mm

93 mm

175 mm

330 mm



80 mm

30 mm

30 mm

Open Top
135 mm

TO OPEN: Tear at notch at bottom of overwrap.
Do not remove unit from overwrap until ready to use.
Do not use if overwrap has been previously open or damaged.
 The overwrap is a moisture and oxygen barrier.

NDC 62756-008-60 Rx only

Infugem™
Gemcitabine
 in 0.9% sodium chloride injection

1,300 mg per 130 mL
 (10 mg/mL)

Caution: Cytotoxic Agent For Intravenous Use Only

Contains one sterile single-dose bag.

Each 100 mL contains:
 Gemcitabine 1,000 mg
 equivalent to Gemcitabine HCl USP 1,138 mg
 Sodium Chloride 900 mg
 Hydrochloric acid and
 Sodium hydroxide added to adjust pH. q.s.

Premixed Intravenous Solution
Usual Dosage: See prescribing information.
Do NOT remove or add medication to IV bag.
 Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). [see USP Controlled Room Temperature].
 Do not freeze.


 SUN
 PHARMA


 OTHER


 62756 00860 1


Distributed by:
Sun Pharmaceutical Industries, Inc.
 Cranbury, NJ 08512

PJPC008
 GUJ/DRUGS/28-D/LVP-6

Manufactured by:
Sun Pharmaceutical Industries Ltd., India

Lot No.:
 EXP.:

* This will be printed on pouch without any background



NDC 62756-008-60

Rx only

Infugem™
Gemcitabine
 in 0.9% sodium chloride injection

1,300 mg per 130 mL
 (10 mg/mL)

Caution: Cytotoxic Agent

For Intravenous Use Only

Contains one sterile single-dose bag.

PJPC0008

Sealed Bottom

75 mm

93 mm

TO OPEN: Tear at notch at bottom of overwrap.
Do not remove unit from overwrap until ready to use.
Do not use if overwrap has been previously open or damaged.
 The overwrap is a moisture and oxygen barrier.

NDC 62756-102-60 Rx only

InfugemTM
Gemcitabine
 in 0.9% sodium chloride injection

1,400 mg per 140 mL
 (10 mg/mL)

Caution: Cytotoxic Agent For Intravenous Use Only

Contains one sterile single-dose bag.

Each 100 mL contains:
 Gemcitabine 1,000 mg
 equivalent to Gemcitabine HCl USP 1,138 mg
 Sodium Chloride 900 mg
 Hydrochloric acid and
 Sodium hydroxide added to adjust pH. q.s.

Premixed Intravenous Solution
Usual Dosage: See prescribing information.
Do NOT remove or add medication to IV bag.
 Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). [see USP Controlled Room Temperature].
 Do not freeze.


 SUN PHARMA


 OTHER


 6275610260 6
 N 3

Distributed by:
Sun Pharmaceutical Industries, Inc.
 Cranbury, NJ 08512

PJPC0009
 GUJ/DRUGS/28-D/LVP-6

Manufactured by:
Sun Pharmaceutical Industries Ltd., India

Lot No.:
 EXP.:

175 mm

330 mm


80 mm

30 mm

30 mm

Open Top
135 mm

*** This will be printed on pouch without any background**



Sealed Bottom

75 mm

93 mm

175 mm

330 mm

(b) (4)

TO OPEN: Tear at notch at bottom of overwrap.
Do not remove unit from overwrap until ready to use.
Do not use if overwrap has been previously open or damaged.
 The overwrap is a moisture and oxygen barrier.

NDC 62756-219-60 Rx only

InfugemTM
Gemcitabine
 in 0.9% sodium chloride injection

1,500 mg per 150 mL
 (10 mg/mL)

Caution: Cytotoxic Agent For Intravenous Use Only

Contains one sterile single-dose bag.

Each 100 mL contains:
 Gemcitabine 1,000 mg
 equivalent to Gemcitabine HCl USP 1,138 mg
 Sodium Chloride 900 mg
 Hydrochloric acid and
 Sodium hydroxide added to adjust pH. q.s.

Premixed Intravenous Solution
Usual Dosage: See prescribing information.
Do NOT remove or add medication to IV bag.
 Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). [see USP Controlled Room Temperature].
 Do not freeze.


 SUN
 PHARMA


 OTHER


 62756 21960 1

Distributed by:
Sun Pharmaceutical Industries, Inc.
 Cranbury, NJ 08512

PJPC0010
 GUJ/DRUGS/28-D/LVP-6

Manufactured by:
Sun Pharmaceutical Industries Ltd., India

Lot No.:
 EXP.:

80 mm

30 mm

30 mm

Open Top
135 mm

*** This will be printed on pouch without any background**



NDC 62756-219-60

Rx only

Infugem™
Gemcitabine
 in 0.9% sodium chloride injection


1,500 mg per 150 mL
 (10 mg/mL)

Caution: Cytotoxic Agent


For Intravenous Use Only


Contains one sterile single-dose bag.

PJPC0010



*** This will be printed on pouch without any background**





(b) (4)

75 mm

80 mm

30 mm

30 mm

Sealed Bottom

93 mm

175 mm

330 mm

Open Top
135 mm

TO OPEN: Tear at notch at bottom of overwrap.
Do not remove unit from overwrap until ready to use.
Do not use if overwrap has been previously open or damaged.
 The overwrap is a moisture and oxygen barrier.

NDC 62756-438-60 Rx only

InfugemTM
Gemcitabine
 in 0.9% sodium chloride injection

1,700 mg per 170 mL
 (10 mg/mL)

Caution: Cytotoxic Agent **For Intravenous Use Only**

Contains one sterile single-dose bag.

Each 100 mL contains:
 Gemcitabine 1,000 mg
 equivalent to Gemcitabine HCl USP 1,138 mg
 Sodium Chloride 900 mg
 Hydrochloric acid and
 Sodium hydroxide added to adjust pH. q.s.

Premixed Intravenous Solution
Usual Dosage: See prescribing information.
Do NOT remove or add medication to IV bag.
 Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). [see USP Controlled Room Temperature].
 Do not freeze.


 SUN
 PHARMA


 OTHER


 6275643860 6


Distributed by:
Sun Pharmaceutical Industries, Inc.
 Cranbury, NJ 08512


PJPC0012
 GUJ/DRUGS/28-D/LVP-6

Lot No.:
 EXP.:


Manufactured by:
Sun Pharmaceutical Industries Ltd., India


* This will be printed on pouch without any background








*** This will be printed on pouch without any background**






*** This will be printed on pouch without any background**







*** This will be printed on pouch without any background**




Sealed Bottom



* This will be printed on pouch without any background



Base




Case Outer




Size: 344x139.5x27 mm
ANDA-US

Case Inner



D1706020_Gemcitabine_120ml_to_160ml_Box_V1_KLD

Base




Case Outer




Size: 344x139.5x27 mm
ANDA-US


Case Inner



D1706020_Gemcitabine_120ml_to_160ml_Box_V1_KLD




Size: 344x139.5x34 mm
ANDA-US



D1706021_Gemcitabine_170ml_to_200ml_Box_V1_KLD

Base




Case Outer




Size: 344x139.5x34 mm
ANDA-US


Case Inner



D1706021_Gemcitabine_170ml_to_200ml_Box_V1_KLD




Size: 344x139.5x34 mm
ANDA-US



D1706021_Gemcitabine_170ml_to_200ml_Box_V1_KLD

Base




Case Outer



Size: 344x169.5x26 mm
ANDA-US

Case Inner



D1706022_Gemcitabine_220ml_Box_V1_KLD

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

JOSEPH E GOOTENBERG
07/16/2018