

**CENTER FOR DRUG EVALUATION AND  
RESEARCH**

***APPLICATION NUMBER:***

**210331Orig1s000**

**LABELING**

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use YUTIQ™ safely and effectively. See full prescribing information for YUTIQ.

**YUTIQ™ (fluocinolone acetonide intravitreal implant) 0.18 mg, for intravitreal injection**

**Initial U.S. Approval: 1963**

**-----INDICATIONS AND USAGE-----**

YUTIQ contains a corticosteroid and is indicated for the treatment of chronic non-infectious uveitis affecting the posterior segment of the eye. (1)

**-----DOSAGE AND ADMINISTRATION-----**

- For ophthalmic intravitreal injection. (0)
- The intravitreal injection procedure should be carried out under aseptic conditions. (0)
- Following the intravitreal injection, patients should be monitored for elevation in intraocular pressure and for endophthalmitis. (0)

**-----DOSAGE FORMS AND STRENGTHS-----**

Non-bioerodible intravitreal implant containing 0.18 mg fluocinolone acetonide in a drug delivery system. (3)

**-----CONTRAINDICATIONS-----**

- Ocular or periocular infections (0)
- Hypersensitivity (02)

**-----WARNINGS AND PRECAUTIONS-----**

- Intravitreal injections have been associated with endophthalmitis, eye inflammation, increased intraocular pressure, and retinal detachments. Patients should be monitored following the injection. (0)
- Use of corticosteroids may produce posterior subcapsular cataracts, increased intraocular pressure, glaucoma, and may enhance the establishment of secondary ocular infections due to bacteria, fungi, or viruses. (0)
- The implant may migrate into the anterior chamber if the posterior lens capsule is not intact. (0)

**-----ADVERSE REACTIONS-----**

In controlled studies, the most common adverse reactions reported were cataract development and increases in intraocular pressure. (0)

**To report SUSPECTED ADVERSE REACTIONS, contact EyePoint Pharmaceuticals US, Inc. at 1-888-812-6198 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**See 17 for PATIENT COUNSELING INFORMATION**

**Revised: 10/2018**

**FULL PRESCRIBING INFORMATION: CONTENTS\*****1 INDICATIONS AND USAGE****2 DOSAGE AND ADMINISTRATION**

2.1 General Dosing Information

2.2 Administration

**3 DOSAGE FORMS AND STRENGTHS****4 CONTRAINDICATIONS**

4.1 Ocular or Periocular Infections

4.2 Hypersensitivity

**5 WARNINGS AND PRECAUTIONS**

5.1 Intravitreal Injection-related Effects

5.2 Steroid-related Effects

5.3 Risk of Implant Migration

**6 ADVERSE REACTIONS**

6.1 Clinical Studies Experience

**8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

**11 DESCRIPTION****12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

**13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**14 CLINICAL STUDIES****16 HOW SUPPLIED/STORAGE AND HANDLING****17 PATIENT COUNSELING INFORMATION**

\*Sections or subsections omitted from the full prescribing information are not listed.

## FULL PRESCRIBING INFORMATION

### 1. INDICATIONS AND USAGE

YUTIQ™ (fluocinolone acetonide intravitreal implant) 0.18 mg is indicated for the treatment of chronic non-infectious uveitis affecting the posterior segment of the eye.

### 2. DOSAGE AND ADMINISTRATION

#### 2.1 General Dosing Information

For ophthalmic intravitreal injection.

#### 2.2 Administration

The intravitreal injection procedure should be carried out under aseptic conditions, which include use of sterile gloves, a sterile drape, a sterile caliper, and a sterile eyelid speculum (or equivalent). Adequate anesthesia and a broad-spectrum microbicide should be given prior to the injection.

The injection procedure for YUTIQ is as follows:

1. Just prior to injection, administer topical and/or subconjunctival anesthesia at the injection site (inferotemporal quadrant recommended).
2. Administer 2-3 drops of a broad-spectrum microbicide into the lower fornix. The lids may be scrubbed with cotton-tipped applicators soaked with a broad-spectrum microbicide. Place a sterile lid speculum. Have the patient look up and apply additional microbicide solution to the injection site. Allow 30-60 seconds for the topical antiseptic to dry prior to injection of YUTIQ.
3. Optimal placement of YUTIQ is inferior to the optic disc and posterior to the equator of the eye. Measure 4 millimeters inferotemporal from the limbus with the aid of callipers for point of entry into the sclera.
4. Using sterile procedure, open the sterile foil pouch containing YUTIQ.
5. Remove the YUTIQ applicator from the sterile pouch by grasping the barrel of the applicator; do not grasp the plunger.
6. Remove the black plunger stop from the plunger.
7. Carefully remove the protective cap from the needle and inspect the needle tip to ensure it is not bent.
8. Remove the trombone wire from the distal end of the needle. Prior to injection, keep the applicator tip above the horizontal plane to ensure that the YUTIQ implant does not fall out of the applicator.
9. Gently displace the conjunctiva so that after withdrawing the needle, the conjunctival and scleral needle entry sites will not align. Care should be taken to avoid contact between the needle and the lid margin or lashes.
10. Insert the needle through the conjunctiva and sclera up to the positive stop of the applicator.
11. Depress the plunger at the back of the applicator fully to deliver the YUTIQ implant into the back of the eye.
12. Remove the YUTIQ applicator from the eye and discard in biohazard sharps container.
13. Remove the lid speculum and perform indirect ophthalmoscopy to verify adequate central retinal artery perfusion, absence of any other complications, and to verify the placement of the implant. Scleral depression may enhance visualisation of the implant. Immediate measurement of intraocular pressure (IOP) may be performed at the discretion of the ophthalmologist.

Following the injection, patients should be monitored for change in intraocular pressure and for endophthalmitis. Monitoring may consist of a check for perfusion of the optic nerve head immediately after the injection, tonometry within 30 minutes following the injection, and biomicroscopy between two and seven days following the injection. Patients should be instructed to report without delay any symptoms suggestive of endophthalmitis.

### **3. DOSAGE FORMS AND STRENGTHS**

YUTIQ is a non-bioerodible intravitreal implant in a drug delivery system containing 0.18 mg fluocinolone acetonide, designed to release fluocinolone acetonide at an initial rate of 0.25 mcg/day, and lasting 36 months.

### **4. CONTRAINDICATIONS**

#### **4.1 Ocular or Periocular Infections**

YUTIQ is contraindicated in patients with active or suspected ocular or periocular infections including most viral disease of the cornea and conjunctiva including active epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, varicella, mycobacterial infections and fungal diseases.

#### **4.2 Hypersensitivity**

YUTIQ is contraindicated in patients with known hypersensitivity to any components of this product.

### **5. WARNINGS AND PRECAUTIONS**

#### **5.1 Intravitreal Injection-related Effects**

Intravitreal injections, including those with YUTIQ, have been associated with endophthalmitis, eye inflammation, increased or decreased intraocular pressure, and choroidal or retinal detachments. Hypotony has been observed within 24 hours of injection and has resolved within 2 weeks. Patients should be monitored following the intravitreal injection [see *Patient Counseling Information* (17)].

#### **5.2 Steroid-related Effects**

Use of corticosteroids including YUTIQ may produce posterior subcapsular cataracts, increased intraocular pressure and glaucoma. Use of corticosteroids may enhance the establishment of secondary ocular infections due to bacteria, fungi, or viruses.

Corticosteroids are not recommended to be used in patients with a history of ocular herpes simplex because of the potential for reactivation of the viral infection.

#### **5.3 Risk of Implant Migration**

Patients in whom the posterior capsule of the lens is absent or has a tear are at risk of implant migration into the anterior chamber.

## 6. ADVERSE REACTIONS

### 6.1 Clinical Studies Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse reactions associated with ophthalmic steroids including YUTIQ include cataract formation and subsequent cataract surgery, elevated intraocular pressure, which may be associated with optic nerve damage, visual acuity and field defects, secondary ocular infection from pathogens including herpes simplex, and perforation of the globe where there is thinning of the cornea or sclera.

Studies 1 and 2 were multicenter, randomized, sham injection-controlled, masked trials in which patients with non-infectious uveitis affecting the posterior segment of the eye were treated once with either YUTIQ or sham injection, and then received standard care for the duration of the study. Study 3 was a multicenter, randomized, masked trial in which patients with non-infectious uveitis affecting the posterior segment of the eye were all treated once with YUTIQ, administered by one of two different applicators, and then received standard care for the duration of the study.

Table 1 summarizes data available from studies 1, 2 and 3 through 12 months for study eyes treated with YUTIQ (n=226) or sham injection (n=94). The most common ocular (study eye) and non-ocular adverse reactions are shown in Table 1 and Table 2.

**Table 1: Ocular Adverse Reactions Reported in  $\geq 1\%$  of Subject Eyes and Non-Ocular Adverse Reactions Reported in  $\geq 2\%$  of Patients**


Ocular		
ADVERSE REACTIONS	YUTIQ (N=226 Eyes) n (%)	Sham Injection (N=94 Eyes) n (%)
Cataract <sup>1</sup>	63/113 (56%)	13/56 (23%)
Visual Acuity Reduced	33 ( 15%)	11 (12%)
Macular Edema	25 ( 11%)	33 (35%)
Uveitis	22 ( 10%)	33 (35%)
Conjunctival Hemorrhage	17 ( 8%)	5 ( 5%)
Eye Pain	17 ( 8%)	12 (13%)
Hypotony Of Eye	16 ( 7%)	1 ( 1%)
Anterior Chamber Inflammation	12 ( 5%)	6 ( 6%)
Dry Eye	10 ( 4%)	3 ( 3%)
Vitreous Opacities	9 ( 4%)	8 ( 9%)
Conjunctivitis	9 ( 4%)	5 ( 5%)

<b>Ocular</b>		
<b>ADVERSE REACTIONS</b>	<b>YUTIQ (N=226 Eyes) n (%)</b>	<b>Sham Injection (N=94 Eyes) n (%)</b>
Posterior Capsule Opacification	8 ( 4%)	3 ( 3%)
Ocular Hyperemia	8 ( 4%)	7 ( 7%)
Vitreous Haze	7 ( 3%)	4 ( 4%)
Foreign Body Sensation In Eyes	7 ( 3%)	2 ( 2%)
Vitritis	6 ( 3%)	8 ( 9%)
Vitreous Floaters	6 ( 3%)	5 ( 5%)
Eye Pruritus	6 ( 3%)	5 ( 5%)
Conjunctival Hyperemia	5 ( 2%)	2 ( 2%)
Ocular Discomfort	5 ( 2%)	1 ( 1%)
Macular Fibrosis	5 ( 2%)	2 ( 2%)
Glaucoma	4 ( 2%)	1 ( 1%)
Photopsia	4 ( 2%)	2 ( 2%)
Vitreous Hemorrhage	4 ( 2%)	0
Iridocyclitis	3 ( 1%)	7 ( 7%)
Eye Inflammation	3 ( 1%)	2 ( 2%)
Choroiditis	3 ( 1%)	1 ( 1%)
Eye Irritation	3 ( 1%)	1 ( 1%)
Visual Field Defect	3 ( 1%)	0
Lacrimation Increased	3 ( 1%)	0
<b>Non-ocular</b>		
<b>ADVERSE REACTIONS</b>	<b>YUTIQ (N=214 Patients) n (%)</b>	<b>Sham Injection (N=94 Patients) n (%)</b>
Nasopharyngitis	10 ( 5%)	5 ( 5%)
Hypertension	6 ( 3%)	1 ( 1%)
Arthralgia	5 ( 2%)	1 ( 1%)
1. Includes cataract, cataract subcapsular and lenticular opacities in study eyes that were phakic at baseline. 113 of the 226 YUTIQ study eyes were phakic at baseline; 56 of 94 sham-controlled study eyes were phakic at baseline.		

**Table 2: Summary of Elevated IOP Related Adverse Reactions**

ADVERSE REACTIONS	YUTIQ (N=226 Eyes) n (%)	Sham (N=94 Eyes) n (%)
IOP elevation $\geq$ 10 mmHg from Baseline	50 (22%)	11 (12%)
IOP elevation > 30 mmHg	28 (12%)	3 (3%)
Any IOP-lowering medication	98 (43%)	39 (41%)
Any surgical intervention for elevated IOP	5 (2%)	2 (2%)

**Figure 1: Mean IOP During the Studies**



## 8. USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Adequate and well-controlled studies with YUTIQ have not been conducted in pregnant women to inform drug associated risk. Animal reproduction studies have not been conducted with YUTIQ. It is not known whether YUTIQ can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Corticosteroids have been shown to be teratogenic in laboratory animals when administered systemically at relatively low dosage levels. YUTIQ should be given to a pregnant woman only if the potential benefit justifies the potential risk to the fetus.

All pregnancies have a risk of birth defect, loss, or other adverse outcomes. In the United States general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

## 8.2 Lactation

### Risk Summary

Systemically administered corticosteroids are present in human milk and can suppress growth, interfere with endogenous corticosteroid production. Clinical or nonclinical lactation studies have not been conducted with YUTIQ. It is not known whether intravitreal treatment with YUTIQ could result in sufficient systemic absorption to produce detectable quantities of fluocinolone acetonide in human milk, or affect breastfed infants or milk production. The developmental and health benefits of breastfeeding should be considered, along with the mother's clinical need for YUTIQ and any potential adverse effects on the breastfed child from YUTIQ.

## 8.4 Pediatric Use

Safety and effectiveness of YUTIQ in pediatric patients have not been established.


## 8.5 Geriatric Use

No overall differences in safety or effectiveness have been observed between elderly and younger patients.

## 11. DESCRIPTION

YUTIQ is a sterile non-bioerodible intravitreal implant containing 0.18 mg fluocinolone acetonide in a 36-month sustained-release drug delivery system. YUTIQ is designed to release fluocinolone acetonide at an initial rate of 0.25 mcg/day. YUTIQ is preloaded into a single-dose applicator to facilitate injection of the implant directly into the vitreous. The drug substance is a synthetic corticosteroid, fluocinolone acetonide.

The chemical name for fluocinolone acetonide is (6 $\alpha$ ,11 $\beta$ , 16 $\alpha$ )-6,9-difluoro-11,21-dihydroxy-16,17-[(1-methylethylidene)bis-(oxy)]-pregna-1,4-diene-3,20-dione. Its chemical structure is:



MW 452.50; molecular formula C<sub>24</sub>H<sub>30</sub>F<sub>2</sub>O<sub>6</sub>

Fluocinolone acetonide is a white or almost white, microcrystalline powder, practically insoluble in water, soluble in methanol, ethanol, chloroform and acetone, and sparingly soluble in ether.

Each YUTIQ consists of a light brown 3.5mm x 0.37mm implant containing 0.18 mg of the active ingredient fluocinolone acetonide and the following inactive ingredients: polyimide tube, polyvinyl alcohol, silicone adhesive and water for injection.



## 12. CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Corticosteroids inhibit inflammatory responses to a variety of inciting agents including multiple inflammatory cytokines. They inhibit edema, fibrin deposition, capillary dilation, leukocyte migration, capillary proliferation, fibroblast proliferation, deposition of collagen, and scar formation associated with inflammation.

Corticosteroids are thought to act by inhibition of phospholipase A<sub>2</sub> via induction of inhibitory proteins collectively called lipocortins. It is postulated that these proteins control biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting release of the common precursor, arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A<sub>2</sub>.

## 13. NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term animal studies have not been conducted to determine the carcinogenic potential or the effect on fertility of YUTIQ.

Fluocinolone acetonide was not genotoxic *in vitro* in the Ames test (*S. typhimurium* and *E. coli*) and the mouse lymphoma TK assay, or *in vivo* in the mouse bone marrow micronucleus assay.



## 14. CLINICAL STUDIES

The efficacy of YUTIQ was assessed in two randomized (2:1, YUTIQ: sham-injection), multi-centre, double-masked, parallel-groups studies (NCT #01694186 and #02746991) that enrolled patients with non-infectious uveitis affecting the posterior segment of the eye. The primary efficacy endpoint in both trials was the proportion of patients who experienced a recurrence of uveitis in the study eye within 6 months of follow-up; recurrence was also assessed at 12 months. Recurrence of uveitis was defined as either deterioration in visual acuity, vitreous haze attributable to non-infectious uveitis or the need for rescue medications.

**Table 3: Efficacy Results of Recurrence of Uveitis in Randomized Study Eyes**

	Study 1		Study 2	
	YUTIQ N = 87	Sham N = 42	YUTIQ N = 101	Sham N = 52
Eyes with recurrence within 6 months, n (%)	16 (18%)	33 (79%)	22 (22%)	28 (54%)
Difference (95% CI) in recurrence rates	60% (41%, 73%)		32% (15%, 48%)	
P-value	< 0.01		< 0.01	
Eyes with recurrence within 12 months, n (%)	24 (28%)	36 (86%)	33 (33%)	31 (60%)
Difference (95% CI) in recurrence rates	58% (40%, 70%)		27% (9%, 43%)	

**Figure 2: Time to First Recurrence of Uveitis (ITT: All Randomized Patients)**



## **16. HOW SUPPLIED/STORAGE AND HANDLING**

YUTIQ™ (fluocinolone acetonide intravitreal implant) 0.18 mg is supplied in a sterile single-dose preloaded applicator with a 25-gauge needle, packaged in a sealed sterile foil pouch inside a sealed Tyvek pouch inside a carton box.

NDC 71879-136-01

**Storage:** Store at 15° C to 30° C (59° F to 86° F).

## **17. PATIENT COUNSELING INFORMATION**

### **Steroid-related Effects**

Advise patients that a cataract may occur after treatment with YUTIQ. If this occurs, advise patients that their vision will decrease, and they will need an operation to remove the cataract and restore their vision.

Advise patients that they may develop increased intraocular pressure with YUTIQ treatment, and the increased IOP may need to be managed with eye drops or surgery.

### **Intravitreal Injection-related Effects**

Advise patients that in the days following intravitreal injection of YUTIQ, they are at risk for potential complications including, but not limited to, the development of endophthalmitis or changes in intraocular pressure.

### **When to Seek Physician Advice**

Advise patients that if the eye becomes red, sensitive to light, painful, or develops a change in vision, they should seek immediate care from an ophthalmologist.

### **Driving and Using Machines**

Inform patients that they may experience temporary visual blurring after receiving an intravitreal injection. Advise patients not to drive or use machines until this has been resolved.

Manufactured by:

EyePoint Pharmaceuticals US, Inc.

480 Pleasant Street

Watertown, MA 02472 USA

Patented.

## Front Panel Box

**YUTIQ<sup>TM</sup>**  
(flucinolone acetone)  
intravitreal implant 0.18 mg

**Contains:** one sterile intravitreal implant in a SINGLE USE ONLY applicator with 25 gauge needle.

**Each implant contains:**

**Active:** 0.18 mg flucinolone acetone  
**Inactive:** polyimide tube, polyvinyl alcohol, silicone adhesive and water for injection. Do not use if inner pouch is damaged.

Dispose of the applicator safely in biohazard sharps container.

Rx Only. Store at 15° C to 30° C (59° F to 86° F).  
Keep out of reach of children.

## Back Panel Box

Part Number: XX-XXX-XX  
Revision Date: YYY-MM-DD  
Lot Number: XX-XXXX  
Expiration Date: YYY-MM-DD

EvePoint Pharmaceuticals US, Inc.  
480 Pleasant Street, Suite B300  
Watertown, MA 02472  
[www.evepointpharma.com](http://www.evepointpharma.com)

Reserved for  
serialization

Box Top Spine

Contains one sterile intravitreal implant in a SINGLE USE ONLY applicator with 25 gauge needle. Each implant contains: Active: 0.18 mg fluocinolone acetonide  
Inactive: polyimide tube, polyvinyl alcohol, silicone adhesive and water for injection. Do not use if inner pouch is damaged.

**YUTIQ**<sup>TM</sup>  
fluocinolone acetonide intravitreal implant 0.18 mg

Box Bottom Spine

**YUTIQ**<sup>TM</sup>  
fluocinolone acetonide intravitreal implant 0.18 mg





Box Left Side

**YUTIQ**<sup>TM</sup>  
(fluocinolone acetonide intravitreal implant) 0.18 mg

Box Right Side

**YUTIQ**<sup>TM</sup>  
(fluocinolone acetonide intravitreal implant) 0.18 mg


### Outer Tyvek Pouch Label

<p><b>NDC 71879-136-01</b></p>  <p><b>Store at 15° C to 30° C (59° F to 86° F).</b> Keep out of reach of children.</p> <p>Eyepoint Pharmaceuticals US, Inc.</p> <p><b>Contains:</b> <b>One sterile-SINGLE USE ONLY Applicator</b></p> <p>For Intravitreal Injection Use Only. Read the package insert before use. Do not use if inner pouch is damaged. Use applicator immediately after opening. Dispose of the applicator safely in biohazard sharps container.</p>	<p>Part Number: XX-XXX-XX Revision Date: YYYY-MM-DD Lot Number: XX-XXXX Expiration Date: YYYY-MM-DD</p> 
--	---

### Inner Foil Pouch Label

**YUTIQ™**  
(fluocinolone acetonide  
intravitreal implant)  
0.18 mg

### Applicator w/Embossed Name



-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

WILEY A CHAMBERS  
10/12/2018