

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

210655Orig1s000

LABELING

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use **PERSERIS™** safely and effectively. See full prescribing information for **PERSERIS**.

PERSERIS (risperidone) for extended-release injectable suspension, for subcutaneous use.

Initial U.S. Approval: 1993

WARNING: INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

See full prescribing information for complete boxed warning.

- Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death.
- **PERSERIS™ is not approved for use in patients with dementia-related psychosis. (5.1)**

-----INDICATIONS AND USAGE-----

PERSERIS is an atypical antipsychotic indicated for the treatment of schizophrenia in adults. (1)

-----DOSAGE AND ADMINISTRATION-----

- Establish tolerability with oral risperidone. (2.1)
- **PERSERIS** may be initiated at a dose of 90 mg or 120 mg. (2.1)
- Supplementation with oral risperidone is not recommended. (2.1)
- Prior to use, the product is constituted by coupling the liquid and powder syringes and passing the contents back and forth between the syringes. (2.4)
- Failure to fully mix the medication could result in incorrect dosage. (2.4)
- Administer monthly by subcutaneous injection in the abdomen by a healthcare professional. Do not administer by any other route. (2.4)
- Do not administer more than one dose (90 mg or 120 mg total) per month. (2.1)

-----DOSAGE FORMS AND STRENGTHS-----

For extended-release injectable suspension: 90 mg and 120 mg risperidone. (3)

-----CONTRAINDICATIONS-----

Known hypersensitivity to risperidone, paliperidone, or other components of **PERSERIS**. (4)

-----WARNINGS AND PRECAUTIONS-----

- Cerebrovascular Adverse Reactions, Including Stroke in Elderly Patients with Dementia-Related Psychosis: Increased risk of cerebrovascular adverse reactions (e.g., stroke, transient ischemic attack), including fatalities. **PERSERIS** is not approved for use in patients with dementia-related psychosis. (5.2)

- Neuroleptic Malignant Syndrome (NMS): Manage with immediate discontinuation and close monitoring. (5.3)
- Tardive Dyskinesia: Discontinue treatment if clinically appropriate. (5.4)
- Metabolic Changes: Monitor for hyperglycemia, dyslipidemia, and weight gain. (5.5)
- Hyperprolactinemia: Prolactin elevations occur and persist during chronic administration. Long-standing hyperprolactinemia, when associated with hypogonadism, may lead to decreased bone density in females and males. (5.6)
- Orthostatic Hypotension: Monitor heart rate and blood pressure and warn patients with known cardiovascular disease or cerebrovascular disease, and risk of dehydration or syncope. (5.7)
- Leukopenia, Neutropenia, and Agranulocytosis: Perform complete blood counts (CBC) in patients with a history of a clinically significant low white blood cell count (WBC) or history of leukopenia or neutropenia. Consider discontinuing **PERSERIS** if a clinically significant decline in WBC occurs in absence of other causative factors. (5.9)
- Potential for Cognitive and Motor Impairment: Use caution when operating machinery. (5.10)
- Seizures: Use caution in patients with a history of seizures or with conditions that lower the seizure threshold. (5.11)

-----ADVERSE REACTIONS-----

The most common adverse reactions in clinical trials ($\geq 5\%$ and greater than twice placebo) were increased weight, sedation/somnolence, and musculoskeletal pain. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Indivior Inc. at 1-877-782-6966 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----DRUG INTERACTIONS-----

- Carbamazepine and other strong CYP3A4 inducers decrease plasma concentrations of risperidone. (2.3, 7.1)
- Fluoxetine, paroxetine, and other strong CYP2D6 inhibitors increase risperidone plasma concentration. (2.3, 7.1)

-----USE IN SPECIFIC POPULATIONS-----

- Pregnancy May cause extrapyramidal and/or withdrawal symptoms in neonates with third trimester exposure. (8.1)
- Renal or Hepatic Impairment: Carefully titrate on oral risperidone up to at least 3 mg before initiating treatment with **PERSERIS** at a dose of 90 mg. (8.6, 8.7)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 07/2018

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

- 1 INDICATIONS AND USAGE**
- 2 DOSAGE AND ADMINISTRATION**
 - 2.1 Recommended Dosage
 - 2.2 Dosage Recommendations for Patients with Renal or Hepatic Impairment
 - 2.3 Dosage Recommendations for Concomitant Use with Strong CYP2D6 Inhibitors and Strong CYP3A4 Inducers
 - 2.4 Instructions for Use
- 3 DOSAGE FORMS AND STRENGTHS**
- 4 CONTRAINDICATIONS**
- 5 WARNINGS AND PRECAUTIONS**
 - 5.1 Increased Mortality in Elderly Patients with Dementia-Related Psychosis
 - 5.2 Cerebrovascular Adverse Events, Including Stroke, in Elderly Patients with Dementia-Related Psychosis
 - 5.3 Neuroleptic Malignant Syndrome (NMS)
 - 5.4 Tardive Dyskinesia
 - 5.5 Metabolic Changes
 - 5.6 Hyperprolactinemia
 - 5.7 Orthostatic Hypotension
 - 5.8 Falls
 - 5.9 Leukopenia, Neutropenia, and Agranulocytosis
 - 5.10 Potential for Cognitive and Motor Impairment
 - 5.11 Seizures
 - 5.12 Dysphagia
 - 5.13 Priapism
 - 5.14 Body Temperature Regulation
- 6 ADVERSE REACTIONS**
 - 6.1 Clinical Trials Experience
 - 6.2 Postmarketing Experience
- 7 DRUG INTERACTIONS**
 - 7.1 Drugs Having Clinically Important Interactions with PERSERIS
 - 7.2 Drugs Having No Clinically Important Interactions with PERSERIS

- 8 USE IN SPECIFIC POPULATIONS**
 - 8.1 Pregnancy
 - 8.2 Lactation
 - 8.3 Females and Males of Reproductive Potential
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use
 - 8.6 Renal Impairment
 - 8.7 Hepatic Impairment
- 10 OVERDOSAGE**
 - 10.1 Human Experience
 - 10.2 Management of Overdosage
- 11 DESCRIPTION**
- 12 CLINICAL PHARMACOLOGY**
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics
 - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY**
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES**
- 16 HOW SUPPLIED/STORAGE AND HANDLING**
- 17 PATIENT COUNSELING INFORMATION**

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS

Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. PERSERIS™ is not approved for the treatment of patients with dementia-related psychosis and has not been studied in this population [see Warnings and Precautions (5.1)].

1 INDICATIONS AND USAGE

PERSERIS™ is indicated for the treatment of schizophrenia in adults [see Clinical Studies (14)].

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

PERSERIS is to be administered as an abdominal subcutaneous injection only. Do not administer by any other route.

Each injection must be administered by a healthcare professional using the prepackaged injection syringe and enclosed safety needle [see Dosage and Administration (2.4)].

For patients who have never taken risperidone, establish tolerability with oral risperidone prior to starting PERSERIS.

Initiate PERSERIS at a dose of 90 mg or 120 mg once monthly by subcutaneous injection. Do not administer more than one dose (90 mg or 120 mg total) per month.

Based on average plasma concentrations (C_{avg}) of risperidone and total active moiety, PERSERIS 90 mg corresponds to 3 mg/day oral risperidone and PERSERIS 120 mg corresponds to 4 mg/day oral risperidone. Patients who are on stable oral risperidone doses lower than 3 mg/day or higher than 4 mg/day may not be candidates for PERSERIS [see Clinical Pharmacology (12.3) and Clinical Studies (14)].

Neither a loading dose nor any supplemental oral risperidone is recommended. A patient who misses a dose should receive the next dose as soon as possible.

2.2 Dosage Recommendations for Patients with Renal or Hepatic Impairment

PERSERIS has not been studied in patients with renal or hepatic impairment and should be used with caution in these special populations. Prior to initiating treatment with PERSERIS in these patients, it is advisable that patients be carefully titrated up to at least 3 mg daily of oral risperidone. If patients can tolerate 3 mg of oral risperidone and are psychiatrically stable, a dose of PERSERIS 90 mg may be considered [see Use in Specific Populations (8.6, 8.7) and Clinical Pharmacology (12.3)].

2.3 Dosage Recommendations for Concomitant Use with Strong CYP2D6 Inhibitors and Strong CYP3A4 Inducers

Co-administration with Strong CYP2D6 Inhibitors

When initiation of fluoxetine or paroxetine is considered, patients may be placed on the lowest dose (90 mg) of PERSERIS between 2 to 4 weeks before the planned start of fluoxetine or paroxetine therapy to adjust for the expected increase in plasma concentrations of risperidone.

When fluoxetine or paroxetine is initiated in patients receiving PERSERIS 90 mg, it is recommended to continue treatment with 90 mg unless clinical judgment necessitates interruption of PERSERIS treatment [see *Drug Interactions (7.1)*].

Co-administration with Strong CYP3A4 Inducers

At the initiation of therapy with carbamazepine or other known hepatic enzyme inducers, patients should be closely monitored during the first 4 to 8 weeks. In patients receiving PERSERIS 90 mg, consider increasing the dose to 120 mg. In patients receiving PERSERIS 120 mg, additional oral risperidone therapy may need to be considered.

On discontinuation of carbamazepine or other strong CYP3A4 hepatic enzyme inducers, the dosage of PERSERIS or any additional oral risperidone therapy should be re-evaluated and, if necessary, decreased to adjust for the expected increase in plasma concentration of risperidone.

For patients treated with PERSERIS 90 mg and discontinuing from carbamazepine or other strong CYP3A4 enzyme inducers, it is recommended to continue treatment with the 90 mg dose unless clinical judgment necessitates interruption of PERSERIS treatment [see *Drug Interactions (7.1)*].

2.4 Instructions for Use

Important Information

- For abdominal subcutaneous injection, only. Do not administer by any other route.
- To be administered by a healthcare professional only.
- Please read the instructions carefully before handling this product.
- Allow package to come to room temperature for at least 15 minutes prior to preparation.
- Only prepare medication when you are ready to administer the dose.
- As a universal precaution, always wear gloves.


1 CHECK CONTENTS

See Figure 1

- One Liquid Syringe (**L**) prefilled with the delivery system. Inspect liquid solution for foreign particles. This is the syringe you will use to inject the patient.
- One Powder Syringe (**P**) prefilled with Risperidone powder. Inspect syringe for consistency of powder color and for foreign particles.
- One sterile 18-gauge, 5/8-inch safety needle.

Parenteral drug products should always be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

Figure 1




2 TAP POWDER SYRINGE

See Figure 2

Hold the Powder Syringe upright and tap the barrel of the syringe to dislodge the packed powder.

NOTE: Powder can become packed during shipping.

Figure 2




3 UNCAP LIQUID AND POWDER SYRINGES

See Figure 3

Remove the cap from the Liquid Syringe, then remove the cap from the Powder Syringe.

Holding both syringes in your non-dominant hand can help with this step.

Figure 3



4 CONNECT THE SYRINGES


See Figure 4

Place the Liquid Syringe on top of the Powder Syringe (to prevent powder spillage) and connect the syringes by twisting approximately $\frac{3}{4}$ turn.

Do not over tighten.

Keep your fingers off the plungers during this step to avoid spillage of the medication.

Figure 4





5 MIX THE PRODUCT

See Figure 5

Failure to fully mix the medication could result in incorrect dosage.

Figure 5



Premixing

- Transfer the contents of the Liquid Syringe into the Powder Syringe.
- Gently push the Powder Syringe plunger until you feel resistance (to wet powder and avoid compacting).
- Repeat this gentle back-and-forth process for **5 cycles**.

Complete mixing

- Continue mixing the syringes for an additional **55 cycles**.
- This mixing can be more vigorous than when premixing.
- **Figure 5** illustrates a correct full cycle.

When fully mixed, the product should be a cloudy suspension that is uniform in color. It can vary from white to yellow-green in color. If you see any clear areas in the mixture, continue to mix until the distribution of the color is uniform. The product is designed to deliver risperidone 90 mg or 120 mg.

6 PREPARE INJECTION SYRINGE


See Figure 6

Failure to aspirate the liquid from the Powder Syringe may result in incorrect dosage.

- First, transfer all contents into the Liquid Syringe.
- Next, perform the following actions **SIMULTANEOUSLY**:
 - maintain slight pressure on the Powder Syringe plunger and
 - pull back gently on the Liquid Syringe plunger while twisting the syringes apart.
- Finally, attach the safety needle by twisting until finger tight.

Check that medication is uniform in color and free from foreign particles.

Figure 6



7 PREPARE THE ABDOMINAL INJECTION SITE

See Figure 7


Choose an injection site on the abdomen with adequate subcutaneous tissue that is free of skin conditions (e.g., nodules, lesions, excessive pigment). It is recommended that the patient is in the supine position.

Do not inject into an area where the skin is irritated, reddened, bruised, infected or scarred in any way.

Clean the injection site well with an alcohol pad.

To help minimize irritation, rotate injection sites following a pattern similar to the illustration (Figure 7).

Figure 7



8 REMOVE EXCESS AIR FROM SYRINGE

See Figure 8

Hold the syringe upright for several seconds to allow air bubbles to rise.

Remove needle cover and slowly depress the plunger to push out the excess air from the syringe.

If medication is seen at the needle tip, pull back slightly on the plunger to prevent medication spillage.

Due to the viscous nature of the medication, bubbles will not rise as quickly as those in an aqueous solution.

Figure 8




9 PINCH INJECTION SITE

See Figure 9

Pinch the skin around the injection area. Be sure to pinch enough skin to accommodate the size of the needle. Lift the adipose tissue from the underlying muscle to prevent accidental intramuscular injection.

Figure 9



10 INJECT THE MEDICATION

See Figure 10


Insert needle fully into the subcutaneous tissue.

Inject the medication slow and steady.

PERSERIS is for subcutaneous administration only. Do not inject by any other route.

NOTE: Actual angle of injection will depend on the amount of subcutaneous tissue.

Figure 10



11 WITHDRAW NEEDLE

See Figure 11

Withdraw the needle at the same angle used for insertion and release pinched skin.

Do not rub the injection area after the injection. If there is bleeding, apply a gauze pad or bandage but use minimal pressure.

Figure 11



12 LOCK THE NEEDLE GUARD AND DISPOSE OF SYRINGE

See Figure 12

Lock the needle guard into place by pushing it against a hard surface such as a table.

Dispose of all syringe components in a secure sharps disposal container.

Figure 12



13 INSTRUCT THE PATIENT

See Figure 13

Advise the patient that they may have a lump for several weeks that will decrease in size over time. It is important that the patient not rub or massage the injection site and to be aware of the placement of any belts or clothing waistbands.

Figure 13



3 DOSAGE FORMS AND STRENGTHS

PERSERIS™ (risperidone) for extended-release injectable suspension for subcutaneous use is available in strengths of 90 mg and 120 mg.

Each strength is provided as a kit which includes: one pre-filled syringe containing a white to yellow risperidone powder in a sealed pouch, one pre-filled syringe containing a colorless to yellow delivery system in a sealed pouch, and one 18-gauge, 5/8-inch needle.

4 CONTRAINDICATIONS

PERSERIS is contraindicated in patients with a known hypersensitivity to risperidone, its metabolite, paliperidone, or to any of its components. Hypersensitivity reactions, including anaphylactic reactions and angioedema, have been reported in patients treated with risperidone or paliperidone.

5 WARNINGS AND PRECAUTIONS

5.1 Increased Mortality in Elderly Patients with Dementia-Related Psychosis

Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. Analyses of 17 placebo-controlled trials (modal duration of 10 weeks), largely in patients taking atypical antipsychotic drugs, revealed a risk of death in drug-treated patients between 1.6 to 1.7 times the risk of death in placebo-treated patients. Over the course of a typical 10-week controlled trial, the rate of death in drug-treated patients was about 4.5%, compared to a rate of about 2.6% in the placebo group. Although the causes of death were varied, most of the deaths appeared to be either cardiovascular (e.g., heart failure, sudden death) or infectious (e.g., pneumonia) in nature. Observational studies suggest that, similar to atypical antipsychotic drugs, treatment with conventional antipsychotic drugs may increase mortality. The extent to which the findings of increased mortality in observational studies may be attributed to the antipsychotic drug as opposed to some characteristic(s) of the patients is not clear. PERSERIS is not approved for the treatment of patients with dementia-related psychosis.

5.2 Cerebrovascular Adverse Reactions, Including Stroke, in Elderly Patients with Dementia-Related Psychosis

Cerebrovascular adverse reactions (e.g., stroke, transient ischemic attack), including fatalities, were reported in patients (mean age 85 years; range 73 to 97) in trials of oral risperidone in elderly patients with dementia-related psychosis. In placebo-controlled trials, there was a significantly higher incidence of cerebrovascular adverse reactions in patients treated with oral risperidone compared to patients treated with placebo. PERSERIS is not approved for the treatment of patients with dementia-related psychosis.

5.3 Neuroleptic Malignant Syndrome (NMS)

A potentially fatal symptom complex sometimes referred to as NMS has been reported in association with antipsychotic drugs. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status, and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmia). Additional signs may include elevated creatine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure.

The diagnostic evaluation of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to identify cases in which the clinical presentation includes both serious medical illness (e.g., pneumonia, systemic infection, etc.) and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever, and primary central nervous system pathology.

The management of NMS should include: (1) immediate discontinuation of antipsychotic drugs and other drugs not essential to concurrent therapy; (2) intensive symptomatic treatment and medical monitoring; and (3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment regimens for uncomplicated NMS.

If a patient requires antipsychotic drug treatment after recovery from NMS, reintroduction of drug therapy should be carefully considered. The patient should be carefully monitored, since recurrences of NMS have been reported.

5.4 Tardive Dyskinesia

A syndrome of potentially irreversible, involuntary, dyskinetic movements may develop in patients treated with antipsychotic drugs. Although the prevalence of the syndrome appears to be highest among the elderly, especially elderly women, it is impossible to rely upon prevalence estimates to predict, at the inception of antipsychotic treatment, which patients are likely to develop the syndrome. Whether antipsychotic drug products differ in their potential to cause tardive dyskinesia is unknown.

The risk of developing tardive dyskinesia and the likelihood that it will become irreversible are believed to increase as the duration of treatment and the total cumulative dose of antipsychotic drugs administered to the patient increase. However, the syndrome can develop, although much less commonly, after relatively brief treatment periods at low doses.

The syndrome may remit, partially or completely, if antipsychotic treatment is withdrawn. Antipsychotic treatment, itself, however, may suppress (or partially suppress) the signs and symptoms of the syndrome and thereby may possibly mask the underlying process. The effect that symptomatic suppression has upon the long-term course of the syndrome is unknown.

Given these considerations, PERSERIS should be prescribed in a manner that is most likely to minimize the occurrence of tardive dyskinesia. Chronic antipsychotic treatment should generally be reserved for patients who suffer from a chronic illness that: (1) is known to respond to antipsychotic drugs, and (2) for whom alternative, equally effective, but potentially less harmful treatments are not available or appropriate. In patients who do require chronic treatment, the smallest dose and the shortest duration of treatment producing a satisfactory clinical response should be sought. The need for continued treatment should be reassessed periodically.

If signs and symptoms of tardive dyskinesia appear in a patient treated with PERSERIS, drug discontinuation should be considered. However, some patients may require treatment with PERSERIS despite the presence of the syndrome.

5.5 Metabolic Changes

Atypical antipsychotic drugs have been associated with metabolic changes that may increase cardiovascular/cerebrovascular risk. These metabolic changes include hyperglycemia, dyslipidemia, and body weight gain. While all of the drugs in the class have been shown to produce some metabolic changes, each drug has its own specific risk profile.

Hyperglycemia and Diabetes Mellitus

Hyperglycemia and diabetes mellitus, in some cases extreme and associated with ketoacidosis or hyperosmolar coma or death, have been reported in patients treated with atypical antipsychotics including risperidone. Assessment of the relationship between atypical antipsychotic use and glucose abnormalities is complicated by the possibility of an increased background risk of diabetes mellitus in patients with schizophrenia and the increasing incidence of diabetes mellitus in the general population. Given these confounders, the relationship between atypical antipsychotic use and hyperglycemia-related adverse events is not completely understood. However, epidemiological studies suggest an increased risk of treatment-emergent hyperglycemia-related adverse events in patients treated with the atypical antipsychotics. Precise risk estimates for hyperglycemia-related adverse events in patients treated with atypical antipsychotics are not available.

Patients with an established diagnosis of diabetes mellitus who are started on atypical antipsychotics, including PERSERIS, should be monitored regularly for worsening of glucose control. Patients with risk factors for diabetes mellitus (e.g., obesity, family history of diabetes) who are starting treatment with atypical antipsychotics, including PERSERIS, should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Any patient treated with atypical antipsychotics, including PERSERIS, should be monitored for symptoms of hyperglycemia including polydipsia, polyuria, polyphagia, and weakness. Patients who develop symptoms of hyperglycemia during treatment with atypical antipsychotics, including PERSERIS, should undergo fasting blood glucose testing. In some cases, hyperglycemia has resolved when the atypical antipsychotic, including risperidone, was discontinued; however, some patients required continuation of anti-diabetic treatment despite discontinuation of risperidone.

Data from an 8-week double-blind, placebo-controlled study with PERSERIS in adult subjects with schizophrenia are presented in Table 1.

Table 1. Changes in Fasting Glucose from Baseline to End of Study (EOS) and Postbaseline Abnormal Values of Glucose > 126 mg/dL in an 8-Week Double-Blind, Placebo-Controlled Study in Adult Subjects with Schizophrenia

	PERSERIS 90 mg	PERSERIS 120 mg	Placebo
	n = 98	n = 106	n = 96
Serum Glucose, mg/dL, mean [†]			
Mean Change from Baseline to EOS	5.7	6.3	-0.9
Glucose, > 126 mg/dL			
Proportion of Subjects with Postbaseline Abnormal Values [‡]	12/104 (11.5%)	14/111 (12.6%)	8/109 (7.3%)

[†] The “n”s in the Serum Glucose mean row are the number of subjects with data at baseline and EOS visits.

[‡] Data shown as number of subjects with at least one postbaseline value as denominator and number of subjects satisfying the predefined criterion as numerator.

Similar changes from baseline in serum glucose were observed in subjects receiving PERSERIS during an open-label, 12-month long-term safety study. Additionally, the mean HbA_{1c} increased from 5.6 to 5.7% over the 12 months.

Dyslipidemia

Undesirable alterations in lipids have been observed in patients treated with atypical antipsychotics.

Data from an 8-week double-blind, placebo-controlled study with PERSERIS in adult subjects with schizophrenia are presented in Table 2.

Table 2. Changes in Cholesterol from Baseline to End of Study (EOS) and Postbaseline Abnormal Values of Cholesterol ≥ 300 mg/dL in an 8-Week Double-Blind, Placebo-Controlled Study in Adult Subjects with Schizophrenia

	PERSERIS 90 mg	PERSERIS 120 mg	Placebo
	n = 98	n = 106	n = 96
Cholesterol, mg/dL, mean [†]			
Mean Change from Baseline to EOS	-0.5	-0.5	1.1
Cholesterol, ≥ 300 mg/dL			
Proportion of Subjects with Postbaseline Abnormal Values [‡]	2/104 (1.9%)	2/111 (1.8%)	2/109 (1.8%)

† The “n”s in the Cholesterol mean row are the number of subjects with data at baseline and EOS visits.

‡ Data shown as number of subjects with at least one postbaseline value as denominator and number of subjects satisfying the predefined criterion as numerator.

Weight Gain

Weight gain has been observed with atypical antipsychotic use. Clinical monitoring of weight is recommended.

Data from an 8-week double-blind, placebo-controlled study with PERSERIS in adult subjects with schizophrenia are presented in Table 3.

Table 3. Changes in Body Weight from Baseline to End of Study (EOS) and $\geq 7\%$ Increase from Baseline in an 8-Week Double-Blind, Placebo-Controlled Study in Adult Subjects with Schizophrenia

	PERSERIS 90 mg	PERSERIS 120 mg	Placebo
Weight †	n = 105	n = 112	n = 107
Mean Change from Baseline to EOS, kg	4.4	5.3	2.6
Weight Gain			
$\geq 7\%$ Increase from Baseline‡	35/107 (32.7%)	48/114 (42.1%)	20/111 (18.0%)

† The “n”s in the Weight Change mean row are the number of subjects with data at baseline and end of study visits.

‡ Data shown as number of subjects with at least one postbaseline value as denominator and number of subjects satisfying the predefined criterion as numerator.

In an open-label, 12-month long-term safety study, for all subjects receiving PERSERIS, mean weight increased approximately 2 kg from baseline to Day 85, then remained stable for the remainder of the study.

5.6 Hyperprolactinemia

As with other drugs that antagonize dopamine D₂ receptors, risperidone elevates prolactin levels and the elevation persists during chronic administration. Risperidone is associated with higher levels of prolactin elevation than other antipsychotic agents.

Hyperprolactinemia may suppress hypothalamic GnRH, resulting in reduced pituitary gonadotropin secretion. This, in turn, may inhibit reproductive function by impairing gonadal steroidogenesis in both female and male patients. Galactorrhea, amenorrhea, gynecomastia, and impotence have been reported in patients receiving prolactin-elevating compounds. Long-standing hyperprolactinemia when associated with hypogonadism may lead to decreased bone density in both female and male subjects.

Tissue culture experiments indicate that approximately one-third of human breast cancers are prolactin dependent *in vitro*, a factor of potential importance if the prescription of these drugs is contemplated in a patient with previously detected breast cancer. An increase in pituitary gland, mammary gland, and pancreatic islet cell neoplasia (mammary adenocarcinomas, pituitary and pancreatic adenomas) was observed in the risperidone carcinogenicity studies conducted in mice and rats [see *Nonclinical Toxicology* (13.1)]. Neither clinical studies nor epidemiologic studies conducted to date have shown an

association between chronic administration of this class of drugs and tumorigenesis in humans; the available evidence is considered too limited to be conclusive at this time.

5.7 Orthostatic Hypotension

Risperidone may induce orthostatic hypotension associated with dizziness, tachycardia, and in some patients, syncope, probably reflecting its alpha-adrenergic antagonistic properties.

PERSERIS should be used with particular caution in (1) patients with known cardiovascular disease (history of myocardial infarction or ischemia, heart failure, or conduction abnormalities), cerebrovascular disease, and conditions which would predispose patients to hypotension, e.g., dehydration and hypovolemia, and (2) in the elderly and patients with renal or hepatic impairment. Monitoring of orthostatic vital signs should be considered in all such patients, and a dose reduction should be considered if hypotension occurs. Clinically significant hypotension has been observed with concomitant use of oral risperidone and antihypertensive medication.

5.8 Falls

Somnolence, postural hypotension, motor instability, and sensory instability have been reported with the use of antipsychotics, including PERSERIS, which may lead to falls and, consequently, fractures or other fall-related injuries. For patients, particularly the elderly, with diseases, conditions, or medications that could exacerbate these effects, assess the risk of falls when initiating antipsychotic treatment and recurrently for patients on long-term antipsychotic therapy.

5.9 Leukopenia, Neutropenia, and Agranulocytosis

In clinical trial and/or postmarketing experience, events of leukopenia/neutropenia have been reported temporally related to antipsychotic agents, including risperidone. Agranulocytosis has also been reported.

Possible risk factors for leukopenia/neutropenia include pre-existing low white blood cell count (WBC) and a history of drug-induced leukopenia/neutropenia. Patients with a history of a clinically significant low WBC or a drug-induced leukopenia/neutropenia should have their complete blood count (CBC) monitored frequently during the first few months of therapy and discontinuation of PERSERIS should be considered at the first sign of a clinically significant decline in WBC in the absence of other causative factors.

Patients with clinically significant neutropenia should be carefully monitored for fever or other symptoms or signs of infection and treated promptly if such symptoms or signs occur. Patients with severe neutropenia (absolute neutrophil count $< 1000/\text{mm}^3$) should discontinue PERSERIS and have their WBC followed until recovery.

5.10 Potential for Cognitive and Motor Impairment

In an 8-week, double-blind, placebo-controlled study, somnolence/sedation was reported by 7.0% and 7.7% of subjects treated with PERSERIS 90 mg and 120 mg, respectively.

Since risperidone has the potential to impair judgment, thinking, or motor skills, patients should be cautioned about operating hazardous machinery, including automobiles, until they are reasonably certain that treatment with PERSERIS does not affect them adversely.

5.11 Seizures

Seizures have been observed during pre-marketing studies of risperidone in adult patients with schizophrenia. PERSERIS should be used cautiously in patients with a history of seizures or other conditions that potentially lower the seizure threshold.

5.12 Dysphagia

Esophageal dysmotility and aspiration have been associated with antipsychotic drug use. Aspiration pneumonia is a common cause of morbidity and mortality in patients with advanced Alzheimer's dementia. PERSERIS and other antipsychotic drugs should be used cautiously in patients at risk for aspiration pneumonia [see Boxed Warning and Warnings and Precautions (5.1)].

5.13 Priapism

Priapism has been reported during postmarketing surveillance for other risperidone products. Severe priapism may require surgical intervention.

5.14 Body Temperature Regulation

Disruption of body temperature regulation has been attributed to antipsychotic agents. Both hyperthermia and hypothermia have been reported in association with oral risperidone use. Caution is advised when prescribing PERSERIS for patients who will be exposed to temperature extremes.

6 ADVERSE REACTIONS

The following are discussed in more detail in previous sections of the labeling:

- Increased Mortality in Elderly Patients with Dementia-Related Psychosis [see Boxed Warning and Warnings and Precautions (5.1)]
- Cerebrovascular Adverse Events, Including Stroke, in Elderly Patients with Dementia-Related Psychosis [see Warnings and Precautions (5.2)]
- Neuroleptic Malignant Syndrome (NMS) [see Warnings and Precautions (5.3)]
- Tardive Dyskinesia [see Warnings and Precautions (5.4)]
- Metabolic Changes [see Warnings and Precautions (5.5)]
- Hyperprolactinemia [see Warnings and Precautions (5.6)]
- Orthostatic Hypotension [see Warnings and Precautions (5.7)]
- Falls [see Warnings and Precautions (5.8)]
- Leukopenia, Neutropenia and Agranulocytosis [see Warnings and Precautions (5.9)]
- Potential for Cognitive and Motor Impairment [see Warnings and Precautions (5.10)]
- Seizures [see Warnings and Precautions (5.11)]
- Dysphagia [see Warnings and Precautions (5.12)]
- Priapism [see Warnings and Precautions (5.13)]
- Body Temperature Regulation [see Warnings and Precautions (5.14)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of PERSERIS was evaluated in a total of 814 adult subjects with schizophrenia who received at least 1 dose of PERSERIS during the clinical development program. A total of 322 subjects were exposed to PERSERIS for at least 6 months, of which 234 subjects were exposed to PERSERIS for at least 12 months; 281 and 176 of these, respectively, received the 120 mg dose.

Adverse drug reactions in adult subjects with schizophrenia ($\geq 5\%$ in any PERSERIS-treated group and greater than placebo) during the 8-week double-blind, placebo-controlled study) were weight increased, constipation, sedation/somnolence, pain in extremity, back pain, akathisia, anxiety, and musculoskeletal

pain. In addition, the frequency of reported injection site reactions was similar across treatment groups with both PERSERIS and placebo; the most common ($\geq 5\%$) of which were injection site pain, and erythema. The systemic safety profile for PERSERIS, was consistent with the known safety profile of oral risperidone.

Commonly-Observed Adverse Drug Reactions in Double-Blind, Placebo-Controlled Clinical Studies – Schizophrenia

Adverse Reactions with an incidence of 2% or more and greater than placebo are shown in Table 4.

Table 4. Adverse Drug Reactions in 2% or More of PERSERIS-Treated Subjects (and Greater than Placebo) in an 8-Week Double-Blind, Placebo-Controlled Study

System Organ Class Preferred Term	PERSERIS 90 mg (n = 115)	PERSERIS 120 mg (n = 117)	Placebo (n = 118)
Percentage of Subjects Reporting ADR			
Gastrointestinal disorders			
Constipation	7.0	7.7	5.1
Abdominal discomfort	2.6	2.6	1.7
Dry mouth	1.7	2.6	1.7
Investigations			
Weight increased	13.0	12.8	3.4
Metabolism and nutrition disorders			
Increased appetite	1.7	3.4	1.7
Musculoskeletal and connective tissue disorders			
Back pain	3.5	6.8	4.2
Pain in extremity	0.9	7.7	5.1
Musculoskeletal pain	5.2	5.1	2.5
Musculoskeletal stiffness	2.6	0.9	1.7
Muscle spasms	0	2.6	0
Nervous system disorders			
Sedation*	7.0	7.7	0
Akathisia	2.6	6.8	4.2
Extrapyramidal disorder	4.3	1.7	0.8

System Organ Class Preferred Term	PERSERIS 90 mg (n = 115)	PERSERIS 120 mg (n = 117)	Placebo (n = 118)
Percentage of Subjects Reporting ADR			
Psychiatric disorders			
Anxiety	2.6	6.8	5.1

* Sedation includes sedation and somnolence

Other Adverse Drug Reactions Observed During the Clinical Trial Evaluation of PERSERIS

The following list does not include reactions: 1) already listed in previous tables or elsewhere in labeling, 2) which are part of the disease state, 3) for which a drug cause was remote, 4) which were so general as to be uninformative, or 5) which were not considered to have significant clinical implications.

Blood and Lymphatic System Disorders: neutropenia

Ear and Labyrinth Disorders: vertigo

Endocrine Disorders: hyperprolactinemia

Eye Disorders: blepharospasm

Gastrointestinal Disorders: nausea, dyspepsia, vomiting, diarrhea, abdominal pain upper, salivary hypersecretion, hypoesthesia oral, tongue movement disturbance

General Disorders and Administration Site Conditions: injection site reaction (including injection site pain, induration, pruritus, bruising, erythema, inflammation, swelling and irritation) fatigue, edema peripheral, asthenia, chest discomfort

Investigations: blood prolactin increased, blood glucose increased, glycosylated hemoglobin increased, electrocardiogram abnormal, electrocardiogram QT prolonged, blood creatine phosphokinase increased

Metabolism and Nutrition Disorders: diabetes mellitus, decreased appetite

Musculoskeletal, Connective Tissue, and Bone Disorders: arthralgia, muscle twitching, joint stiffness, trismus

Nervous System Disorders: headache, dizziness, tremor, drooling, dyskinesia, lethargy, dystonia, hypoesthesia, oromandibular dystonia, tardive dyskinesia, cogwheel rigidity, dysarthria, balance disorder, parkinsonian rest tremor, parkinsonism, slow speech

Psychiatric Disorders: insomnia, libido decreased, bruxism, restlessness, anorgasmia, loss of libido

Reproductive System and Breast Disorders: erectile dysfunction, galactorrhea, breast tenderness, breast pain, amenorrhea, breast engorgement, ejaculation delayed, ejaculation disorder, gynecomastia, hypomenorrhea, breast discharge, breast enlargement, ejaculation failure, menstruation delayed, menstruation irregular, polymenorrhea

Skin and Subcutaneous Tissue Disorders: night sweats

Vascular Disorders: hypertension, hypotension, orthostatic hypotension

Other Adverse Reactions Observed During the Clinical Trial Evaluations of Oral Risperidone

The following is a list of additional ADRs that have been reported during the clinical trial evaluation of oral risperidone, regardless of frequency of occurrence:

Blood and Lymphatic System Disorders: anemia, granulocytopenia

Cardiac Disorders: tachycardia, sinus bradycardia, sinus tachycardia, atrioventricular block first degree, bundle branch block left, bundle branch block right, atrioventricular block

Ear and Labyrinth Disorders: ear pain, tinnitus

Eye Disorders: vision blurred, oculogyration, ocular hyperemia, eye discharge, conjunctivitis, eye rolling, eyelid edema, eye swelling, eyelid margin crusting, dry eye, lacrimation increased, photophobia, glaucoma, visual acuity reduced

Gastrointestinal Disorders: dysphagia, fecaloma, fecal incontinence, gastritis, lip swelling, cheilitis, apyptalism

General Disorders: thirst, gait disturbance, chest pain, influenza-like illness, pitting edema, edema, chills, sluggishness, malaise, face edema, discomfort, generalized edema, drug withdrawal syndrome, peripheral coldness, feeling abnormal

Immune System Disorders: drug hypersensitivity

Infections and Infestations: nasopharyngitis, upper respiratory tract infection, sinusitis, urinary tract infection, pneumonia, influenza, ear infection, viral infection, pharyngitis, tonsillitis, bronchitis, eye infection, localized infection, cystitis, cellulitis, otitis media, onychomycosis, acarodermatitis, bronchopneumonia, respiratory tract infection, tracheobronchitis, otitis media chronic

Investigations: body temperature increased, alanine aminotransferase increased, heart rate increased, eosinophil count increased, white blood cell count decreased, hemoglobin decreased, blood creatine phosphokinase increased, hematocrit decreased, body temperature decreased, blood pressure decreased, transaminases increased,

Metabolism and Nutrition Disorders: polydipsia, anorexia

Musculoskeletal, Connective Tissue, and Bone Disorders: joint swelling, musculoskeletal chest pain, posture abnormal, myalgia, neck pain, muscular weakness, muscle rigidity, muscle contracture, rhabdomyolysis

Nervous System Disorders: dizziness postural, disturbance in attention, unresponsive to stimuli, depressed level of consciousness, movement disorder, hypokinesia, bradykinesia, transient ischemic attack, coordination abnormal, cerebrovascular accident, masked facies, speech disorder, syncope, loss of consciousness, muscle contractions involuntary, Parkinson's disease, tongue paralysis, akinesia, cerebral ischemia, cerebrovascular disorder, neuroleptic malignant syndrome, diabetic coma, head titubation

Psychiatric Disorders: agitation, blunted affect, confusional state, middle insomnia, nervousness, sleep disorder, listlessness

Renal and Urinary Disorders: enuresis, dysuria, pollakiuria, urinary incontinence

Reproductive System and Breast Disorders: vaginal discharge, menstrual disorder, retrograde ejaculation, sexual dysfunction

Respiratory, Thoracic, and Mediastinal Disorders: nasal congestion, dyspnea, epistaxis, wheezing, pneumonia aspiration, sinus congestion, dysphonia, productive cough, pulmonary congestion, respiratory tract congestion, rales, respiratory disorder, hyperventilation, nasal edema

Skin and Subcutaneous Tissue Disorders: rash, dry skin, erythema, skin discoloration, skin lesion, pruritus, skin disorder, rash erythematous, rash papular, acne, hyperkeratosis, seborrheic dermatitis, rash generalized, rash maculopapular

Vascular Disorders: flushing

Discontinuations Due to Adverse Drug Reactions (ADRs)

There was no single adverse reaction leading to discontinuation that occurred at a rate of $\geq 2\%$ in PERSERIS-treated patients and greater than placebo.

Dose Dependency of Adverse Drug Reactions in Clinical Trials

Changes in Body Weight

Data from the double-blind placebo-controlled study indicated there was a dose-dependent increase in mean changes in weight from baseline to postdose assessments in the PERSERIS 90 mg and 120 mg groups compared with the placebo group [see *Warnings and Precautions (5.5)*, *Adverse Reactions (6.1, Table 4)*].

Increased Prolactin

In the 8-week double-blind, placebo-controlled study, there was a typical increase in mean prolactin levels in fasting blood samples from baseline to the EOS assessments in both the PERSERIS 90 mg and 120 mg groups, while mean prolactin for the placebo group remained stable during the study. Changes in mean prolactin were dose-dependent and more pronounced in female subjects than male subjects.

Extrapyramidal Symptoms (EPS)

Several methods were used to measure EPS, including: (1) the Barnes Akathisia Rating Scale (BARS) global clinical rating score which evaluates akathisia, (2) the Abnormal Involuntary Movement Scale (AIMS) scores which evaluates dyskinesia, (3) the Simpson-Angus Scale (SAS) global score which broadly evaluates parkinsonism, and (4) the incidence of spontaneous reports of EPS-related adverse reactions.

In the 8-week double-blind, placebo-controlled study, the mean changes from baseline in BARS, AIMS, and SAS total scores were comparable between PERSERIS- and placebo-treated patients. At all postbaseline assessments, mean changes from baseline were between -0.1 and 0.2 (inclusive) for the BARS, between 0 and 0.2 (inclusive) for the AIMS and between -0.1 and 0.2 (inclusive) for the SAS.

The rates of ADRs associated with EPS were similar across treatment groups, including placebo. There was a higher incidence of akathisia in the PERSERIS 120 mg (6.8%) group compared with the PERSERIS 90 mg (2.6%) and placebo group (4.2%); reports of extrapyramidal disorders were higher in the PERSERIS 90 mg group (4.3%) compared with the PERSERIS 120 mg (1.7%) and placebo group (0.8%). In contrast, there was a higher incidence of dystonia in the placebo group (2.5%) compared with the PERSERIS groups (0 and 0.9%, respectively).

Dystonia

Symptoms of dystonia, prolonged abnormal contractions of muscle groups, may occur in susceptible individuals during the first few days of treatment. Dystonic symptoms include: spasm of the neck

muscles, sometimes progressing to tightness of the throat, swallowing difficulty, difficulty breathing, and/or protrusion of the tongue. Although these symptoms can occur at low doses, they occur more frequently and with greater severity with high potency and at higher doses of first generation antipsychotic drugs. An elevated risk of acute dystonia has been observed in males and younger age groups.

Changes in ECG

In the 8-week double-blind, placebo-controlled study, there were no clinically relevant differences in mean changes from baseline to EOS in ECG parameters, including QT_cF (Fridericia's corrected QT interval), QRS and PR intervals, and heart rate, in subjects in either PERSERIS treatment group (90 mg and 120 mg) compared with placebo. Similarly, in the 12-month, long-term safety study, there were no clinically relevant changes in mean ECG interval values from baseline to postdose assessments.

Pain Assessment and Local Injection Site Reactions

Local injection site pain was assessed using subject-reported VAS scales (0 = no pain to 100 = unbearably painful). In the 8-week, double-blind placebo-controlled study, the mean subject-reported injection site pain VAS scores were similar for all treatment groups following both injections. Pain scores decreased from a mean of 27 (VAS score) 1 minute after the first dose to a range of 3 to 7 (VAS score) 30 to 60 minutes postdose. In the 12-month, long-term safety study, the 1-minute postdose injection site pain VAS scores were highest on Day 1 (mean of 25) and decreased over time with subsequent injections (14 to 16 following last injection).

The local injection site was assessed by appropriately trained personnel. Throughout the clinical development program, the maximum reported intensity at any time point for each injection site assessment (pain, tenderness, inflammation/swelling and erythema) was none or mild for most subjects receiving PERSERIS.

Most subjects ($\geq 79\%$) reported no tenderness and most who had tenderness reported mild severity. Less than 1% of subjects had moderate tenderness at any time point and 1 subject at Injections 1, 2, and 5 had severe tenderness. At each time point, most subjects ($\geq 75\%$) reported no pain on injection. Of subjects who did have pain on injection, almost all of these were mild at each time point; only 1 or 2 subjects at Injections 1, 2, 7, and 12 had moderate pain on injection. At least 92% of subjects reported no erythema on each injection. All reports of erythema were of mild severity except for 2 cases of moderate erythema on Injection 1. Inflammation/swelling had a similar profile, with at least 88% of subjects reporting no inflammation/swelling and only mild symptoms except for 1 case of moderate severity on Injection 1.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post approval use of oral risperidone. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. These adverse reactions include: alopecia, anaphylactic reaction, angioedema, atrial fibrillation, cardiopulmonary arrest, diabetic ketoacidosis in patients with impaired glucose metabolism, dysgeusia, hypoglycemia, hypothermia, ileus, inappropriate antidiuretic hormone secretion, intestinal obstruction, jaundice, mania, pancreatitis, pituitary adenoma, precocious puberty, pulmonary embolism, QT prolongation, sleep apnea syndrome, sudden death, thrombocytopenia, thrombotic thrombocytopenic purpura, urinary retention, and water intoxication.

7 DRUG INTERACTIONS

The interactions of PERSERIS with co-administration of other drugs have not been studied. The drug interaction data provided in this section is based on studies with oral risperidone.

7.1 Drugs Having Clinically Important Interactions with PERSERIS

Table 5 includes clinically significant drug interactions with PERSERIS.

Table 5: Clinically Important Drug Interactions with PERSERIS

Strong CYP2D6 Inhibitors	
<i>Clinical Impact:</i>	Concomitant use of PERSERIS with strong CYP2D6 inhibitors may increase the plasma exposure of risperidone and lower the plasma exposure of a major active metabolite, 9-hydroxyrisperidone [see <i>Clinical Pharmacology</i> (12.3)].
<i>Intervention:</i>	When initiation of strong CYP2D6 inhibitors is considered, patients may be placed on the lowest dose (90 mg) of PERSERIS between 2 to 4 weeks before the planned start of strong CYP2D6 inhibitors to adjust for the expected increase in plasma concentrations of risperidone. When strong CYP2D6 inhibitors is initiated in patients receiving PERSERIS 90 mg, it is recommended to continue treatment with 90 mg unless clinical judgment necessitates interruption of PERSERIS treatment. The effects of discontinuation of strong CYP2D6 inhibitors on the pharmacokinetics of risperidone and 9-hydroxyrisperidone have not been studied [see <i>Clinical Pharmacology</i> (12.3)].
<i>Examples:</i>	paroxetine, fluoxetine, quinidine
Strong CYP3A4 Inducers	
<i>Clinical Impact:</i>	Concomitant use of PERSERIS and a strong CYP3A4 inducer may cause decreases in the combined plasma concentrations of risperidone and 9-hydroxyrisperidone which could lead to decreased efficacy of PERSERIS [see <i>Clinical Pharmacology</i> (12.3)].
<i>Intervention:</i>	Changes in efficacy and safety should be carefully monitored with any dose adjustment of PERSERIS. At the initiation of therapy with a strong CYP3A4 inducer, patients should be closely monitored during the first 4 to 8 weeks. In patients receiving PERSERIS 90 mg, consider increasing the dose to 120 mg. In patients receiving PERSERIS 120 mg, additional oral risperidone therapy may need to be considered. On discontinuation of a strong CYP3A4 inducer, the dosage of PERSERIS or any additional oral risperidone therapy should be re-evaluated and, if necessary, decreased to adjust for the expected increase in plasma concentration of risperidone and 9-hydroxyrisperidone. For patients treated with PERSERIS 90 mg and discontinuing from a strong CYP3A4 inducer, it is recommended to continue treatment with the 90 mg dose unless clinical judgment necessitates interruption of PERSERIS treatment [see <i>Dosage and Administration</i> (2.3)].
<i>Examples:</i>	rifampin, carbamazepine, phenytoin, phenobarbital
Centrally-Acting Drugs and Alcohol	
<i>Clinical Impact:</i>	Due to additive pharmacologic effects, the concomitant use of centrally-acting drugs, including alcohol, may increase nervous system disorders.
<i>Intervention:</i>	Caution should be used when PERSERIS is administered in combination with

	other centrally-acting drugs or alcohol.
<i>Examples:</i>	Antipsychotics, alcohol
Hypotensive Agents	
<i>Clinical Impact:</i>	Because of its potential for inducing hypotension, PERSERIS may enhance the hypotensive effects of other therapeutic agents with this potential.
<i>Intervention:</i>	Caution should be used when PERSERIS is administered in combination with other therapeutic agents with hypotensive effects.
<i>Examples:</i>	Antihypertensive Drugs
Dopamine Agonists	
<i>Clinical Impact:</i>	Agents with central antidopaminergic activity such as PERSERIS may antagonize the pharmacologic effects of dopamine agonists.
<i>Intervention:</i>	Caution should be used when PERSERIS is administered in combination with levodopa and dopamine agonists.
<i>Examples:</i>	carbidopa, levodopa

7.2 Drugs Having No Clinically Important Interactions with PERSERIS

Based on pharmacokinetic studies with oral risperidone, no dosage adjustment of PERSERIS is required when administered concomitantly with amitriptyline, cimetidine, ranitidine, clozapine, topiramate and moderate CYP3A4 inhibitors (erythromycin). Additionally, no dosage adjustment is necessary for lithium, valproate, topiramate, digoxin and CYP2D6 substrates (donepezil and galantamine) when co-administered with PERSERIS [see *Clinical Pharmacology* (12.3)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to atypical antipsychotics, including PERSERIS, during pregnancy. Healthcare professionals are encouraged to register patients by contacting the National Pregnancy Registry for Atypical Antipsychotics at 1-866-961-2388 or online at <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>.

Risk Summary

Neonates exposed to antipsychotic drugs during the third trimester of pregnancy are at risk for extrapyramidal and/or withdrawal symptoms following delivery (*see Clinical Considerations*). Overall available data from published epidemiologic studies of pregnant women exposed to risperidone have not established a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes (*see Data*). There are risks to the mother associated with untreated schizophrenia and with exposure to antipsychotics, including PERSERIS, during pregnancy (*see Clinical Considerations*).

Oral administration of risperidone to pregnant mice caused cleft palate at doses 3 to 4 times the maximum recommended human dose (MRHD) of 16 mg/day with maternal toxicity observed at 4 times the MRHD based on mg/m² body surface area. Risperidone was not teratogenic in rats or rabbits at doses up to 6 times the MRHD based on mg/m² body surface area. Increased stillbirths and decreased birth weight

occurred after oral risperidone administration to pregnant rats at 1.5 times the MRHD based on mg/m² body surface area. Learning was impaired in offspring of rats when the dams were dosed at 0.6 times the MRHD and offspring mortality increased at doses 0.1 to 3 times the MRHD based on mg/m² body surface area.

Subcutaneous administration of the delivery system to pregnant rats and rabbits during the period of organogenesis caused developmental toxicity that included post-implantation loss, decreased number of live fetuses, decreased fetal weight and fetal malformations (external, skeletal, and visceral), at doses that are 52 (rat) and 43 (rabbit) times the delivery system amount present in 120 mg risperidone subcutaneous injectable suspension based on mg/m² body surface area. These effects could be attributed to NMP an excipient in the delivery system based on information in the published literature (*see Data*). Subcutaneous administration of the delivery system to pregnant and lactating rats had no effect on embryo/fetal and postnatal development at doses up to 17 times the delivery system amount present in 120 mg risperidone subcutaneous injectable suspension based on mg/m² body surface area.

The estimated background risks of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Clinical Considerations

Disease-associated maternal and/or embryo/fetal risk

There is a risk to the mother from untreated schizophrenia, including increased risk of relapse, hospitalization, and suicide. Schizophrenia is associated with increased adverse perinatal outcomes, including preterm birth. It is not known if this is a direct result of the illness or other comorbid factors.

Fetal/neonatal adverse reactions

Extrapyramidal and/or withdrawal symptoms, including agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress, and feeding disorder have been reported in neonates who were exposed to antipsychotic drugs, including risperidone, during the third trimester of pregnancy. These symptoms have varied in severity. Monitor neonates for extrapyramidal and/or withdrawal symptoms and manage symptoms appropriately. Some neonates recovered within hours or days without specific treatment; others required prolonged hospitalization.

Data

Human Data

Published data from observational studies, birth registries, and case reports on the use of atypical antipsychotics during pregnancy do not report a clear association with antipsychotics and major birth defects. A prospective observational study including 6 women treated with risperidone demonstrated placental passage of risperidone. A retrospective cohort study from a Medicaid database of 9258 women exposed to antipsychotics during pregnancy did not indicate an overall increased risk for major birth defects. There was a small increase in the risk of major birth defects (RR = 1.26, 95% CI 1.02 to 1.56) and of cardiac malformations (RR = 1.26, 95% CI 0.88 to 1.81) in a subgroup of 1566 women exposed to risperidone during the first trimester of pregnancy; however, there is no mechanism of action to explain the difference in malformation rates.

Animal data

No developmental toxicity studies were conducted with subcutaneous risperidone suspension.

Oral administration of risperidone to pregnant mice during organogenesis caused cleft palate at 10 mg/kg/day which is 3 times the MRHD of 16 mg/day based on mg/m² body surface area; maternal toxicity occurred at 4 times the MRHD. Risperidone was not teratogenic when administered orally to rats at 0.6 to 10 mg/kg/day and rabbits at 0.3 to 5 mg/kg/day, which are up to 6 times the MRHD of 16 mg/day risperidone based on mg/m² body surface area. Learning was impaired in offspring of rats dosed orally throughout pregnancy at 1 mg/kg/day which is 0.6 times the MRHD and neuronal cell death increased in fetal brains of offspring of rats dosed during pregnancy at 1 and 2 mg/kg/day which are 0.6 and 1.2 times the MRHD based on mg/m² body surface area; postnatal development and growth of the offspring were also delayed.

Rat offspring mortality increased during the first 4 days of lactation when pregnant rats were dosed throughout gestation at 0.16 to 5 mg/kg/day which are 0.1 to 3 times the MRHD of 16 mg/day based on mg/m² body surface area. It is not known whether these deaths were due to a direct effect on the fetuses or pups or to effects on the dams; a no-effect dose could not be determined. The rate of stillbirths was increased at 2.5 mg/kg or 1.5 times the MRHD based on mg/m² body surface area. In a rat cross-fostering study the number of live offspring was decreased, the number of stillbirths increased, and the birth weight was decreased in offspring of drug-treated pregnant rats. In addition, the number of deaths increased by Day 1 among offspring of drug-treated pregnant rats, regardless of whether or not the offspring were cross-fostered. Risperidone also appeared to impair maternal behavior in that offspring body weight gain and survival (from Day 1 to 4 of lactation) were reduced in offspring born to control but reared by drug-treated dams. All of these effects occurred at 5 mg/kg which is 3 times the MRHD based on mg/m² and the only dose tested in the study.

Subcutaneous administration of the delivery system to pregnant rats and rabbits during the period of organogenesis caused maternal toxicity (decreased body weight, weight gain and food intake), post-implantation loss, decrease in number of live fetuses and decrease in fetal weight at doses that are 52 (rat), and 43 (rabbit) times the delivery system amount present in monthly 120 mg risperidone subcutaneous injectable suspension based on mg/m² body surface area. Developmental toxicity in both rat and rabbit included skeletal and visceral malformations at doses 35 (rat), and 43 (rabbit) times the delivery system amount present in monthly 120 mg risperidone subcutaneous injectable suspension based on mg/m² body surface area. The NOAEL dose for these effects in both species is 17 times the delivery system amount present in monthly 120 mg risperidone subcutaneous injectable suspension based on mg/m² body surface area. These effects could be related to NMP, an excipient present in the delivery system. In published animal developmental toxicity studies, NMP administered orally daily to pregnant rats during organogenesis produced developmental toxicity below maternally toxic levels and resulted in dose-dependent decrease in fetal body weights, increased incidence of post-implantation loss, incomplete ossification and increased incidence of external, visceral and skeletal malformations. These toxicities occurred at doses that are ~3 to 12 times the NMP amount present in monthly 120 mg risperidone subcutaneous injectable suspension based on mg/m² body surface area.

8.2 Lactation

Risk Summary

Limited data from published literature reports the presence of risperidone and its metabolite, 9-hydroxyrisperidone, in human breast milk at relative infant dose ranging between 2.3 and 4.7% of the maternal weight-adjusted dosage. There are reports of sedation, failure to thrive, jitteriness, and extrapyramidal symptoms (tremors and abnormal muscle movements) in breastfed infants exposed to risperidone (*see Clinical Considerations*). There is no information on the effects of risperidone on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for PERSERIS and any potential adverse effects on the breastfed child from PERSERIS or from the mother's underlying condition.

Clinical Considerations

Infants exposed to PERSERIS through breastmilk should be monitored for excess sedation, failure to thrive, jitteriness, and extrapyramidal symptoms (tremors and abnormal muscle movements).

8.3 Females and Males of Reproductive Potential

Infertility

Females

Based on the pharmacologic action of risperidone (D₂ receptor antagonism), treatment with PERSERIS may result in an increase in serum prolactin levels, which may lead to a reversible reduction in fertility in females of reproductive potential [see *Warnings and Precautions* (5.6)].

8.4 Pediatric Use

Safety and effectiveness of PERSERIS have not been established in pediatric patients.

8.5 Geriatric Use

Clinical studies of PERSERIS in the treatment of schizophrenia did not include patients aged 65 and older to determine whether or not they respond differently from younger patients.

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Elderly patients with dementia-related psychosis treated with PERSERIS are at an increased risk of death compared to placebo. PERSERIS is not approved for the treatment of patients with dementia related psychosis [see *Boxed Warning and Warnings and Precautions* (5.1, 5.2)].

8.6 Renal Impairment

In patients with renal impairment, carefully titrate with oral risperidone (up to at least 3 mg) before initiating treatment with PERSERIS at a dose of 90 mg [see *Dosage and Administration* (2.2) and *Clinical Pharmacology* (12.3)].

PERSERIS was not studied in patients with renal impairment, however, such effect has been investigated with oral risperidone.

8.7 Hepatic Impairment

In patients with hepatic impairment, carefully titrate with oral risperidone (up to at least 3 mg) before initiating treatment with PERSERIS at a dose of 90 mg [see *Dosage and Administration* (2.2) and *Clinical Pharmacology* (12.3)].

PERSERIS was not studied in patients with hepatic impairment, however, such effect has been investigated with oral risperidone.

10 OVERDOSAGE

10.1 Human Experience

No cases of overdose were reported in premarketing studies with PERSERIS. Because PERSERIS is to be administered by healthcare professionals, the potential for overdose by patients is low.

10.2 Management of Overdosage

In case of overdose, consult a Poison Control Center at 1-800-222-1222.

In case of acute overdose, establish and maintain an airway and ensure adequate oxygenation and ventilation. Cardiovascular monitoring should commence immediately and should include continuous electrocardiographic monitoring to detect possible arrhythmias. If antiarrhythmic therapy is administered, disopyramide, procainamide, and quinidine carry a theoretical hazard of QT prolonging effects that might be additive to those of risperidone. Similarly, it is reasonable to expect that the alpha-blocking properties of bretylium might be additive to those of risperidone, resulting in problematic hypotension.


There is no specific antidote to risperidone. Appropriate supportive measures should be instituted. Hypotension and circulatory collapse should be treated with appropriate measures, such as intravenous fluids and/or sympathomimetic agents (epinephrine and dopamine should not be used, since beta stimulation may worsen hypotension in the setting of risperidone-induced alpha blockade). In cases of severe extrapyramidal symptoms, anticholinergic medication should be administered. Close medical supervision and monitoring should continue until the patient recovers.

Consider the long-acting nature of PERSERIS when assessing treatment needs and recovery.

11 DESCRIPTION

PERSERIS contains risperidone, an atypical antipsychotic. Risperidone belongs to the chemical class of benzisoxazole derivatives. The chemical designation 3-[2-[4-(6-fluoro-1,2-benzisoxazol-3-yl) piperidin-1-yl] ethyl]-2-methyl-6,7,8,9-tetrahydropyrido[1,2-a] pyrimidin-4-one. Its molecular formula is $C_{23}H_{27}FN_4O_2$ and its molecular weight is 410.5 g/mol.

The structural formula is:



Risperidone is a white to off-white powder. It is practically insoluble in water and soluble in methanol and 0.1 N HCl.

PERSERIS is available as a sterile two-syringe mixing system; a liquid syringe prefilled with the delivery system, a colorless to yellow solution. The delivery system provides the monthly extended-release delivery of risperidone in PERSERIS. It is comprised of poly (DL-lactide-co-glycolide) polymer and *N*-methyl-2-pyrrolidone. The powder syringe is prefilled with risperidone (white to yellow). Prior to use, the product is constituted by coupling the liquid and powder syringes and passing the contents back and forth between the syringes [see *Dosage and Administration* (2.4)]. On completion of the mixing cycles, the combined mixture resides in the liquid syringe. A sterile, safety needle is affixed to the liquid syringe and the expressible syringe contents are injected subcutaneously into the abdomen. The product should be prepared immediately prior to use for subcutaneous injection.

After mixing, PERSERIS is available as an extended release injectable suspension, for subcutaneous use, in the following strengths of risperidone: 90 mg and 120 mg.

Table 6. PERSERIS Constituted Product Delivered Mass

Component	PERSERIS 90 mg	PERSERIS 120 mg
Risperidone	90 mg	120 mg
PLGH	228 mg	304 mg
N-methyl-pyrrolidine	282 mg	376 mg
Total mass	600 mg	800 mg
Total volume	0.6 mL	0.8 mL

PLGH poly D,L(lactide co-glycolide); 80:20 molar ratio of lactide to glycolide

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism of action of risperidone, in schizophrenia, is unclear. The drug's therapeutic activity in schizophrenia could be mediated through a combination of dopamine Type 2 (D_2) and serotonin Type 2 ($5HT_2$) receptor antagonism. The clinical effect from risperidone results from the combined concentrations of risperidone and its major metabolite, 9-hydroxyrisperidone (paliperidone) [see *Clinical Pharmacology* (12.3)]. Antagonism at receptors other than D_2 and $5HT_2$ may explain some of the other effects of risperidone.

12.2 Pharmacodynamics

Risperidone is a monoaminergic antagonist with high affinity (K_i of 0.12 to 7.3 nM) for the serotonin Type 2 ($5HT_2$), dopamine Type 2 (D_2), α_1 and α_2 adrenergic, and H_1 histaminergic receptors. Risperidone showed low to moderate affinity (K_i of 47 to 253 nM) for the serotonin $5HT_{1C}$, $5HT_{1D}$, and $5HT_{1A}$ receptors, weak affinity (K_i of 620 to 800 nM) for the dopamine D_1 and haloperidol-sensitive sigma site, and no affinity (when tested at concentrations $> 10^{-5}$ M) for cholinergic muscarinic or β_1 and β_2 adrenergic receptors.

12.3 Pharmacokinetics

The pharmacokinetics of risperidone and total active moiety following subcutaneous injection of PERSERIS was evaluated in subjects with clinically stable schizophrenia after single doses (60 mg, 90 mg, and 120 mg) ($n = 101$) and repeated doses (60 mg, 90 mg, and 120 mg) ($n = 45$) separated by 28 days for up to 3 injections following oral risperidone. Risperidone plasma concentrations had a T_{max} of 4 to 6 hours and approached steady-state levels after the first subcutaneous injection of PERSERIS. Similar pattern was observed for 9-hydroxyrisperidone and total active moiety. Steady-state plasma concentrations were reached by the end of the second injection for risperidone, 9-hydroxyrisperidone, and total active moiety and were maintained for 4 weeks after the last injection. Mean accumulation ratios for risperidone ranged from 1.2 to 1.7 based on AUC, and from 0.9 to 1.3 based on overall C_{max} , indicating no or modest accumulation. For 9-hydroxyrisperidone, accumulation ratios ranged from 1.2 to 1.6 (AUC) and 0.99 to 1.3 (overall C_{max}). For total active moiety, accumulation ratios ranged from 1.2 to 1.6 (AUC_{tau}) and 0.97 to 1.3 (overall C_{max}).

Total active moiety concentrations reached clinically relevant levels after the first injection without use of a loading dose or any supplemental oral risperidone.

Following multiple doses of PERSERIS, plasma exposure (AUC_{tau} and C_{max}) of risperidone, 9-hydroxyrisperidone, and total active moiety increased in an approximately dose proportional manner over the dose range of 60 to 120 mg. At steady-state, a 2-fold increase in dose resulted in a 1.7-fold increase in C_{max} (6.33 to 10.9 ng/mL) and AUC_{tau} (2262 to 3891 ng*hr/mL) for risperidone. For 9-hydroxyrisperidone, a 2-fold increase in dose resulted in a 2.1-fold increase in C_{max} (13.7 to 28.9 ng/mL) and 2-fold increase in AUC_{tau} (5706 to 11658 ng*hr/mL). For total active moiety, a 2-fold increase in

dose resulted in a 2.0-fold increase in $C_{\max}$ (19.6 to 38.5 ng/mL) and a 1.9-fold increase in AUC_{τ} (8102 to 15370 ng*hr/mL).

Plasma exposures at steady-state were compared between oral risperidone and PERSERIS. Based on average plasma concentrations (C_{avg}) of risperidone and total active moiety, 90 mg PERSERIS corresponds to 3 mg oral risperidone and 120 mg PERSERIS corresponds to 4 mg oral risperidone.

Absorption

PERSERIS contains risperidone in a liquid delivery system. Following subcutaneous injection, it forms a depot that provides sustained plasma levels of risperidone over the monthly dosing interval.

After single subcutaneous injection, PERSERIS shows two absorption peaks for risperidone in plasma. The first peak of risperidone occurs with a $T_{\max}$ of 4 to 6 hours and is due to an initial release of the drug during the depot formation process. A second peak of risperidone is observed at 10 to 14 days post-dose and is associated with the slow release of risperidone from the subcutaneous depot. The first and second peaks of risperidone are of similar magnitude. For both 9-hydroxyrisperidone and total active moiety, the median $T_{\max}$ of the first peak ranges from 4 to 48 hours and the second peak ranges from 7 to 11 days.

Distribution

Following a subcutaneous injection of PERSERIS, the apparent volume of distribution is large. The extensively large values are because PERSERIS is administered as a depot injection. Risperidone is bound to albumin and α 1-acid glycoprotein. The plasma protein binding of risperidone is approximately 90%, and that of its major metabolite, 9-hydroxyrisperidone, is 77%. Neither risperidone nor 9-hydroxyrisperidone displace each other from plasma binding sites.

Elimination

Metabolism

Risperidone is extensively metabolized in the liver. The main metabolic pathway is through hydroxylation of risperidone to 9-hydroxyrisperidone by the enzyme cytochrome CYP2D6 with minor contribution by CYP3A4. A minor metabolic pathway is through N-dealkylation. The main metabolite, 9-hydroxyrisperidone, has similar pharmacological activity as risperidone. Consequently, the clinical effect of the drug results from the combined concentrations of risperidone plus 9-hydroxyrisperidone).

CYP2D6, is the enzyme responsible for metabolism of many neuroleptics, antidepressants, antiarrhythmics, and other drugs. CYP2D6 is subject to genetic polymorphism (about 6 to 8% of Caucasians, and a very low percentage of Asians, have little or no activity and are “poor metabolizers”) and to inhibition by a variety of substrates and some non-substrates, notably quinidine. Extensive CYP2D6 metabolizers convert risperidone rapidly into 9-hydroxyrisperidone, whereas poor CYP2D6 metabolizers convert it much more slowly. Plasma exposure to total active moiety was similar in CYP2D6 extensive, intermediate and poor metabolizers following subcutaneous injection with PERSERIS, supporting no need for dose adjustment based on genotype of CYP2D6.

Excretion

Risperidone and its metabolites are eliminated via the urine and, to a much lesser extent, via the feces. As illustrated by a mass balance study of a single 1 mg oral dose of ^{14}C -risperidone administered as solution to 3 healthy male volunteers, total recovery of radioactivity at 1 week was 84%, including 70% in the urine and 14% in the feces.

Following a single subcutaneous injection of PERSERIS, the apparent terminal half-life of risperidone ranges between 9 and 11 days on average. This half-life is related to the slow release of risperidone from

the subcutaneous depot and subsequent absorption of risperidone into the systemic circulation. The mean apparent terminal half-life ranges between 8 to 9 days for both 9-hydroxyrisperidone and total active moiety, on average.

Drug Interaction Studies

No specific drug interaction studies have been performed with PERSERIS. The drug interaction data provided in this section is based on studies with oral risperidone. Effects of other drugs on the exposures of risperidone, 9-hydroxyrisperidone and total active moiety as well as the effects of risperidone on the exposures of other drugs is summarized below.

Effects of Other Drugs on Risperidone, 9-hydroxyrisperidone and Total Active Moiety Pharmacokinetics

Strong CYP2D6 Inhibitors (Fluoxetine and Paroxetine)

Fluoxetine (20 mg once daily) and paroxetine (20 mg once daily), potent CYP2D6 inhibitors, have been shown to increase the plasma concentration of risperidone by 2.5- to 2.8-fold and 3- to 9-fold, respectively. Fluoxetine did not affect the plasma concentration of 9-hydroxyrisperidone. Paroxetine lowered the concentration of 9-hydroxyrisperidone by about 10%. The effects of discontinuation of concomitant fluoxetine or paroxetine therapy on the pharmacokinetics of risperidone and 9-hydroxyrisperidone have not been studied.

Moderate CYP3A4 Inhibitor (Erythromycin)

There were no significant interactions between oral risperidone and erythromycin, a moderate CYP3A4 inhibitor.

Strong CYP3A4 Inducer (Carbamazepine)

Carbamazepine co-administration with oral risperidone decreased the steady-state plasma concentrations of risperidone and 9-hydroxyrisperidone by about 50%. Plasma concentrations of carbamazepine did not appear to be affected. Co-administration of other known CYP3A4 enzyme inducers (e.g., phenytoin, rifampin, and phenobarbital) with risperidone may cause similar decreases in the combined plasma concentrations of risperidone and 9-hydroxyrisperidone, which could lead to decreased efficacy of PERSERIS.

Amitriptyline, Cimetidine, Ranitidine, Clozapine, Topiramate

Clinically meaningful pharmacokinetic interaction between PERSERIS and other drugs, such as amitriptyline, cimetidine, ranitidine and clozapine, is not expected.

- Amitriptyline did not affect the pharmacokinetics of risperidone or of risperidone and 9-hydroxyrisperidone combined following concomitant administration with oral risperidone.
- Cimetidine and ranitidine increased the bioavailability of oral risperidone by 64% and 26%, respectively. However, cimetidine did not affect the AUC of risperidone and 9-hydroxyrisperidone combined, whereas ranitidine increased the AUC of risperidone and 9-hydroxyrisperidone combined by 20%.
- Chronic administration of clozapine with oral risperidone have shown to affect the clearance of risperidone, however, clinical relevance is unknown.
- There was no clinically relevant effect of oral risperidone (1 to 6 mg/day) on the pharmacokinetics of topiramate 400 mg/day.

Effects of Oral Risperidone on Pharmacokinetics of Other Drugs

Lithium

Repeated doses of oral risperidone (3 mg twice daily) did not affect the exposure (AUC) or peak plasma concentrations ($C_{\max}$) of lithium (n = 13).

Valproate

Repeated doses of oral risperidone (4 mg once daily) did not affect the pre-dose or average plasma concentrations and exposure (AUC) of valproate (1000 mg/day in three divided doses) compared to placebo (n = 21). However, there was a 20% increase in valproate peak plasma concentration ($C_{\max}$) after concomitant administration of oral risperidone.

Topiramate

Oral risperidone administered at doses from 1 to 6 mg/day concomitantly with topiramate 400 mg/day resulted in a 23% decrease in risperidone $C_{\max}$ and a 33% decrease in risperidone AUC_{0-12} hour at steady state. Minimal reductions in the exposure to risperidone and 9-hydroxyrisperidone combined, and no change for 9-hydroxyrisperidone were observed. This interaction is unlikely to be of clinical significance. There was no clinically relevant effect of oral risperidone on the pharmacokinetics of topiramate.

Digoxin

Oral risperidone (0.25 mg twice daily) did not show a clinically relevant effect on the pharmacokinetics of digoxin.

CYP2D6 Substrates (Donepezil and Galantamine)

In vitro studies indicate that risperidone is a relatively weak inhibitor of CYP2D6. Therefore, PERSERIS is not expected to substantially inhibit the clearance of drugs that are metabolized by this enzymatic pathway. In drug interaction studies, oral risperidone did not significantly affect the pharmacokinetics of donepezil and galantamine, which are metabolized by CYP2D6.

Specific Populations

Based on population pharmacokinetic analyses, age, sex and race do not have a clinically meaningful effect on the pharmacokinetics of PERSERIS.

Renal Impairment

PERSERIS was not studied in patients with renal impairment, however, such effect has been investigated with oral risperidone. In patients with moderate to severe renal disease treated with oral risperidone, the apparent clearance (CL/F) of total active moiety was decreased by 60% in patients with moderate to severe renal disease compared with young healthy subjects [see *Use in Specific Populations* (8.6)].

Hepatic Impairment

The effect of hepatic impairment on the pharmacokinetics of PERSERIS has not been studied.

The effect of hepatic impairment on the pharmacokinetics of oral risperidone has been evaluated in a dedicated phase I study. While the pharmacokinetics of risperidone in subjects with liver disease were comparable to those in young healthy subjects, the mean free fraction of risperidone in plasma was increased by about 35% because of the diminished concentration of both albumin and α_1 -acid glycoprotein [see *Use in Specific Populations* (8.7)].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

No carcinogenicity studies were conducted with subcutaneous risperidone suspension. Carcinogenicity studies were conducted with oral risperidone in mice and rats. Risperidone was administered in the diet at doses of 0.63, 2.5, and 10 mg/kg for 18 months to mice and for 25 months to rats. These doses are equivalent to approximately 0.2, 0.75, and 3 times (mice) and 0.4, 1.5, and 6 times (rats) the MHRD of 16 mg/day, based on a mg/m² body surface area. A maximum tolerated dose was not achieved in male mice. There were statistically significant increases in pituitary gland adenomas, endocrine pancreas adenomas, and mammary gland adenocarcinomas. The table below summarizes the multiples of the human dose on a mg/m² (mg/kg) basis at which these tumors occurred.

Table 7. Summary of Tumor Occurrence at the Multiples of the Human Dose on a mg/m² (mg/kg) Basis with Oral Risperidone Dosing

Tumor Type	Species	Sex	Multiples of Maximum Human Dose in mg/m ² (mg/kg)	
			Lowest Effect Level	Highest No-Effect Level
Pituitary adenomas	mouse	Female	0.75 (9.4)	0.2 (2.4)
Endocrine pancreas adenomas	rat	Male	1.5 (9.4)	0.4 (2.4)
Mammary gland adenocarcinomas	mouse	Female	0.2 (2.4)	none
	rat	Female	0.4 (2.4)	none
	rat	Male	6.0 (37.5)	1.5 (9.4)
Mammary gland neoplasm, Total	rat	Male	1.5 (9.4)	0.4 (2.4)

Antipsychotic drugs have been shown to chronically elevate prolactin levels in rodents. Serum prolactin levels were not measured during the risperidone carcinogenicity studies; however, measurements during subchronic toxicity studies showed that risperidone elevated serum prolactin levels 5- to 6-fold in mice and rats at the same doses used in the carcinogenicity studies. An increase in mammary, pituitary, and endocrine pancreas neoplasms has been found in rodents after chronic administration of other antipsychotic drugs and is considered to be prolactin-mediated. The relevance for human risk of the findings of prolactin-mediated endocrine tumors in rodents is unclear [see *Warnings and Precautions* (5.6)].

Mutagenesis

No evidence of mutagenic or clastogenic potential for risperidone was found in the *in vitro* tests of Ames gene mutation, the mouse lymphoma assay, rat hepatocyte DNA-repair assay, the chromosomal aberration test in human lymphocytes, Chinese hamster ovary cells, or in the *in vivo* oral micronucleus test in mice and the sex-linked recessive lethal test in *Drosophila*.

No evidence of mutagenic potential was observed with risperidone subcutaneous injectable suspension or its delivery system alone at doses of 150 mg/kg risperidone or 943 mg/kg delivery system in an *in vivo* micronucleus test in rats. The safety margins of risperidone were 12 to 19 times the maximum monthly plasma risperidone concentration observed for humans at the monthly MRHD of 120 mg risperidone based on plasma exposure, and 13 times the delivery system amount present in monthly 120 mg risperidone.

Impairment of Fertility

No mating and fertility studies were conducted with subcutaneous risperidone suspension. Oral risperidone (0.16 to 5 mg/kg) impaired mating, but not fertility, in rat reproductive studies at doses 0.1 to 3 times the maximum recommended human dose (MRHD), of 16 mg/day based on mg/m² body surface

area. The effect appeared to be in females, since impaired mating behavior was not noted in the male fertility study. In a subchronic study in Beagle dogs in which risperidone was administered orally at doses of 0.31 to 5 mg/kg, sperm motility and concentration were decreased at doses 0.6 to 10 times the MRHD based on mg/m² body surface area. Dose-related decreases were also noted in serum testosterone at the same doses. Serum testosterone and sperm parameters partially recovered, but remained decreased after treatment was discontinued. A no-effect dose could not be determined in either rat or dog.

Subcutaneous administration of the delivery system to rats had no effect on fertility parameters in either sex up to a dose that is 17 (delivery system), and 23 (NMP) times the amount present in monthly 120 mg risperidone subcutaneous injectable suspension based on mg/m² body surface area, respectively.

14 CLINICAL STUDIES

Efficacy for PERSERIS was demonstrated in an 8-week, randomized, double-blind, placebo-controlled study (Study 1, NCT #02109562). The study evaluated the efficacy, safety and tolerability of PERSERIS (90 and 120 mg subcutaneous every 4 weeks) compared with placebo in adults (age 18 to 55 years, inclusive) experiencing acute exacerbations of schizophrenia. Patients were required to have a Positive and Negative Syndrome Scale (PANSS) total score of 80 to 120 inclusive (moderate to severely ill) at the screening visit, occurring 3 to 8 days before the start of double-blind treatment, without an improvement in the PANSS total score of $\geq 20\%$ between screening and the first dosing day.

At the screening visit, all patients received two doses of 0.25 mg oral risperidone 24 hours apart to establish tolerability. Patients were then placed in an inpatient setting, if not already hospitalized, and tapered off their current oral antipsychotic medication (if they were taking one) over a period of 3 to 8 days. Patients were randomized to receive 2 doses of subcutaneous PERSERIS (90 mg or 120 mg) or placebo 28 days apart (on Day 1 and Day 29). No supplemental oral risperidone was permitted during the study.

The primary endpoint was the change in PANSS total score from baseline to end of study (Day 57). Both PERSERIS 90 and 120 mg doses demonstrated a statistically significant improvement compared with placebo based on the primary endpoint (Table 8). The results at each scheduled visit are displayed in Figure 14.

Characteristics of the patient population were balanced across the treatment groups. The mean baseline PANSS total score ranged from 94 to 96 across the groups. Most patients were male (74 to 83% per group), and the mean ages were 40 to 43 in each group. Most patients in this study were black or African American (71 to 75% per group). Of the 354 subjects randomized to treatment, 337 were included in the intent-to-treat (ITT) population, and 259 (73%) completed the study.

Subgroup analyses by gender, age, and race did not suggest any clear evidence of differential responsiveness to PERSERIS.

Table 8: Primary Efficacy Analysis Results for Study 1

Treatment Group	N (# ITT subjects)	Primary Efficacy Measure: PANSS		
		Mean Baseline Score (SD)	LS Mean Change from Baseline (SE)	Placebo-subtracted Difference ^a (95% CI)
PERSERIS 90 mg*	111	95.5 (9.23)	-19.86 (1.56)	-6.50 (-10.87, -2.13) *
PERSERIS 120 mg*	114	94.9 (8.09)	-23.61 (1.58)	-10.24 (-14.64, -5.85) *


Placebo 112 94.1 (8.89) -13.37 (1.58) --

ITT: intent-to-treat; SD: standard deviation; SE: standard error; LS Mean: least-squares mean; CI: unadjusted confidence interval

*Difference (drug minus placebo) in least-squares mean change from baseline

*Doses that are statistically significantly superior to placebo

Figure 14. Least Square Mean Change from Baseline (+/- Standard Error) in PANSS Total Scores by Days



The secondary efficacy endpoint was defined as the CGI-S score at Day 57. Both PERSERIS treatment groups demonstrated statistically significantly better CGI-S scores versus placebo.

16 HOW SUPPLIED/STORAGE AND HANDLING

PERSERIS (risperidone) for extended-release injectable suspension, for subcutaneous use is, when fully mixed, a viscous suspension that varies from white to yellow-green and is available in dosage strengths of 90 mg and 120 mg.

PERSERIS 90 mg is supplied in a single-dose kit, packaged in a carton (NDC 12496-0090-1), containing the following:

- One pouch with a sterile syringe (labelled 'P') prefilled with risperidone powder
- One pouch with a sterile syringe (labelled 'L') prefilled with the delivery system, and desiccant.
- One 18-gauge, 5/8-inch sterile safety needle.

PERSERIS 120 mg is supplied in a single-dose kit, packaged in a carton (NDC 12496-0120-1), containing the following:

- One pouch with a sterile syringe (labelled ‘P’) prefilled with risperidone powder.
- One pouch with a sterile syringe (labelled ‘L’) prefilled with the delivery system, and desiccant.
- One 18-gauge, 5/8-inch sterile safety needle.

Storage and Handling

Store in refrigerator at 2° to 8°C (36° to 46°F). Allow PERSERIS kit to come to room temperature, 20°C to 25°C (68°F to 77°F), for at least 15 minutes prior to mixing.

PERSERIS may be stored in its unopened original packaging at room temperature, 20°C to 25°C (68°F to 77°F), for up to 7 days prior to administration. After removal from the refrigerator, use PERSERIS within 7 days or discard.

17 PATIENT COUNSELING INFORMATION

Physicians are advised to discuss with patients for whom they prescribe PERSERIS all relevant safety information including, but not limited to, the following:

Instructions for General Use

Advise patients not to rub or massage the injection site and to be aware of the placement of any belts or clothing waistbands [*see Dosage and Administration (2.4)*].

Orthostatic Hypotension

Advise patients of the risk of orthostatic hypotension and instruct in nonpharmacologic interventions that help to reduce the occurrence of orthostatic hypotension (e.g., sitting on the edge of the bed for several minutes before attempting to stand in the morning and slowly rising from a seated position) [*see Warnings and Precautions (5.7)*].

Potential for Cognitive and Motor Impairment

Inform patients that PERSERIS has the potential to impair judgment, thinking, or motor skills. Advise caution about operating hazardous machinery, including automobiles, until they are reasonably certain that treatment with PERSERIS does not affect them adversely [*see Warnings and Precautions (5.10)*].

Concomitant Medication

Advise patients to inform their physicians if they are taking, or plan to take, any prescription or over-the-counter drugs, since there is a potential for interaction [*see Drug Interactions (7)*].

Alcohol

Advise patients to avoid alcohol during treatment with PERSERIS [*see Drug Interactions (7.1)*].

Metabolic Changes

Advise patients that treatment with PERSERIS can be associated with diabetes mellitus and hyperglycemia, dyslipidemia and weight gain [*see Warning and Precautions (5.5)*].

Pregnancy

Advise patients to notify their healthcare professional if they become pregnant or intend to become pregnant during treatment with PERSERIS. Advise patients that PERSERIS may cause extrapyramidal and/or withdrawal symptoms in a neonate. Advise patients that there is a pregnancy registry that monitors

pregnancy outcomes in women exposed to PERSERIS during pregnancy *[see Use in Specific Populations (8.1)]*.

Lactation

Advise breastfeeding women using PERSERIS to monitor infants for somnolence, failure to thrive, jitteriness, and extrapyramidal symptoms (tremors and abnormal muscle movements) and to seek medical care if they notice these signs *[see Use in Specific Populations (8.2)]*.

Infertility

Advise females of reproductive potential that PERSERIS may impair fertility due to an increase in serum prolactin levels. The effects on fertility are reversible *[see Use in Specific Populations (8.3)]*.

Manufactured for:

Indivior Inc., North Chesterfield, VA 23235.

Powder syringe manufactured by Patheon Manufacturing Services, Greenville, NC 27834.

Liquid syringe manufactured by AMRI Global, Burlington, MA 01803.

PERSERIS™ is a trademark of Indivior UK Limited. All rights reserved.

INSTRUCTIONS FOR USE

PERSERIS™ (risperidone) for extended-release injectable suspension

90 mg or 120 mg, sterile, single dose

For abdominal subcutaneous injection only.

Do not administer by any other route.


Additional information, including an instructional video, are available at perserishcp.com. The additional information has not been evaluated or approved by the FDA.

IMPORTANT INFORMATION:

- To be administered by a healthcare professional only.
- Please read the instructions carefully before handling this product.
- Allow package to come to room temperature for at least 15 minutes prior to preparation.
- Only prepare medication when you are ready to administer the dose.
- As a universal precaution, always wear gloves.

1 CHECK CONTENTS

- One **Liquid Syringe** (**L**) pre-filled with the delivery system – Inspect liquid solution for foreign particles. This is the syringe you will use to inject the patient.
- One **Powder Syringe** (**P**) prefilled with Risperidone powder – Inspect syringe for consistency of powder color and for foreign particles.
- One sterile 18-gauge, 5/8 inch safety needle
- Parenteral drug products should always be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.




2 TAP POWDER SYRINGE

- Hold the **Powder Syringe** upright and tap the barrel of the syringe to dislodge the packed powder.

NOTE

Powder can become packed during shipping.




3 UNCAP LIQUID AND POWDER SYRINGES

- Remove the cap from the **Liquid Syringe**, then remove the cap from the **Powder Syringe**.

TIP

Holding both syringes in your non-dominant hand can help with this step.




4 CONNECT THE SYRINGES

- Place the **Liquid Syringe** on top of the **Powder Syringe** (to prevent powder spillage) and connect the syringes by twisting approximately $\frac{3}{4}$ turn. Do not over tighten.

CAUTION

Keep your fingers off the plungers during this step to avoid spillage of the medication.



5 MIX THE PRODUCT

WARNING

Failure to fully mix the medication could result in incorrect dosage.

Premixing:



- Transfer the contents of the **Liquid Syringe** into the **Powder Syringe**.
- Gently push the **Powder Syringe** plunger until you feel resistance (to wet powder and avoid compacting).
- Repeat this gentle back-and-forth process for **5 Cycles**.

Complete mixing:

- Continue mixing the syringes for an additional **55 Cycles**.
- This mixing can be more vigorous than when premixing.

TIP

Graphic to the right illustrates a correct full cycle.



6 PREPARE INJECTION SYRINGE


WARNING

Failure to aspirate the liquid from the **Powder Syringe** may result in incorrect dosage.

- First, transfer all contents into the **Liquid Syringe**.
- Next, perform the following actions **SIMULTANEOUSLY**:
 - maintain slight pressure on the **Powder Syringe** plunger and
 - pull back **gently** on the **Liquid Syringe** plunger while twisting the syringes apart.
- Finally, attach the safety needle by twisting until finger tight.

NOTE

Check that medication is uniform in color and free from foreign particles.



Turn the Page 

PERSERIS™


Instructions for Use

7 PREPARE THE ABDOMINAL INJECTION SITE

- Choose an injection site on the abdomen with adequate subcutaneous tissue that is free of skin conditions (e.g. nodules, lesions, excessive pigment). It is recommended that the patient is in the supine position.
- **Do not** inject into an area where the skin is irritated, reddened, bruised, infected or scarred in any way.
- Clean the injection site well with an alcohol pad.

TIP

To help minimize irritation, rotate injection sites following a pattern similar to the illustration.



8 REMOVE EXCESS AIR FROM SYRINGE

- Hold the syringe upright for several seconds to allow air bubbles to rise.
- Remove needle cover and slowly depress the plunger to push out the excess air from the syringe.
- If medication is seen at the needle tip, pull back slightly on the plunger to prevent medication spillage.


NOTE

Due to the viscous nature of the medication, bubbles will not rise as quickly as those in an aqueous solution.



9 PINCH INJECTION SITE

- Pinch the skin around the injection area. Be sure to pinch enough skin to accommodate the size of the needle. Lift the adipose tissue from the underlying muscle to prevent accidental intramuscular injection.



10 INJECT THE MEDICATION


- Insert needle fully into the subcutaneous tissue.
- Inject the medication slow and steady.



PERSERIS™ is for subcutaneous administration only. Do not inject by any other route.

NOTE

Actual angle of injection will depend on the amount of subcutaneous tissue.




11 WITHDRAW NEEDLE

- Withdraw the needle at the same angle used for insertion and release pinched skin.




Do not rub the injection area after the injection. If there is bleeding, apply a gauze pad or bandage but use minimal pressure.



12 LOCK THE NEEDLE GUARD AND DISPOSE OF SYRINGE

- Lock the needle guard into place by pushing it against a hard surface such as a table.
- Dispose of all syringe components in a secure sharps disposal container.



13 INSTRUCT THE PATIENT

- Advise the patient that they may have a lump for several weeks that will decrease in size over time. It is important that the patient not rub or massage the injection site and to be aware of the placement of any belts or clothing waistbands.



FREQUENTLY ASKED QUESTIONS

Q What is the mixture supposed to look like?

A When fully mixed, the product should be a cloudy suspension that is uniform in color. It can vary from white to yellow-green in color. If you see any clear areas in the mixture, continue to mix until the distribution of the color is uniform.

Q Why is it important to put the liquid syringe on top?

A The liquid is viscous and will not fall out when placing on top. However, powder may be lost if the powder syringe is inverted, which could affect the ultimate dosage.

Q What do I do if there is residual left in the powder syringe?

A A small ring of residual powder may be observed in the end of the powder syringe barrel after mixing. This is normal. If you have followed the instructions carefully and the rest of the mixture is a cloudy suspension that is uniform in color, proceed with the injection.

Q What do I do if a foreign particle is found in either of the syringes?

A Do not use if you suspect foreign particles in either of the syringes.

Q How soon must I inject after mixing PERSERIS™?

A This product should be used immediately following preparation.

Q Can I inject into the arm or leg?

A This injection is only approved for injection into the abdomen.

Q How do I get rid of large air gaps?

A Small bubbles, also known as champagne bubbles, are not a problem and common with this medication. Large air gaps, however, can be minimized by pulling back on the plunger rod to pop air bubbles prior to expelling the air very slowly. Air should be expelled very carefully to avoid loss of medication.

Q Should I massage or put my finger on the injection site following the injection?

A It is not advisable to palpate or massage the area following the injection.

Q Will the deposit be palpable?

A Depending on the patient's subcutaneous tissue, the deposit may be more or less palpable. Patients should be advised that a bump may be palpable (decreasing in size) for several weeks.

Manufactured for: Indivior Inc., North Chesterfield, VA 23235.

Powder syringe manufactured by Patheon Manufacturing Services, Greenville, NC 27834.

Liquid syringe manufactured by AMRI Global, Burlington, MA 01803.

PERSERIS™ is a trademark of Indivior UK Limited.

All rights reserved.

NDC 12496-0120-2



120 mg

**For abdominal subcutaneous
injection only.**


**Please read complete
instructions prior to use.**

**Please see accompanying
Full Prescribing Information.**

Rx only

Sterile

Single-dose only



POUCH CONTENTS

Powder Syringe: Prefilled with risperidone powder designed to deliver 120 mg risperidone after mixing with the delivery system. Allow pouch to come to room temperature for at least 15 minutes prior to mixing. Contents of Powder Syringe must be mixed with liquid provided in the other prefilled Liquid Syringe, prior to use.



Manufactured for Indivior Inc., North Chesterfield, VA 23235
by Patheon Mfr. Services, Greenville, NC 27834
Rev. July 2018

NDC 12496-0776-2

Rx only

Sterile

Single-dose only

Delivery System



for **PERSERIS™ 120 mg**

For abdominal subcutaneous injection only.

Please read complete instructions prior to use.

Please see accompanying Full Prescribing Information.

POUCH CONTENTS

Liquid Syringe: Prefilled with delivery system designed to deliver 0.8 mL containing inactive ingredients poly D,L (lactide co-glycolide) with an 80:20 molar ratio of lactide to glycolide (304 mg) and *N*-methyl-pyrrolidone (376 mg) after mixing with powder syringe. Allow pouch to come to room temperature for at least 15 minutes prior to mixing. Contents of Liquid Syringe must be mixed with powder provided in the other prefilled Powder Syringe, prior to use.



Manufactured for Indivior Inc., North Chesterfield, VA 23235
by AMRI Global, Burlington, MA 01803
Rev. July 2018

NDC 12496-0090-2



90 mg

**For abdominal subcutaneous
injection only.**


**Please read complete
instructions prior to use.**

**Please see accompanying
Full Prescribing Information.**

Rx only

Sterile

Single-dose only



POUCH CONTENTS

Powder Syringe: Prefilled with risperidone powder designed to deliver 90 mg risperidone after mixing with the delivery system. Allow pouch to come to room temperature for at least 15 minutes prior to mixing. Contents of Powder Syringe must be mixed with liquid provided in the other prefilled Liquid Syringe, prior to use.



Manufactured for Indivior Inc., North Chesterfield, VA 23235
by Patheon Mfr. Services, Greenville, NC 27834
Rev. July 2018

NDC 12496-0589-2

Rx only

Sterile

Single-dose only

Delivery System



for PERSERIS™ 90 mg

For abdominal subcutaneous injection only.

Please read complete instructions prior to use.

Please see accompanying Full Prescribing Information.

POUCH CONTENTS

Liquid Syringe: Prefilled with delivery system designed to deliver 0.6 mL containing inactive ingredients poly D,L (lactide co-glycolide) with an 80:20 molar ratio of lactide to glycolide (228 mg) and *N*-methyl-pyrrolidone (282 mg) after mixing with powder syringe. Allow pouch to come to room temperature for at least 15 minutes prior to mixing. Contents of Liquid Syringe must be mixed with powder provided in the other prefilled Powder Syringe, prior to use.



Manufactured for Indivior Inc., North Chesterfield, VA 23235
by AMRI Global, Burlington, MA 01803
Rev. July 2018

Sterile

PERSERIS™

(risperidone)

120 mg

**For abdominal
subcutaneous
injection only.**



Powder Syringe

Mfd. by Patheon Mfr. Services

FOR POSITIVE ONLY	IN
123456789	123456789

LOT XXXXXXXXX

EXP XX/XXXX

Unvarnished area for wrap

NDC 12496-0776-5

Rx only

Sterile

Delivery System

for PERSERIS™ 120 mg

For abdominal
subcutaneous
injection only.

L

Liquid
Syringe

FOR POSITIVE
ONLY

120 mg/mL

LOT XXXXXXXX
EXP XX/XXXX

Unvarnished area for wrap

Mfd. by AMRI Global

NDC 12496-0090-5

Rx only

Sterile

PERSERIS™
(risperidone)

90 mg

**For abdominal
subcutaneous
injection only.**



**Powder
Syringe**

Mfd. by Patheon Mfr.
Services



LOT XXXXXXXX
EXP XX/XXXX

Unvarnished area for wrap

NDC 12496-0589-5

Rx only

Sterile

Delivery System

for PERSERIS™ 90 mg

For abdominal
subcutaneous
injection only.

L

Liquid
Syringe

FOR POSITIVE
ONLY

12.88 mm dia

LOT XXXXXXXXX
EXP XX/XXXX

Unvarnished area for wrap

Mfd. by AMRI Global

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

MITCHELL V Mathis
07/27/2018