

# CENTER FOR DRUG EVALUATION AND RESEARCH

## Approval Package for:

### *APPLICATION NUMBER:*

**017862Orig1s063**

*Trade Name:* REGLAN

*Generic or Proper Name:* metoclopramide injection

*Sponsor:* Baxter Healthcare Corporation

*Approval Date:* November 18, 2010

*Indication:* **Diabetic Gastroparesis (Diabetic Gastric Stasis)**  
REGLAN is indicated for the relief of symptoms associated with acute and recurrent diabetic gastric stasis.

**The Prevention of Nausea and Vomiting Associated with Emetogenic Cancer Chemotherapy**

REGLAN Injection is indicated for the prophylaxis of vomiting associated with emetogenic cancer chemotherapy.

**The Prevention of Postoperative Nausea and Vomiting**

REGLAN Injection is indicated for the prophylaxis of postoperative nausea and vomiting in those circumstances where nasogastric suction is undesirable.

**Small Bowel Intubation**

REGLAN Injection may be used to facilitate small bowel intubation in adults and pediatric patients in whom the tube does not pass the pylorus with conventional maneuvers.

**Radiological Examination**

REGLAN Injection may be used to stimulate gastric emptying and intestinal transit of barium in cases where delayed emptying interferes with radiological examination of the stomach and/or small intestine.

# CENTER FOR DRUG EVALUATION AND RESEARCH

## 017862Orig1s063

### CONTENTS

|                                                           |
|-----------------------------------------------------------|
| <b>Reviews / Information Included in this NDA Review.</b> |
|-----------------------------------------------------------|

|                                                         |          |
|---------------------------------------------------------|----------|
| <b>Approval Letter</b>                                  | <b>X</b> |
| <b>Other Action Letters</b>                             |          |
| <b>Labeling</b>                                         | <b>X</b> |
| <b>REMS</b>                                             |          |
| <b>Summary Review</b>                                   |          |
| <b>Officer/Employee List</b>                            |          |
| <b>Office Director Memo</b>                             |          |
| <b>Cross Discipline Team Leader Review</b>              |          |
| <b>Medical Review(s)</b>                                |          |
| <b>Chemistry Review(s)</b>                              |          |
| <b>Environmental Assessment</b>                         |          |
| <b>Pharmacology Review(s)</b>                           |          |
| <b>Statistical Review(s)</b>                            |          |
| <b>Microbiology / Virology Review(s)</b>                |          |
| <b>Clinical Pharmacology/Biopharmaceutics Review(s)</b> |          |
| <b>Other Reviews</b>                                    | <b>X</b> |
| <b>Risk Assessment and Risk Mitigation Review(s)</b>    |          |
| <b>Proprietary Name Review(s)</b>                       |          |
| <b>Administrative/Correspondence Document(s)</b>        |          |

**CENTER FOR DRUG EVALUATION AND  
RESEARCH**

*APPLICATION NUMBER:*

**017862Orig1s063**

**APPROVAL LETTER**



DEPARTMENT OF HEALTH AND HUMAN SERVICES

Food and Drug Administration  
Silver Spring MD 20993

NDA 17-862/S-063

**SUPPLEMENT APPROVAL**

Baxter Healthcare Corporation  
Attention: Eleanor Lescano  
Specialist, Global Regulatory Affairs  
2 Esterbrook Lane  
Cherry Hill, NJ 08003-4099

Dear Ms. Lescano:

Please refer to your Supplemental New Drug Application (sNDA) dated September 24, 2010, received September 27, 2010, submitted under section 505(b) of the Federal Food, Drug, and Cosmetic Act (FDCA) for Reglan (metoclopramide) Injection.

This "Prior Approval" supplemental new drug application provides for revisions to the **Warnings, Tardive Dyskinesia** section of the package insert per our July 20, 2010, request.

We have completed our review of this supplemental application. It is approved, effective on the date of this letter, for use as recommended in the enclosed, agreed-upon labeling text.

**CONTENT OF LABELING**

As soon as possible, but no later than 14 days from the date of this letter, submit, using the FDA automated drug registration and listing system (eLIST), the content of labeling [21 CFR 314.50(l)] in structured product labeling (SPL) format, as described at <http://www.fda.gov/ForIndustry/DataStandards/StructuredProductLabeling/default.htm>, that is identical to the enclosed labeling and include the labeling changes proposed in any pending "Changes Being Effected" (CBE) supplements. Information on submitting SPL files using eLIST may be found in the guidance for industry titled "SPL Standard for Content of Labeling Technical Qs and As" at <http://www.fda.gov/downloads/DrugsGuidanceComplianceRegulatoryInformation/Guidances/UCM072392.pdf>.

The SPL will be accessible from publicly available labeling repositories.

Also within 14 days, amend all pending supplemental applications for this NDA, including pending "Changes Being Effected" (CBE) supplements, for which FDA has not yet issued an

action letter, with the content of labeling [21 CFR 314.50(l)(1)(i)] in MS Word format that includes the changes approved in this supplemental application.

### **LETTERS TO HEALTH CARE PROFESSIONALS**

If you decide to issue a letter communicating important safety-related information about this drug product (i.e., a “Dear Health Care Professional” letter), we request that you submit, at least 24 hours prior to issuing the letter, an electronic copy of the letter to this NDA to the following address:

MedWatch Program  
Office of Special Health Issues  
Food and Drug Administration  
10903 New Hampshire Ave  
Building 32, Mail Stop 5353  
Silver Spring, MD 20993

### **REPORTING REQUIREMENTS**

We remind you that you must comply with reporting requirements for an approved NDA (21 CFR 314.80 and 314.81).

If you have any questions, call Heather Buck, Regulatory Project Manager, at (301) 796-1413.

Sincerely,

*{See appended electronic signature page}*

Joyce Korvick, M.D., M.P.H.  
Deputy Director for Safety  
Division of Gastroenterology Products  
Office of Drug Evaluation III  
Center for Drug Evaluation and Research

ENCLOSURE: Content of Labeling

-----  
**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**  
-----

/s/  
-----

JOYCE A KORVICK  
11/18/2010

**CENTER FOR DRUG EVALUATION AND  
RESEARCH**

***APPLICATION NUMBER:***

**017862Orig1s063**

**LABELING**

# REGLAN Injection (metoclopramide injection, USP)

R<sub>x</sub> only

## WARNING: TARDIVE DYSKINESIA


Treatment with metoclopramide can cause tardive dyskinesia, a serious movement disorder that is often irreversible. The risk of developing tardive dyskinesia increases with duration of treatment and total cumulative dose.

Metoclopramide therapy should be discontinued in patients who develop signs or symptoms of tardive dyskinesia. There is no known treatment for tardive dyskinesia. In some patients, symptoms may lessen or resolve after metoclopramide treatment is stopped.

Treatment with metoclopramide for longer than 12 weeks should be avoided in all but rare cases where therapeutic benefit is thought to outweigh the risk of developing tardive dyskinesia. See WARNINGS.

## DESCRIPTION

Metoclopramide hydrochloride is a white crystalline, odorless substance, freely soluble in water. Chemically, it is 4-amino-5-chloro-N-[2-(diethylamino)ethyl]-2-methoxy benzamide monohydrochloride monohydrate. Molecular weight: 354.3.



REGLAN Injection (metoclopramide injection, USP) is a clear, colorless, sterile solution with a pH of 4.5-6.5 for intravenous (IV) or intramuscular (IM) administration.

This product is light sensitive. It should be inspected before use and discarded if either color or particulate is observed.

2 mL single dose vials; 10 mL and 30 mL single dose vials

Each 1 mL contains: Metoclopramide base 5 mg (as the monohydrochloride monohydrate), Sodium Chloride, USP 8.5 mg, Water for Injection, USP q.s. pH adjusted, when necessary, with hydrochloric acid and/or sodium hydroxide.

## CLINICAL PHARMACOLOGY

Metoclopramide stimulates motility of the upper gastrointestinal tract without stimulating gastric, biliary, or pancreatic secretions. Its mode of action is unclear. It seems to sensitize

tissues to the action of acetylcholine. The effect of metoclopramide on motility is not dependent on intact vagal innervation, but it can be abolished by anticholinergic drugs.

Metoclopramide increases the tone and amplitude of gastric (especially antral) contractions, relaxes the pyloric sphincter and the duodenal bulb, and increases peristalsis of the duodenum and jejunum resulting in accelerated gastric emptying and intestinal transit. It increases the resting tone of the lower esophageal sphincter. It has little, if any, effect on the motility of the colon or gallbladder.

In patients with gastroesophageal reflux and low LESP (lower esophageal sphincter pressure), single oral doses of metoclopramide produce dose-related increases in LESP. Effects begin at about 5 mg and increase through 20 mg (the largest dose tested). The increase in LESP from a 5 mg dose lasts about 45 minutes and that of 20 mg lasts between 2 and 3 hours. Increased rate of stomach emptying has been observed with single oral doses of 10 mg.

The antiemetic properties of metoclopramide appear to be a result of its antagonism of central and peripheral dopamine receptors. Dopamine produces nausea and vomiting by stimulation of the medullary chemoreceptor trigger zone (CTZ), and metoclopramide blocks stimulation of the CTZ by agents like l-dopa or apomorphine which are known to increase dopamine levels or to possess dopamine-like effects. Metoclopramide also abolishes the slowing of gastric emptying caused by apomorphine.

Like the phenothiazines and related drugs, which are also dopamine antagonists, metoclopramide produces sedation and may produce extrapyramidal reactions, although these are comparatively rare (see **WARNINGS**). Metoclopramide inhibits the central and peripheral effects of apomorphine, induces release of prolactin and causes a transient increase in circulating aldosterone levels, which may be associated with transient fluid retention.

The onset of pharmacological action of metoclopramide is 1 to 3 minutes following an intravenous dose, 10 to 15 minutes following intramuscular administration, and 30 to 60 minutes following an oral dose; pharmacological effects persist for 1 to 2 hours.

### **Pharmacokinetics**

Metoclopramide is rapidly and well absorbed. Relative to an intravenous dose of 20 mg, the absolute oral bioavailability of metoclopramide is  $80\% \pm 15.5\%$  as demonstrated in a crossover study of 18 subjects. Peak plasma concentrations occur at about 1-2 hr after a single oral dose. Similar time to peak is observed after individual doses at steady state.

In a single dose study of 12 subjects, the area under the drug concentration-time curve increases linearly with doses from 20 to 100 mg. Peak concentrations increase linearly with dose; time to peak concentrations remains the same; whole body clearance is unchanged; and the elimination rate remains the same. The average elimination half-life in individuals with normal renal function is 5-6 hr. Linear kinetic processes adequately describe the absorption and elimination of metoclopramide.

Approximately 85% of the radioactivity of an orally administered dose appears in the urine within 72 hr. Of the 85% eliminated in the urine, about half is present as free or conjugated metoclopramide.

The drug is not extensively bound to plasma proteins (about 30%). The whole body volume of distribution is high (about 3.5 L/kg) which suggests extensive distribution of drug to the tissues.

Renal impairment affects the clearance of metoclopramide. In a study with patients with varying degrees of renal impairment, a reduction in creatinine clearance was correlated with a reduction in plasma clearance, renal clearance, non-renal clearance, and increase in elimination half-life. The kinetics of metoclopramide in the presence of renal impairment remained linear however. The reduction in clearance as a result of renal impairment suggests that adjustment downward of maintenance dosage should be done to avoid drug accumulation.

#### Adult Pharmacokinetic Data

| Parameter              | Value     |
|------------------------|-----------|
| Vd (L/kg)              | ~ 3.5     |
| Plasma Protein Binding | ~ 30%     |
| t <sub>1/2</sub> (hr)  | 5-6       |
| Oral Bioavailability   | 80%±15.5% |

In pediatric patients, the pharmacodynamics of metoclopramide following oral and intravenous administration are highly variable and a concentration-effect relationship has not been established.

There are insufficient reliable data to conclude whether the pharmacokinetics of metoclopramide in adults and the pediatric population are similar. Although there are insufficient data to support the efficacy of metoclopramide in pediatric patients with symptomatic gastroesophageal reflux (GER) or cancer chemotherapy-related nausea and vomiting, its pharmacokinetics have been studied in these patient populations.

In an open-label study, six pediatric patients (age range, 3.5 weeks to 5.4 months) with GER received a metoclopramide 0.15 mg/kg oral solution every 6 hours for 10 doses. The mean peak plasma concentration of metoclopramide after the tenth dose was 2-fold (56.8 µg/L) higher compared to that observed after the first dose (29 µg/L) indicating drug accumulation with repeated dosing. After the tenth dose, the mean time to reach peak concentrations (2.2 hr), half-life (4.1 hr), clearance (0.67 L/h/kg), and volume of distribution (4.4 L/kg) of metoclopramide were similar to those observed after the first dose. In the youngest patient (age, 3.5 weeks), metoclopramide half-life after the first and the tenth dose (23.1 and 10.3 hr, respectively) was significantly longer compared to other infants due to reduced clearance. This may be attributed to immature hepatic and renal systems at birth.

Single intravenous doses of metoclopramide 0.22 to 0.46 mg/kg (mean, 0.35 mg/kg) were administered over 5 minutes to 9 pediatric cancer patients receiving chemotherapy (mean age, 11.7 years; range, 7 to 14 yr) for prophylaxis of cytotoxic-induced vomiting. The metoclopramide plasma concentrations extrapolated to time zero ranged from 65 to 395 µg/L (mean, 152 µg/L). The mean elimination half-life, clearance, and volume of distribution of metoclopramide were 4.4 hr (range, 1.7 to 8.3 hr), 0.56 L/h/kg (range, 0.12 to 1.20 L/h/kg), and 3.0 L/kg (range, 1.0 to 4.8 L/kg), respectively.

In another study, nine pediatric cancer patients (age range, 1 to 9 yr) received 4 to 5 intravenous infusions (over 30 minutes) of metoclopramide at a dose of 2 mg/kg to control emesis. After the last dose, the peak serum concentrations of metoclopramide ranged from 1060 to 5680 µg/L. The mean elimination half-life, clearance, and volume of distribution of metoclopramide were 4.5 hr (range, 2.0 to 12.5 hr), 0.37 L/h/kg (range, 0.10 to 1.24 L/h/kg), and 1.93 L/kg (range, 0.95 to 5.50 L/kg), respectively.

### Pediatric Pharmacokinetic Studies

| Reference | Dose, Route                                                       | t <sub>1/2</sub><br>(hr) | Cl<br>(L/hr/kg)   | Vd<br>(L/kg)<br>(Dose/Cp0) | C <sub>max</sub><br>(µg/L) |
|-----------|-------------------------------------------------------------------|--------------------------|-------------------|----------------------------|----------------------------|
| 1.        | 0.35 mg/kg,<br>IV over 5 min                                      | 4.4±0.56                 | 0.56±0.10         | 3.0±0.38                   | 152±31                     |
| 2.        | 2 mg/kg<br>30 min IV<br>infusion 4-5<br>times<br>within 9.5 hours | 4.5 <sup>a</sup>         | 0.37 <sup>a</sup> | 1.93 <sup>a</sup>          | 1060 to 5680 <sup>a</sup>  |

<sup>a</sup>. SEM not available.

1. Bateman, DN, et al. *Br J Clin Pharmac* 15:557-559, 1983.

2. Ford, C. *Clin Pharmac Ther* 43:196, 1988.

### INDICATIONS AND USAGE

#### Diabetic Gastroparesis (Diabetic Gastric Stasis)

REGLAN Injection (metoclopramide hydrochloride, USP) is indicated for the relief of symptoms associated with acute and recurrent diabetic gastric stasis.

#### The Prevention of Nausea and Vomiting Associated with Emetogenic Cancer Chemotherapy

REGLAN Injection is indicated for the prophylaxis of vomiting associated with emetogenic cancer chemotherapy.

#### The Prevention of Postoperative Nausea and Vomiting

REGLAN Injection is indicated for the prophylaxis of postoperative nausea and vomiting in those circumstances where nasogastric suction is undesirable.

#### Small Bowel Intubation

REGLAN Injection may be used to facilitate small bowel intubation in adults and pediatric patients in whom the tube does not pass the pylorus with conventional maneuvers.

#### Radiological Examination

REGLAN Injection may be used to stimulate gastric emptying and intestinal transit of barium in cases where delayed emptying interferes with radiological examination of the stomach and/or small intestine.

### CONTRAINDICATIONS

Metoclopramide should not be used whenever stimulation of gastrointestinal motility might be dangerous, e.g., in the presence of gastrointestinal hemorrhage, mechanical obstruction, or perforation.

Metoclopramide is contraindicated in patients with pheochromocytoma because the drug may cause a hypertensive crisis, probably due to release of catecholamines from the tumor. Such hypertensive crises may be controlled by phentolamine.

Metoclopramide is contraindicated in patients with known sensitivity or intolerance to the drug.

Metoclopramide should not be used in epileptics or patients receiving other drugs which are likely to cause extrapyramidal reactions, since the frequency and severity of seizures or extrapyramidal reactions may be increased.

## **WARNINGS**

### **Neuroleptic Malignant Syndrome (NMS)**

There have been rare reports of an uncommon but potentially fatal symptom complex sometimes referred to as Neuroleptic Malignant Syndrome (NMS) associated with metoclopramide. Clinical manifestations of NMS include hyperthermia, muscle rigidity, altered consciousness, and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis and cardiac arrhythmias).

The diagnostic evaluation of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to identify cases where the clinical presentation includes both serious medical illness (e.g., pneumonia, systemic infection, etc.) and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, malignant hyperthermia, drug fever and primary central nervous system (CNS) pathology.

The management of NMS should include 1) immediate discontinuation of metoclopramide and other drugs not essential to concurrent therapy, 2) intensive symptomatic treatment and medical monitoring, and 3) treatment of any concomitant serious medical problems for which specific treatments are available. Bromocriptine and dantrolene sodium have been used in treatment of NMS, but their effectiveness have not been established (see **ADVERSE REACTIONS**).

### **Extrapyramidal Symptoms (EPS)**

#### ***Acute Dystonic Reactions***

Acute dystonic reactions occur in approximately 1 in 500 patients treated with the usual adult dosages of 30-40 mg/day of metoclopramide. These usually are seen during the first 24-48 hours of treatment with metoclopramide, occur more frequently in pediatric patients and adult patients less than 30 years of age and are even more frequent at the higher doses used in prophylaxis of vomiting due to cancer chemotherapy. These symptoms may include involuntary movements of limbs and facial grimacing, torticollis, oculogyric crisis, rhythmic protrusion of tongue, bulbar type of speech, trismus, or dystonic reactions resembling tetanus. Rarely, dystonic reactions may present as stridor and dyspnea, possibly due to laryngospasm. If these symptoms should occur, inject 50 mg Benadryl® (diphenhydramine hydrochloride) intramuscularly, and they usually will subside. Cogentin® (benztropine mesylate), 1 to 2 mg intramuscularly, may also be used to reverse these reactions.

#### ***Tardive Dyskinesia (see Boxed Warnings)***

Treatment with metoclopramide can cause tardive dyskinesia (TD), a potentially irreversible and disfiguring disorder characterized by involuntary movements of the face, tongue, or extremities. The risk of developing tardive dyskinesia increases with the duration of treatment and the total cumulative dose. An analysis of utilization patterns showed that about 20% of patients who used metoclopramide took it for longer than 12 weeks. Treatment with metoclopramide for longer than the recommended 12 weeks should be avoided in all but rare cases where therapeutic benefit is thought to outweigh the risk of developing TD.

Although the risk of developing TD in the general population may be increased among the elderly, women, and diabetics, it is not possible to predict which patients will develop metoclopramide-induced TD. Both the risk of developing TD and the likelihood that TD will become irreversible increase with duration of treatment and total cumulative dose.

Metoclopramide should be discontinued in patients who develop signs or symptoms of TD. There is no known effective treatment for established cases of TD, although in some patients, TD may remit, partially or completely, within several weeks to months after metoclopramide is withdrawn.

Metoclopramide itself may suppress, or partially suppress, the signs of TD, thereby masking the underlying disease process. The effect of this symptomatic suppression upon the long-term course of TD is unknown. Therefore, metoclopramide should not be used for the symptomatic control of TD.

### ***Parkinsonian-like Symptoms***

Parkinsonian-like symptoms, including bradykinesia, tremor, cogwheel rigidity, or mask-like facies, have occurred more commonly within the first 6 months after beginning treatment with metoclopramide, but occasionally after longer periods. These symptoms generally subside within 2-3 months following discontinuance of metoclopramide. Patients with preexisting Parkinson's disease should be given metoclopramide cautiously, if at all, since such patients may experience exacerbation of parkinsonian symptoms when taking metoclopramide.

### **Depression**

Mental depression has occurred in patients with and without prior history of depression. Symptoms have ranged from mild to severe and have included suicidal ideation and suicide. Metoclopramide should be given to patients with a prior history of depression only if the expected benefits outweigh the potential risks.

## **PRECAUTIONS**

### **General**

In one study in hypertensive patients, intravenously administered metoclopramide was shown to release catecholamines; hence, caution should be exercised when metoclopramide is used in patients with hypertension.

Intravenous injections of undiluted metoclopramide should be made slowly allowing 1 to 2 minutes for 10 mg since a transient but intense feeling of anxiety and restlessness, followed by drowsiness, may occur with rapid administration.

Because metoclopramide produces a transient increase in plasma aldosterone, certain patients, especially those with cirrhosis or congestive heart failure, may be at risk of developing fluid retention and volume overload. If these side effects occur at any time during metoclopramide therapy, the drug should be discontinued.

Intravenous administration of REGLAN Injection diluted in a parenteral solution should be made slowly over a period of not less than 15 minutes.

Giving a promotility drug such as metoclopramide theoretically could put increased pressure on suture lines following a gut anastomosis or closure. This possibility should be considered and weighed when deciding whether to use metoclopramide or nasogastric suction in the prevention of postoperative nausea and vomiting.

### **Information for Patients**

A patient Medication Guide is available for REGLAN Injection. The prescriber or health professional should instruct patients, their families, and their caregivers to read the Medication Guide and should assist them in understanding its contents. Patients should be given the

opportunity to discuss the contents of the Medication Guide and to obtain answers to any questions they may have. Refer to accompanying Medication Guide.

Metoclopramide may impair the mental and/or physical abilities required for the performance of hazardous tasks such as operating machinery or driving a motor vehicle. The ambulatory patient should be cautioned accordingly.

### **Drug Interactions**

The effects of metoclopramide on gastrointestinal motility are antagonized by anticholinergic drugs and narcotic analgesics. Additive sedative effects can occur when metoclopramide is given with alcohol, sedatives, hypnotics, narcotics, or tranquilizers.

The finding that metoclopramide releases catecholamines in patients with essential hypertension suggests that it should be used cautiously, if at all, in patients receiving monoamine oxidase inhibitors.

Absorption of drugs from the stomach may be diminished (e.g., digoxin) by metoclopramide, whereas the rate and/or extent of absorption of drugs from the small bowel may be increased (e.g., acetaminophen, tetracycline, levodopa, ethanol, cyclosporine).

Gastroparesis (gastric stasis) may be responsible for poor diabetic control in some patients. Exogenously administered insulin may begin to act before food has left the stomach and lead to hypoglycemia. Because the action of metoclopramide will influence the delivery of food to the intestines and thus the rate of absorption, insulin dosage or timing of dosage may require adjustment.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

A 77-week study was conducted in rats with oral doses up to about 40 times the maximum recommended human daily dose. Metoclopramide elevates prolactin levels and the elevation persists during chronic administration. Tissue culture experiments indicate that approximately one-third of human breast cancers are prolactin-dependent *in vitro*, a factor of potential importance if the prescription of metoclopramide is contemplated in a patient with previously detected breast cancer. Although disturbances such as galactorrhea, amenorrhea, gynecomastia, and impotence have been reported with prolactin-elevating drugs, the clinical significance of elevated serum prolactin levels is unknown for most patients. An increase in mammary neoplasms has been found in rodents after chronic administration of prolactin-stimulating neuroleptic drugs and metoclopramide. Neither clinical studies nor epidemiologic studies conducted to date, however, have shown an association between chronic administration of these drugs and mammary tumorigenesis; the available evidence is too limited to be conclusive at this time.

An Ames mutagenicity test performed on metoclopramide was negative.

### **Pregnancy Category B**

Reproduction studies performed in rats, mice and rabbits by the IM, IV, subcutaneous (SC), and oral routes at maximum levels ranging from 12 to 250 times the human dose have demonstrated no impairment of fertility or significant harm to the fetus due to metoclopramide. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

### **Nursing Mothers**

Metoclopramide is excreted in human milk. Caution should be exercised when metoclopramide is administered to a nursing mother.

### **Pediatric Use**

Safety and effectiveness in pediatric patients have not been established except as stated to facilitate small bowel intubation (see **OVERDOSAGE** and **DOSAGE AND ADMINISTRATION**).

Care should be exercised in administering metoclopramide to neonates since prolonged clearance may produce excessive serum concentrations (see **CLINICAL PHARMACOLOGY — Pharmacokinetics**). In addition, neonates have reduced levels of NADH-cytochrome b<sub>5</sub> reductase which, in combination with the aforementioned pharmacokinetic factors, make neonates more susceptible to methemoglobinemia (see **OVERDOSAGE**).

The safety profile of metoclopramide in adults cannot be extrapolated to pediatric patients. Dystonias and other extrapyramidal reactions associated with metoclopramide are more common in the pediatric population than in adults. (See **WARNINGS** and **ADVERSE REACTIONS — Extrapyramidal Reactions**.)

### **Geriatric Use**

Clinical studies of REGLAN Injection did not include sufficient numbers of subjects aged 65 and over to determine whether elderly subjects respond differently from younger subjects.

The risk of developing parkinsonian-like side effects increases with ascending dose. Geriatric patients should receive the lowest dose of REGLAN Injection that is effective. If parkinsonian-like symptoms develop in a geriatric patient receiving REGLAN Injection, REGLAN Injection should generally be discontinued before initiating any specific anti-parkinsonian agents (see **WARNINGS**).

The elderly may be at greater risk for tardive dyskinesia (see **WARNINGS – Tardive Dyskinesia**).

Sedation has been reported in REGLAN Injection users. Sedation may cause confusion and manifest as over-sedation in elderly (see **CLINICAL PHARMACOLOGY, PRECAUTIONS – Information for Patients** and **ADVERSE REACTIONS – CNS Effects**).

REGLAN Injection is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function (see **DOSAGE AND ADMINISTRATION - Use in Patients With Renal or Hepatic Impairment**).

For these reasons, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased renal function, concomitant disease, or other drug therapy in the elderly (see **Use in Patients With Renal or Hepatic Impairment**).

### **Other Special Populations**

Patients with NADH-cytochrome b<sub>5</sub> reductase deficiency are at an increased risk of developing methemoglobinemia and/or sulfhemoglobinemia when metoclopramide is administered. In patients with G6PD deficiency who experience metoclopramide-induced methemoglobinemia, methylene blue treatment is not recommended (see **OVERDOSAGE**).

## **ADVERSE REACTIONS**

In general, the incidence of adverse reactions correlates with the dose and duration of metoclopramide administration. The following reactions have been reported, although in most instances, data do not permit an estimate of frequency:

### **CNS Effects**

Restlessness, drowsiness, fatigue, and lassitude may occur in patients receiving the recommended prescribed dosage of REGLAN Injection. Insomnia, headache, confusion, dizziness, or mental depression with suicidal ideation also may occur (see **WARNINGS**). In cancer chemotherapy patients being treated with 1-2 mg/kg per dose, incidence of drowsiness is about 70%. There are isolated reports of convulsive seizures without clear-cut relationship to metoclopramide. Rarely, hallucinations have been reported.

### **Extrapyramidal Reactions (EPS)**

Acute dystonic reactions, the most common type of EPS associated with metoclopramide, occur in approximately 0.2% of patients (1 in 500) treated with 30 to 40 mg of metoclopramide per day. In cancer chemotherapy patients receiving 1-2 mg/kg per dose, the incidence is 2% in patients over the ages of 30-35, and 25% or higher in pediatric patients and adult patients less than 30 years of age who have not had prophylactic administration of diphenhydramine. Symptoms include involuntary movements of limbs, facial grimacing, torticollis, oculogyric crisis, rhythmic protrusion of tongue, bulbar type of speech, trismus, opisthotonus (tetanus-like reactions), and, rarely, stridor and dyspnea possibly due to laryngospasm; ordinarily these symptoms are readily reversed by diphenhydramine (see **WARNINGS**).

Parkinsonian-like symptoms may include bradykinesia, tremor, cogwheel rigidity, mask-like facies (see **WARNINGS**).

Tardive dyskinesia most frequently is characterized by involuntary movements of the tongue, face, mouth, or jaw, and sometimes by involuntary movements of the trunk and/or extremities; movements may be choreoathetotic in appearance (see **WARNINGS**).

Motor restlessness (akathisia) may consist of feelings of anxiety, agitation, jitteriness, and insomnia, as well as inability to sit still, pacing, foot tapping. These symptoms may disappear spontaneously or respond to a reduction in dosage.

### **Neuroleptic Malignant Syndrome**

Rare occurrences of neuroleptic malignant syndrome (NMS) have been reported. This potentially fatal syndrome is comprised of the symptom complex of hyperthermia, muscular rigidity, altered consciousness, and autonomic instability (see **WARNINGS**).

### **Endocrine Disturbances**

Galactorrhea, amenorrhea, gynecomastia, impotence secondary to hyperprolactinemia (see **PRECAUTIONS**). Fluid retention secondary to transient elevation of aldosterone (see **CLINICAL PHARMACOLOGY**).

### **Cardiovascular**

Hypotension, hypertension, supraventricular tachycardia, bradycardia, fluid retention, acute congestive heart failure and possible atrioventricular (AV) block (see **CONTRAINDICATIONS** and **PRECAUTIONS**).

### **Gastrointestinal**

Nausea and bowel disturbances, primarily diarrhea.

## **Hepatic**

Rarely, cases of hepatotoxicity, characterized by such findings as jaundice and altered liver function tests, when metoclopramide was administered with other drugs with known hepatotoxic potential.

## **Renal**

Urinary frequency and incontinence.

## **Hematologic**

A few cases of neutropenia, leukopenia, or agranulocytosis, generally without clear-cut relationship to metoclopramide. Methemoglobinemia in adults and especially with overdosage in neonates (see **OVERDOSAGE**). Sulfhemoglobinemia in adults.

## **Allergic Reactions**

A few cases of rash, urticaria, or bronchospasm, especially in patients with a history of asthma. Rarely, angioneurotic edema, including glossal or laryngeal edema.

## **Miscellaneous**

Visual disturbances. Porphyria.

Transient flushing of the face and upper body, without alterations in vital signs, following high doses intravenously.

## **OVERDOSAGE**

Symptoms of overdosage may include drowsiness, disorientation and extrapyramidal reactions. Anticholinergic or antiparkinson drugs or antihistamines with anticholinergic properties may be helpful in controlling the extrapyramidal reactions. Symptoms are self-limiting and usually disappear within 24 hours.

Hemodialysis removes relatively little metoclopramide, probably because of the small amount of the drug in blood relative to tissues. Similarly, continuous ambulatory peritoneal dialysis does not remove significant amounts of drug. It is unlikely that dosage would need to be adjusted to compensate for losses through dialysis. Dialysis is not likely to be an effective method of drug removal in overdose situations.

Unintentional overdose due to misadministration has been reported in infants and children with the use of REGLAN syrup. While there was no consistent pattern to the reports associated with these overdoses, events included seizures, extrapyramidal reactions, and lethargy.

Methemoglobinemia has occurred in premature and full-term neonates who were given overdoses of metoclopramide (1-4 mg/kg/day orally, intramuscularly or intravenously for 1-3 or more days). Methemoglobinemia can be reversed by the intravenous administration of methylene blue. However, methylene blue may cause hemolytic anemia in patients with G6PD deficiency, which may be fatal (see **PRECAUTIONS – Other Special Populations**).

## **DOSAGE AND ADMINISTRATION**

### **For the Relief of Symptoms Associated with Diabetic Gastroparesis (Diabetic Gastric Stasis)**

If only the earliest manifestations of diabetic gastric stasis are present, oral administration of metoclopramide may be initiated. However, if severe symptoms are present, therapy should begin with REGLAN Injection (IM or IV). Doses of 10 mg may be administered slowly by the intravenous route over a 1- to 2-minute period.

Administration of REGLAN Injection (metoclopramide injection, USP) up to 10 days may be required before symptoms subside, at which time oral administration of metoclopramide may be instituted. The physician should make a thorough assessment of the risks and benefits prior to prescribing further metoclopramide treatment.

### **For the Prevention of Nausea and Vomiting Associated with Emetogenic Cancer Chemotherapy**

Intravenous infusions should be made slowly over a period of not less than 15 minutes, 30 minutes before beginning cancer chemotherapy and repeated every 2 hours for two doses, then every 3 hours for three doses.

The initial two doses should be 2 mg/kg if highly emetogenic drugs such as cisplatin or dacarbazine are used alone or in combination. For less emetogenic regimens, 1 mg/kg per dose may be adequate.

For doses in excess of 10 mg, REGLAN Injection should be diluted in 50 mL of a parenteral solution.

The preferred parenteral solution is Sodium Chloride Injection (normal saline), which when combined with REGLAN Injection, can be stored frozen for up to 4 weeks. REGLAN Injection is degraded when admixed and frozen with Dextrose-5% in Water. REGLAN Injection diluted in Sodium Chloride Injection, Dextrose-5% in Water, Dextrose-5% in 0.45% Sodium Chloride, Ringer's Injection, or Lactated Ringer's Injection may be stored up to 48 hours (without freezing) after preparation if protected from light. All dilutions may be stored unprotected from light under normal light conditions up to 24 hours after preparation.

If acute dystonic reactions should occur, inject 50 mg Benadryl® (diphenhydramine hydrochloride) intramuscularly, and the symptoms usually will subside.

### **For the Prevention of Postoperative Nausea and Vomiting**

REGLAN Injection should be given intramuscularly near the end of surgery. The usual adult dose is 10 mg; however, doses of 20 mg may be used.

### **To Facilitate Small Bowel Intubation**

If the tube has not passed the pylorus with conventional maneuvers in 10 minutes, a single dose (undiluted) may be administered slowly by the intravenous route over a 1- to 2-minute period.

The recommended single dose is: Pediatric patients above 14 years of age and adults — 10 mg metoclopramide base. Pediatric patients (6-14 years of age) — 2.5 to 5 mg metoclopramide base; (under 6 years of age) — 0.1 mg/kg metoclopramide base.

### **To Aid in Radiological Examinations**

In patients where delayed gastric emptying interferes with radiological examination of the stomach and/or small intestine, a single dose may be administered slowly by the intravenous route over a 1- to 2-minute period.

For dosage, see intubation above.

### **Use in Patients With Renal or Hepatic Impairment**

Since metoclopramide is excreted principally through the kidneys, in those patients whose creatinine clearance is below 40 mL/min, therapy should be initiated at approximately one-half the recommended dosage. Depending upon clinical efficacy and safety considerations, the dosage may be increased or decreased as appropriate.

See **OVERDOSAGE** section for information regarding dialysis.

Metoclopramide undergoes minimal hepatic metabolism, except for simple conjugation. Its safe use has been described in patients with advanced liver disease whose renal function was normal.

**NOTE:** Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

### **ADMIXTURES COMPATIBILITIES**

REGLAN Injection (metoclopramide injection, USP) is compatible for mixing and injection with the following dosage forms to the extent indicated below:

#### **Physically and Chemically Compatible Up to 48 Hours**

Cimetidine Hydrochloride (SK&F), Mannitol, USP (Abbott), Potassium Acetate, USP (Invenex), Potassium Phosphate, USP (Invenex).

#### **Physically Compatible Up to 48 Hours**

Ascorbic Acid, USP (Abbott), Benztropine Mesylate, USP (MS&D), Cytarabine, USP (Upjohn), Dexamethasone Sodium Phosphate, USP (ESI, MS&D), Diphenhydramine Hydrochloride, USP (Parke-Davis), Doxorubicin Hydrochloride, USP (Adria), Heparin Sodium, USP (ESI), Hydrocortisone Sodium Phosphate (MS&D), Lidocaine Hydrochloride, USP (ESI), Multi-Vitamin Infusion (must be refrigerated-USV), Vitamin B Complex with Ascorbic Acid (Roche).

#### **Physically Compatible Up to 24 Hours**

**(Do not use if precipitation occurs)**

Clindamycin Phosphate, USP (Upjohn), Cyclophosphamide, USP (Mead-Johnson), Insulin, USP (Lilly).

#### **Conditionally Compatible**

**(Use within one hour after mixing or may be infused directly into the same running IV line)**

Ampicillin Sodium, USP (Bristol), Cisplatin (Bristol), Erythromycin Lactobionate, USP (Abbott), Methotrexate Sodium, USP (Lederle), Penicillin G Potassium, USP (Squibb), Tetracycline Hydrochloride, USP (Lederle).

#### **Incompatible**

**(Do Not Mix)**

Cephalothin Sodium, USP (Lilly), Chloramphenicol Sodium, USP (Parke-Davis), Sodium Bicarbonate, USP (Abbott).

### **HOW SUPPLIED**

REGLAN Injection (metoclopramide injection, USP) 5 mg metoclopramide base (as the monohydrochloride monohydrate) per mL; available in:

2 mL single dose vials in cartons of 25 (NDC 60977-451-01),

10 mL single dose vials in cartons of 25 (NDC 60977-451-02),

30 mL single dose vials in cartons of 25 (NDC 60977-451-03).

| <b>Container</b>      | <b>Total Contents #</b> | <b>Concentration #</b> | <b>Administration</b>       |
|-----------------------|-------------------------|------------------------|-----------------------------|
| 2 mL single dose vial | 10 mg                   | 5 mg/mL                | FOR IV or IM ADMINISTRATION |

|                           |        |         |                                              |
|---------------------------|--------|---------|----------------------------------------------|
| 10 mL single<br>dose vial | 50 mg  | 5 mg/mL | FOR IV INFUSION ONLY;<br>DILUTE BEFORE USING |
| 30 mL single<br>dose vial | 150 mg | 5 mg/mL | FOR IV INFUSION ONLY;<br>DILUTE BEFORE USING |

---

# Metoclopramide base (as the monohydrochloride monohydrate)

---

Store vials in carton until used. Do not store open single dose vials for later use, as they contain no preservative.

This product is light sensitive. It should be inspected before use and discarded if either color or particulate is observed.

**Dilutions may be stored unprotected from light under normal light conditions up to 24 hours after preparation.**

**REGLAN Injection should be stored at Controlled Room Temperature, 20°-25°C (68°-77°F) [see USP Controlled Room Temperature].**

Reglan is a trademark of Alaven Pharmaceutical, LLC, its agents or assignees.

ESI logo is a trademark of Baxter International Inc.



Manufactured by

**Baxter Healthcare Corporation**

Deerfield, IL 60015 USA

For Product Inquiry 1 800 933 0303

Revised November 2010

MLT00066,I

**CENTER FOR DRUG EVALUATION AND  
RESEARCH**

*APPLICATION NUMBER:*

**017862Orig1s063**

**OTHER REVIEW(S)**

# REGULATORY PROJECT MANAGER LABELING REVIEW

## Division of Gastroenterology Products

| <u>NDA</u> | <u>SLR</u> | <u>Drug Name</u>                                                                                | <u>Applicant</u> | <u>Submission</u> | <u>Receipt*</u> |
|------------|------------|-------------------------------------------------------------------------------------------------|------------------|-------------------|-----------------|
| 17-862     | S-063      | Reglan Injection (metoclopramide injection, USP), 5 mg/mL, 2 mL Vial, 10 mL Vial and 30 mL Vial | Baxter           | 9/24/2010         | 9/27/2010       |
| 17-854     | S-055      | Reglan (metoclopramide) tablets                                                                 | Alaven           | 9/30/2010         | 10/1/2010       |
| 21-793     | S-008      | Reglan ODT (metoclopramide) orally disintegrating tablets                                       | Alaven           | 9/30/2010         | 10/1/2010       |
| 22-246     | S-001      | Metozolv ODT (metoclopramide hydrochloride), 5 mg & 10 mg                                       | Salix            | 8/6/2010          | 8/9/2010        |

\*SPL was only received for 22-246/S-001 on 8/9/10.

### **Background and Summary**

The above supplements were submitted in response to prior approval supplement (PAS) request letters that the Division issued on July 20, 2010. The letters made the following request:

*We have reassessed the statements regarding Tardive Dyskinesia in the Warnings and Precautions Section of the prescribing information. Inadvertent editing has resulted in inaccuracies regarding the representation of the data reviewed.*

*We request that the following changes in the labeling be made so as to furnish adequate information for the safe and effective use of the drug:*

#### ***Tardive Dyskinesia (see Boxed Warnings)***

*Treatment with metoclopramide can cause tardive dyskinesia (TD), a potentially irreversible and disfiguring disorder characterized by involuntary movements of the face, tongue, or extremities. The risk of developing tardive dyskinesia increases with the duration of treatment and the total cumulative dose. An analysis of utilization patterns showed that about 20% of patients who used metoclopramide took it for longer than 12 weeks. Treatment with metoclopramide for longer than the recommended 12 weeks should be avoided in all but rare cases where therapeutic benefit is thought to outweigh the risk of developing TD.*

*Although the risk of developing TD in the general population may be increased among the elderly, women, and diabetics, it is not possible to predict which patients will develop metoclopramide-induced TD. Both the risk of developing TD and the likelihood that TD will become irreversible increase with duration of treatment and total cumulative dose.*

*Metoclopramide should be discontinued in patients who develop signs or symptoms of TD.*

*There is no known effective treatment for established cases of TD, although in some patients, TD may remit, partially or completely, within several weeks to months after metoclopramide is withdrawn.*

*Metoclopramide itself may suppress, or partially suppress, the signs of TD, thereby masking the underlying disease process. The effect of this symptomatic suppression upon the long-term course of TD is unknown. Therefore, metoclopramide should not be used for the symptomatic control of TD.*

It should be noted that SPL was only received for 22-246/S-001 on 8/9/10. We waived the requirement to submit SPL since the supplement is for a correction that we requested. The sponsors will be reminded to submit final SPL within 14 days upon receipt of the approval letter.

## **Review**

This review focuses on the proposed Word version of the submitted package inserts as compared to those that were last approved. This review also verifies that the newly proposed content matches the language that was requested. Deletions are marked with ~~strikeout~~. Additions are marked with underline.

### **17-862/S-063 Reglan Injection**

#### **Proposed Word versus last approved (6/30/09)**

1. "Injection" was added to the drug name in various places including once under Indications and Usage, and several times under Geriatric Use.
2. The reference to the Medication Guide was reworded under Information for Patients: ~~The complete text of the~~ Refer to accompanying Medication Guide is printed at the end of this document.
3. Minor revisions were made to the trademark and manufacturing information at the end of the label.
4. The newly proposed language under **Warnings, Tardive Dyskinesia** is identical to the language requested by FDA.

**Comments** – The additional changes made are improvements to the label and are acceptable.

### **17-854/S-055 Reglan Tablets**

#### **Proposed Word versus last approved (9/4/09)**

1. Only minor stylistic differences were found.
2. The newly proposed language under **Warnings, Tardive Dyskinesia** is identical to the language requested by FDA.

**Comments** – The label is acceptable.

### **21-793/S-008 Reglan ODT**

#### **Proposed Word versus last approved (9/9/10)**

1. The only difference in the proposed label was (b) (4)
2. The newly proposed language under **Warnings, Tardive Dyskinesia** is identical to the language requested by FDA.

**Comments** – The label is acceptable.

## **22-246/S-001 Metozolv ODT**

### **Proposed Word versus last approved (9/4/09)**

1. The proposed Word version corrected the Patient Counseling Information statement in Highlights as follows: See 17 for PATIENT COUNSELING INFORMATION and the FDA-approved Medication Guide
2. The proposed Word version corrected the Boxed Warning heading in the Table of Contents as follows: WARNING: TARDIVE DYSKINESIA
3. The newly proposed language under **Warnings, Tardive Dyskinesia** is similar to that requested of the sponsor, with the exception of the word “increase” which should be “increases” in the second sentence. The following revision is recommended: “The risk of developing tardive dyskinesia increases with the duration of treatment and the total cumulative dose.”
5. Minor revisions were made including updates to the trademark information throughout the label.

**Comments** – The sponsor will be asked to make the revision as stated above (#3). The additional changes made to the label are improvements to the label and are acceptable.

### **Recommendations**

The sponsor for 22-246/S-001 Metozolv ODT will be asked to make the revision recommended above (see comments). All sponsors will be reminded in the approval letters, to submit final SPL identical to the approved labels. These labeling supplements are recommended for approval pending final review by the Director for Safety.

---

Heather Buck  
Regulatory Project Manager

Supervisory Comment/Concurrence:

---

Richard Ishihara  
Chief, Project Management Staff

HB: 10/20/10  
Revised/Initialed: RI 11/4/10  
Finalized: HB 11/8/10

-----  
**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**  
-----

/s/  
-----

HEATHER G BUCK  
11/09/2010

RICHARD W ISHIHARA  
11/09/2010