

# CENTER FOR DRUG EVALUATION AND RESEARCH

## Approval Package for:

### ***APPLICATION NUMBER:***

**NDA 019591/S-022**

***Trade Name:*** Lariam for oral use, 250mg

***Generic or Proper Name:*** Mefloquine Hydrochloride

***Sponsor:*** Hoffman-LaRoche, Inc.

***Approval Date:*** August 03, 2003

***Indication:*** Indicated for the treatment of mild to moderate acute malaria caused by mefloquine-susceptible strains of *P. falciparum* (both chloroquine-susceptible and resistant strains) or by *Plasmodium vivax*.

# CENTER FOR DRUG EVALUATION AND RESEARCH

NDA 019591/S-022

## CONTENTS

### Reviews / Information Included in this NDA Review.

<b>Approval Letter</b>	<b>X</b>
<b>Other Action Letters</b>	
<b>Labeling</b>	<b>X</b>
<b>REMS</b>	
<b>Summary Review</b>	
<b>Officer/Employee List</b>	
<b>Office Director Memo</b>	
<b>Cross Discipline Team Leader Review</b>	
<b>Clinical Review(s)</b>	
<b>Product Quality Review(s)</b>	
<b>Non-Clinical Review(s)</b>	
<b>Statistical Review(s)</b>	
<b>Clinical Microbiology / Virology Review(s)</b>	
<b>Clinical Pharmacology Review(s)</b>	
<b>Other Reviews</b>	<b>X</b>
<b>Risk Assessment and Risk Mitigation Review(s)</b>	
<b>Proprietary Name Review(s)</b>	
<b>Administrative/Correspondence Document(s)</b>	

**CENTER FOR DRUG EVALUATION AND  
RESEARCH**

*APPLICATION NUMBER:*

**NDA 019591/S-022**

**APPROVAL LETTER**



NDA 19-591/S-022

Hoffman-LaRoche, Inc.  
Attention: Lynn DeVenezia-Tobias  
Program Manager, Drug Regulatory Affairs  
340 Kingsland St.  
Nutley, NJ 07110-1199

Dear Ms. DeVenezia-Tobias:

Please refer to your supplemental new drug application dated July 23, 2003, received July 25, 2003, submitted under section 505(b) of the Federal Food, Drug, and Cosmetic Act for Lariam® (mefloquine hydrochloride) Tablets, 250 mg.

We acknowledge receipt of your submission dated August 15, 2003.

This “changes being effected” supplemental new drug application provides for following revisions to the labeling. Deleted text noted by ~~strike through~~ and additions are double underlined:

#### 1. PRECAUTIONS

- In the **Information for Patients** subsection, the following sentences now appear first:

Medication Guide: As required by law, a Lariam Medication Guide is supplied to patients when Lariam is dispensed. Patients should be instructed to read the MedGuide when Lariam is received. The complete text of the Medication Guide is reprinted at the end of this document.

2. The Lariam Medication Guide approved on July 9, 2003 (with the additional agreed upon proposed labeling changes discussed with you via e-mail on August 4-6, 2003 as follows) has been added to the end of the package insert:

**MEDICATION GUIDE**  
**LARIAM® (LAH-ree-am)**  
**(mefloquine hydrochloride) Tablets**  
**to Prevent Malaria**

**This Medication Guide is intended only for travelers who are taking Lariam to prevent malaria.** The information may not apply to patients who are sick with malaria and who are taking Lariam to treat malaria.

An information wallet card is provided at the end of this Medication Guide. Cut it out and carry it with you when you are taking Lariam.

This Medication Guide was \_\_\_\_\_ revised in August 2003. Please read it before you start taking Lariam and b)(4)----- refill. There may be new information. This Medication Guide does not take the place of talking with your prescriber (doctor or other health care provider) about Lariam and malaria prevention. Only you and your prescriber can decide if Lariam is right for you. If you cannot take Lariam, you may be able to take a different medicine to prevent malaria.

**What is the most important information I should know about Lariam?**

**1. Take Lariam exactly as prescribed to prevent malaria.**

Malaria is an infection that can cause death and is spread to humans through mosquito bites. If you travel to parts of the world where the mosquitoes carry the malaria parasite, you must take a malaria prevention medicine. Lariam is one of a small number of medications approved to prevent and to treat malaria. If taken correctly, Lariam is (b)(4)-effective at preventing malaria but, like all medications, it may produce side effects in some patients.

**2. Lariam can rarely cause serious mental problems in some patients.**

The most frequently reported side effects with Lariam, such as nausea, difficulty sleeping, and bad dreams are usually mild and do not cause people to stop taking the medicine. However, people taking Lariam occasionally experience severe anxiety, feelings that people are against them, hallucinations (seeing or hearing things that are not there, for example), depression, unusual behavior, or feeling disoriented. It (b)(4)-----  
 (b)(4)-----There have been reports that in som-----  
 -----d. Some patients taking Lariam think about killing themselves, and there have been rare reports of suicides.(b)(4)-----It is not known whether Lariam was responsible for these suicides.

**3. You need to take malaria prevention medicine before you travel to a malaria area, while you are in a malaria area, and after you return from a malaria area.**

Medicines approved in the United States for malaria prevention include Lariam, doxycycline, atovaquone/proguanil, hydroxychloroquine, and chloroquine. Not all of these drugs work equally as well in all areas of the world where there is malaria. The chloroquines, for example, do not work in areas where the malaria parasite has developed resistance to chloroquine. Lariam may be effective against malaria that is resistant to chloroquine or other drugs. All drugs to treat malaria have side effects that are different for each one. For example, some may make your skin more sensitive to sunlight (Lariam does not do this). (b)(4)-----  
 (b)(4)-----

----- However, if you use Lariam to  
 -----ression, restlessness, confusion  
 (possible signs of more serious mental problems), or you develop other serious side effects, contact a doctor or other health care provider.(b)(4)-----it may be necessary to stop taking Lariam and use another malaria prevention medicine i- ----d. If you can't get another medicine, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.

***Who should not take Lariam?***

Do not take Lariam to **prevent** malaria if you

- **have depression or had depression recently**
- **have had recent mental illness or problems**, including anxiety disorder, schizophrenia (a severe type of mental illness), or psychosis (losing touch with reality)
- **have or had seizures (epilepsy or convulsions)**
- **are allergic to quinine or quinidine (medicines related to Lariam)**

**Tell your prescriber about all your medical conditions.** Lariam may not be right for you if you have certain conditions, especially the ones listed below:

- **Heart disease.** Lariam may not be right for you.
- **Pregnancy.** Tell your prescriber if you are pregnant or plan to become pregnant. It is dangerous for the mother and for the unborn baby (fetus) to get malaria during pregnancy. Therefore, ask your prescriber if you should take Lariam or another medicine to prevent malaria while you are pregnant.
- **Breast feeding.** Lariam can pass through your milk and may harm the baby. Therefore, ask your prescriber whether you will need to stop breast feeding or use another medicine.
- **Liver problems.**

**Tell your prescriber about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements.** Some medicines may give you a higher chance of having serious side effects from Lariam.

***How should I take Lariam?***

**Take Lariam exactly as prescribed. If you are an adult or pediatric patient weighing 45 kg (99 pounds) or less, your prescriber will tell you the correct dose based on your weight.**

To prevent malaria

- For adults and pediatric patients weighing over 45 kg, Take 1 tablet of Lariam at least 1 week before you travel to a malaria area (or 2 to 3 weeks before you travel to a malaria area, if instructed by your prescriber). This starts the prevention and also helps you see how Lariam affects you and the other medicines you take. **Take 1 Lariam tablet once a week**, on the same day each week, while in a malaria area.
- **Continue taking Lariam for 4 weeks after returning from a malaria area.** If you cannot continue taking Lariam due to side effects or for other reasons, contact your prescriber.
- Take Lariam just after a meal and with at least 1 cup (8 ounces) of water.
- For children, Lariam can be given with water or crushed and mixed with water or sugar water. The prescriber will tell you the correct dose for children based on the child's weight.
- If you are told by a doctor or other health care provider to stop taking Lariam due to side effects or for other reasons, it will be necessary to take another malaria medicine. You must take **malaria prevention medicine before you travel to a malaria area, while you are in a**

**malaria area, and after you return from a malaria area. If you don't have access to a doctor or other health care provider or to another medicine besides Lariam and have to stop taking it, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.**

### ***What should I avoid while taking Lariam?***

- **Halofantrine (marketed under various brand names)**, a medicine used to treat malaria. Taking both of these medicines together can cause serious heart problems that can cause death.
- **Do not become pregnant.** Women should use effective birth control while taking Lariam.
- **Quinine, quinidine, or chloroquine (other medicines used to treat malaria).** Taking these medicines with Lariam could cause changes in your heart rate or increase the risk of seizures.

### ***In addition:***

- **Be careful driving or in other activities** needing alertness and careful movements (fine motor coordination). Lariam can cause dizziness or loss of balance, even after you stop taking it.
- **Be aware that certain vaccines may not work if given while you are taking Lariam.** Your prescriber may want you to finish taking your vaccines at least 3 days before starting Lariam.

### ***What are the possible side effects of Lariam?***

Lariam, like all medicines, may cause side effects in some patients. The most frequently reported side effects with Lariam when used for prevention of malaria include nausea, vomiting, diarrhea, dizziness, difficulty sleeping, and bad dreams. These are usually mild and do not cause people to stop taking the medicine.

Lariam may cause serious mental problems in some patients. (See "What is the most important information I should know about Lariam?").

Lariam may affect your liver and your eyes if you take it for a long time. Your prescriber will tell you if you should have your eyes and liver checked while taking Lariam.

### ***What else should I know about preventing malaria?***

- **Find out whether you need malaria prevention.** Before you travel, talk with your prescriber about your travel plans to determine whether you need to take medicine to prevent malaria. Even in those countries where malaria is present, there may be areas of the country that are free of malaria. In general, malaria is more common in rural (country) areas than in big cities, and it is more common during rainy seasons, when mosquitoes are most common. You can get information about the areas of the world where malaria occurs from the Centers for Disease Control and Prevention (CDC) and from local authorities in the countries you visit. If possible, plan your travel to reduce the risk of malaria.
- **Take medicine to prevent malaria infection.** Without malaria prevention medicine, you have a higher risk of getting malaria. Malaria starts with flu-like symptoms, such as chills, fever, muscle pains, and headaches. However, malaria can make you very sick or cause death if you don't seek medical help immediately. These symptoms may disappear for a while, and you

may think you are well. But, the symptoms return later and then it may be too late for successful treatment.

Malaria can cause confusion, coma, and seizures. It can cause kidney failure, breathing problems, and severe damage to red blood cells. However, malaria can be easily diagnosed with a blood test, and if caught in time, can be effectively treated.

**If you get flu-like symptoms (chills, fever, muscle pains, or headaches) after you return from a malaria area,** get medical help right away and tell your prescriber that you may have been exposed to malaria.

People who have lived for many years in areas with malaria may have some immunity to malaria (they do not get it as easily) and may not take malaria prevention medicine. This does not mean that you don't need to take malaria prevention medicine.

- **Protect against mosquito bites.** Medicines do not always completely prevent your catching malaria from mosquito bites. So protect yourself very well against mosquitoes. Cover your skin with long sleeves and long pants, and use mosquito repellent and bednets while in malaria areas. If you are out in the bush, you may want to pre-wash your clothes with permethrin. This is a mosquito repellent that may be effective for weeks after use. Ask your prescriber for other ways to protect yourself.

***General information about the safe and effective use of Lariam.***

Medicines are sometimes prescribed for conditions not listed in Medication Guides. If you have any concerns about Lariam, ask your prescriber. This Medication Guide contains certain important information for travelers visiting areas with malaria. Your prescriber or pharmacist can give you information about Lariam that was written for health care professionals. Do not use Lariam for a condition for which it was not prescribed. Do not share Lariam with other people.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Manufactured by:  
F. HOFFMANN-LA ROCHE LTD  
Basel, Switzerland

Distributed by:



**Pharmaceuticals**

Roche Laboratories Inc.  
340 Kingsland Street  
Nutley, New Jersey 07110-1199

27898583

Revised: August 2003

Copyright © 2003 by Roche Laboratories Inc. All rights reserved.



**Information wallet card to carry when you are taking Lariam.**



<b>Lariam® (mefloquine hydrochloride) Tablets</b>	
<p><b>You need to take malaria prevention medicine before you travel to a malaria area, while you are in a malaria area, and after you return from a malaria area.</b></p> <p><b>If taken correctly, Lariam is highly effective at preventing malaria but, like all medications, it may produce side effects in some patients.</b></p> <p><b>If you use Lariam to prevent malaria and you develop a sudden onset of anxiety, depression, restlessness, confusion (possible signs of more serious mental problems), or you develop other serious side effects, contact a doctor or other health care provider as it. <u>It</u> may be necessary to stop taking Lariam and use another malaria prevention medicine instead.</b></p> <p><del>(b)(4)</del> _____</p> <p><u>Revised: August 2003</u></p>	<p><b>Other medicines approved in the United States for malaria prevention include: doxycycline, atovaquone/proguanil (<del>Malarone</del>), hydroxychloroquine, and chloroquine. Not all malaria medicines work equally well in malaria areas. The chloroquines, for example, do not work in many parts of the world. If you can't get another medicine, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.</b></p> <p><b>Please read the Medication Guide for additional information on Lariam.</b></p>

The carton labeling now includes the following statement (in red type) located next to the dosage on the front of the carton:

PHARMACIST: Provide a Medication Guide to each patient when dispensing Lariam.

We have completed the review of this supplemental application, as amended, and have concluded that adequate information has been presented to demonstrate that the drug product is safe and effective for use as recommended in the final printed labeling submitted on August 15, 2003. Accordingly, the supplemental application is approved effective on the date of this letter.

If a letter communicating important information about this drug product (i.e., a "Dear Health Care Professional" letter) is issued to physicians and others responsible for patient care, we request that you submit a copy of the letter to this NDA and a copy to the following address:

MEDWATCH, HF-2  
FDA  
5600 Fishers Lane  
Rockville, MD 20857

We remind you that you must comply with the requirements for an approved NDA set forth under 21 CFR 314.80 and 314.81.

If you have any questions, call Robin Anderson, Labeling Reviewer, at (301) 827-2127.

Sincerely,

*{See appended electronic signature page}*

Renata Albrecht, M.D.  
Director  
Division of Special Pathogen and Immunologic Drug  
Products  
Office of Drug Evaluation IV  
Center for Drug Evaluation and Research

-----  
**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**  
-----

/s/

-----  
Steven Gitterman  
8/20/03 10:23:08 PM  
for Renata Albrecht

**CENTER FOR DRUG EVALUATION AND  
RESEARCH**

*APPLICATION NUMBER:*

**NDA 019591/S-022**

**LABELING**

R<sub>x</sub> only**LARIAM<sup>®</sup>**


brand of

**mefloquine hydrochloride****TABLETS****DESCRIPTION**

Lariam (mefloquine hydrochloride) is an antimalarial agent available as 250-mg tablets of mefloquine hydrochloride (equivalent to 228.0 mg of the free base) for oral administration.

Mefloquine hydrochloride is a 4-quinolinemethanol derivative with the specific chemical name of (R\*, S\*)-(±)-α-2-piperidinyl-2,8-bis (trifluoromethyl)-4-quinolinemethanol hydrochloride. It is a 2-aryl substituted chemical structural analog of quinine. The drug is a white to almost white crystalline compound, slightly soluble in water.

Mefloquine hydrochloride has a calculated molecular weight of 414.78 and the following structural formula:



The inactive ingredients are ammonium-calcium alginate, corn starch, crospovidone, lactose, magnesium stearate, microcrystalline cellulose, poloxamer #331, and talc.

**CLINICAL PHARMACOLOGY****Pharmacokinetics****Absorption**

The absolute oral bioavailability of mefloquine has not been determined since an intravenous formulation is not available. The bioavailability of the tablet formation compared with an oral solution was over 85%. The presence of food significantly enhances the rate and extent of absorption, leading to about a 40% increase in bioavailability. In healthy volunteers, plasma concentrations peak 6 to 24 hours (median, about 17 hours) after a single dose of Lariam. In a similar group of volunteers, maximum plasma concentrations in µg/L are roughly equivalent to the dose in milligrams (for example, a single 1000 mg dose produces a maximum concentration of about 1000 µg/L).

## **LARIAM® (mefloquine hydrochloride)**

In healthy volunteers, a dose of 250 mg once weekly, produces maximum steady-state plasma concentrations of 1000 to 2000 µg/L, which are reached after 7 to 10 weeks.

### **Distribution**

In healthy adults, the apparent volume of distribution is approximately 20 L/kg, indicating extensive tissue distribution. Mefloquine may accumulate in parasitized erythrocytes. Experiments conducted in vitro with human blood using concentrations between 50 and 1000 mg/mL showed a relatively constant erythrocyte-to-plasma concentration ratio of about 2 to 1. The equilibrium reached in less than 30 minutes, was found to be reversible. Protein binding is about 98%.

Mefloquine crosses the placenta. Excretion into breast milk appears to be minimal (see **PRECAUTIONS: Nursing Mothers**).

### **Metabolism**

Two metabolites have been identified in humans. The main metabolite, 2,8-bis-trifluoromethyl-4-quinoline carboxylic acid, is inactive in *Plasmodium falciparum*. In a study in healthy volunteers, the carboxylic acid metabolite appeared in plasma 2 to 4 hours after a single oral dose. Maximum plasma concentrations, which were about 50% higher than those of mefloquine, were reached after 2 weeks. Thereafter, plasma levels of the main metabolite and mefloquine declined at a similar rate. The area under the plasma concentration-time curve (AUC) of the main metabolite was 3 to 5 times larger than that of the parent drug. The other metabolite, an alcohol, was present in minute quantities only.

### **Elimination**

In several studies in healthy adults, the mean elimination half-life of mefloquine varied between 2 and 4 weeks, with an average of about 3 weeks. Total clearance, which is essentially hepatic, is in the order of 30 mL/min. There is evidence that mefloquine is excreted mainly in the bile and feces. In volunteers, urinary excretion of unchanged mefloquine and its main metabolite under steady-state condition accounted for about 9% and 4% of the dose, respectively. Concentrations of other metabolites could not be measured in the urine.

### **Pharmacokinetics in Special Clinical Situations**

#### *Children and the Elderly*

No relevant age-related changes have been observed in the pharmacokinetics of mefloquine. Therefore, the dosage for children has been extrapolated from the recommended adult dose.

No pharmacokinetic studies have been performed in patients with renal insufficiency since only a small proportion of the drug is eliminated renally. Mefloquine and its main metabolite are not appreciably removed by hemodialysis. No special chemoprophylactic dosage adjustments are indicated for dialysis patients to achieve concentrations in plasma similar to those in healthy persons.

## **LARIAM® (mefloquine hydrochloride)**

Although clearance of mefloquine may increase in late pregnancy, in general, pregnancy has no clinically relevant effect on the pharmacokinetics of mefloquine.

The pharmacokinetics of mefloquine may be altered in acute malaria.

Pharmacokinetic differences have been observed between various ethnic populations. In practice, however, these are of minor importance compared with host immune status and sensitivity of the parasite.

During long-term prophylaxis (>2 years), the trough concentrations and the elimination half-life of mefloquine were similar to those obtained in the same population after 6 months of drug use, which is when they reached steady state.

In vitro and in vivo studies showed no hemolysis associated with glucose-6-phosphate dehydrogenase deficiency (see **ANIMAL TOXICOLOGY**).

### **Microbiology**

#### **Mechanism of Action**

Mefloquine is an antimalarial agent which acts as a blood schizonticide. Its exact mechanism of action is not known.

#### **Activity In Vitro and In Vivo**

Mefloquine is active against the erythrocytic stages of *Plasmodium* species (see **INDICATIONS AND USAGE**). However, the drug has no effect against the exoerythrocytic (hepatic) stages of the parasite. Mefloquine is effective against malaria parasites resistant to chloroquine (see **INDICATIONS AND USAGE**).

#### **Drug Resistance**

Strains of *P. falciparum* with decreased susceptibility to mefloquine can be selected in vitro or in vivo. Resistance of *P. falciparum* to mefloquine has been reported in areas of multi-drug resistance in South East Asia. Increased incidences of resistance have also been reported in other parts of the world.

#### **Cross Resistance**

Cross-resistance between mefloquine and halofantrine and cross-resistance between mefloquine and quinine have been observed in some regions.

## **INDICATIONS AND USAGE**

### **Treatment of Acute Malaria Infections**

Lariam is indicated for the treatment of mild to moderate acute malaria caused by mefloquine-susceptible strains of *P. falciparum* (both chloroquine-susceptible and resistant strains) or by *Plasmodium vivax*. There are insufficient clinical data to document the effect of mefloquine in malaria caused by *P. ovale* or *P. malariae*.

## **LARIAM® (mefloquine hydrochloride)**

*Note:* Patients with acute *P. vivax* malaria, treated with Lariam, are at high risk of relapse because Lariam does not eliminate exoerythrocytic (hepatic phase) parasites. To avoid relapse, after initial treatment of the acute infection with Lariam, patients should subsequently be treated with an 8-aminoquinoline derivative (eg, primaquine).

### **Prevention of Malaria**

Lariam is indicated for the prophylaxis of *P. falciparum* and *P. vivax* malaria infections, including prophylaxis of chloroquine-resistant strains of *P. falciparum*.

### **CONTRAINDICATIONS**

Use of Lariam is contraindicated in patients with a known hypersensitivity to mefloquine or related compounds (eg, quinine and quinidine) or to any of the excipients contained in the formulation. Lariam should not be prescribed for prophylaxis in patients with active depression, a recent history of depression, generalized anxiety disorder, psychosis, or schizophrenia or other major psychiatric disorders, or with a history of convulsions.

### **WARNINGS**

**In case of life-threatening, serious or overwhelming malaria infections due to *P. falciparum*, patients should be treated with an intravenous antimalarial drug. Following completion of intravenous treatment, Lariam may be given to complete the course of therapy.**

**Data on the use of halofantrine subsequent to administration of Lariam suggest a significant, potentially fatal prolongation of the QTc interval of the ECG. Therefore, halofantrine must not be given simultaneously with or subsequent to Lariam. No data are available on the use of Lariam after halofantrine (see PRECAUTIONS: Drug Interactions).**

**Mefloquine may cause psychiatric symptoms in a number of patients, ranging from anxiety, paranoia, and depression to hallucinations and psychotic behavior. On occasions, these symptoms have been reported to continue long after mefloquine has been stopped. Rare cases of suicidal ideation and suicide have been reported though no relationship to drug administration has been confirmed. To minimize the chances of these adverse events, mefloquine should not be taken for prophylaxis in patients with active depression or with a recent history of depression, generalized anxiety disorder, psychosis, or schizophrenia or other major psychiatric disorders. Lariam should be used with caution in patients with a previous history of depression.**

**During prophylactic use, if psychiatric symptoms such as acute anxiety, depression, restlessness or confusion occur, these may be considered prodromal to a more serious event. In these cases, the drug must be discontinued and an alternative medication should be substituted.**



## **LARIAM® (mefloquine hydrochloride)**

**Concomitant administration of Lariam and quinine or quinidine may produce electrocardiographic abnormalities.**

**Concomitant administration of Lariam and quinine or chloroquine may increase the risk of convulsions.**

### **PRECAUTIONS**

#### **General**

Hypersensitivity reactions ranging from mild cutaneous events to anaphylaxis cannot be predicted.

In patients with epilepsy, Lariam may increase the risk of convulsions. The drug should therefore be prescribed only for curative treatment in such patients and only if there are compelling medical reasons for its use (see **PRECAUTIONS: Drug Interactions**).

Caution should be exercised with regard to activities requiring alertness and fine motor coordination such as driving, piloting aircraft, operating machinery, and deep-sea diving, as dizziness, a loss of balance, or other disorders of the central or peripheral nervous system have been reported during and following the use of Lariam. These effects may occur after therapy is discontinued due to the long half-life of the drug. Lariam should be used with caution in patients with psychiatric disturbances because mefloquine use has been associated with emotional disturbances (see **ADVERSE REACTIONS**).

In patients with impaired liver function the elimination of mefloquine may be prolonged, leading to higher plasma levels.

This drug has been administered for longer than 1 year. If the drug is to be administered for a prolonged period, periodic evaluations including liver function tests should be performed. Although retinal abnormalities seen in humans with long-term chloroquine use have not been observed with mefloquine use, long-term feeding of mefloquine to rats resulted in dose-related ocular lesions (retinal degeneration, retinal edema and lenticular opacity at 12.5 mg/kg/day and higher) (see **ANIMAL TOXICOLOGY**). Therefore, periodic ophthalmic examinations are recommended.

Parenteral studies in animals show that mefloquine, a myocardial depressant, possesses 20% of the antifibrillatory action of quinidine and produces 50% of the increase in the PR interval reported with quinine. The effect of mefloquine on the compromised cardiovascular system has not been evaluated. However, transitory and clinically silent ECG alterations have been reported during the use of mefloquine. Alterations included sinus bradycardia, sinus arrhythmia, first degree AV-block, prolongation of the QTc interval and abnormal T waves (see also cardiovascular effects under **PRECAUTIONS: Drug Interactions** and **ADVERSE REACTIONS**). The benefits of Lariam therapy should be weighed against the possibility of adverse effects in patients with cardiac disease.

## **LARIAM® (mefloquine hydrochloride)**

### **Laboratory Tests**

Periodic evaluation of hepatic function should be performed during prolonged prophylaxis.

### **Information for Patients**

Medication Guide: As required by law, a Lariam Medication Guide is supplied to patients when Lariam is dispensed. Patients should be instructed to read the MedGuide when Lariam is received. The complete text of the MedGuide is reprinted at the end of this document.

Patients should be advised:

- that malaria can be a life-threatening infection in the traveler;
- that Lariam is being prescribed to help prevent or treat this serious infection;
- that in a small percentage of cases, patients are unable to take this medication because of side effects, and it may be necessary to change medications;
- that when used as prophylaxis, the first dose of Lariam should be taken 1 week prior to arrival in an endemic area;
- that if the patients experience psychiatric symptoms such as acute anxiety, depression, restlessness or confusion, these may be considered prodromal to a more serious event. In these cases, the drug must be discontinued and an alternative medication should be substituted;
- that no chemoprophylactic regimen is 100% effective, and protective clothing, insect repellents, and bednets are important components of malaria prophylaxis;
- to seek medical attention for any febrile illness that occurs after return from a malarious area and to inform their physician that they may have been exposed to malaria.

### **Drug Interactions**

Drug-drug interactions with Lariam have not been explored in detail. There is one report of cardiopulmonary arrest, with full recovery, in a patient who was taking a beta blocker (propranolol) (see **PRECAUTIONS: General**). The effects of mefloquine on the compromised cardiovascular system have not been evaluated. The benefits of Lariam therapy should be weighed against the possibility of adverse effects in patients with cardiac disease.

Because of the danger of a potentially fatal prolongation of the QTc interval, halofantrine must not be given simultaneously with or subsequent to Lariam (see **WARNINGS**).

Concomitant administration of Lariam and other related compounds (eg, quinine, quinidine and chloroquine) may produce electrocardiographic abnormalities and increase the risk of convulsions (see **WARNINGS**). If these drugs are to be used in the initial treatment of severe malaria, Lariam administration should be delayed at least 12 hours after the last dose. There is evidence that the use of halofantrine after mefloquine causes a significant lengthening of the QTc interval. Clinically significant QTc prolongation has not been found with mefloquine alone.

## **LARIAM® (mefloquine hydrochloride)**

This appears to be the only clinically relevant interaction of this kind with Lariam, although theoretically, coadministration of other drugs known to alter cardiac conduction (eg, anti-arrhythmic or beta-adrenergic blocking agents, calcium channel blockers, antihistamines or H<sub>1</sub>-blocking agents, tricyclic antidepressants and phenothiazines) might also contribute to a prolongation of the QTc interval. There are no data that conclusively establish whether the concomitant administration of mefloquine and the above listed agents has an effect on cardiac function.

In patients taking an anticonvulsant (eg, valproic acid, carbamazepine, phenobarbital or phenytoin), the concomitant use of Lariam may reduce seizure control by lowering the plasma levels of the anticonvulsant. Therefore, patients concurrently taking antiseizure medication and Lariam should have the blood level of their antiseizure medication monitored and the dosage adjusted appropriately (see **PRECAUTIONS: General**).

When Lariam is taken concurrently with oral live typhoid vaccines, attenuation of immunization cannot be excluded. Vaccinations with attenuated live bacteria should therefore be completed at least 3 days before the first dose of Lariam.

No other drug interactions are known. Nevertheless, the effects of Lariam on travelers receiving comedication, particularly diabetics or patients using anticoagulants, should be checked before departure.

In clinical trials, the concomitant administration of sulfadoxine and pyrimethamine did not alter the adverse reaction profile.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### **Carcinogenesis**

The carcinogenic potential of mefloquine was studied in rats and mice in 2-year feeding studies at doses of up to 30 mg/kg/day. No treatment-related increases in tumors of any type were noted.

#### **Mutagenesis**

The mutagenic potential of mefloquine was studied in a variety of assay systems including: Ames test, a host-mediated assay in mice, fluctuation tests and a mouse micronucleus assay. Several of these assays were performed with and without prior metabolic activation. In no instance was evidence obtained for the mutagenicity of mefloquine.

#### **Impairment of Fertility**

Fertility studies in rats at doses of 5, 20, and 50 mg/kg/day of mefloquine have demonstrated adverse effects on fertility in the male at the high dose of 50 mg/kg/day, and in the female at doses of 20 and 50 mg/kg/day. Histopathological lesions were noted in the epididymides from male rats at doses of 20 and 50 mg/kg/day. Administration of 250 mg/week of mefloquine (base) in adult males for 22 weeks failed to reveal any deleterious effects on human spermatozoa.

## **LARIAM® (mefloquine hydrochloride)**

### **Pregnancy**

#### **Teratogenic Effects**

Pregnancy Category C. Mefloquine has been demonstrated to be teratogenic in rats and mice at a dose of 100 mg/kg/day. In rabbits, a high dose of 160 mg/kg/day was embryotoxic and teratogenic, and a dose of 80 mg/kg/day was teratogenic but not embryotoxic. There are no adequate and well-controlled studies in pregnant women. However, clinical experience with Lariam has not revealed an embryotoxic or teratogenic effect. Mefloquine should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Women of childbearing potential who are traveling to areas where malaria is endemic should be warned against becoming pregnant. Women of childbearing potential should also be advised to practice contraception during malaria prophylaxis with Lariam and for up to 3 months thereafter. However, in the case of unplanned pregnancy, malaria chemoprophylaxis with Lariam is not considered an indication for pregnancy termination.

### **Nursing Mothers**

Mefloquine is excreted in human milk in small amounts, the activity of which is unknown. Based on a study in a few subjects, low concentrations (3% to 4%) of mefloquine were excreted in human milk following a dose equivalent to 250 mg of the free base. Because of the potential for serious adverse reactions in nursing infants from mefloquine, a decision should be made whether to discontinue the drug, taking into account the importance of the drug to the mother.

### **Pediatric Use**

Use of Lariam to treat acute, uncomplicated *P. falciparum* malaria in pediatric patients is supported by evidence from adequate and well-controlled studies of Lariam in adults with additional data from published open-label and comparative trials using Lariam to treat malaria caused by *P. falciparum* in patients younger than 16 years of age. The safety and effectiveness of Lariam for the treatment of malaria in pediatric patients below the age of 6 months have not been established.

In several studies, the administration of Lariam for the treatment of malaria was associated with early vomiting in pediatric patients. Early vomiting was cited in some reports as a possible cause of treatment failure. If a second dose is not tolerated, the patient should be monitored closely and alternative malaria treatment considered if improvement is not observed within a reasonable period of time (see **DOSAGE AND ADMINISTRATION**).

### **Geriatric Use**

Clinical studies of Lariam did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. Since electrocardiographic abnormalities have been observed in individuals treated with Lariam (see **PRECAUTIONS**) and underlying cardiac disease is

## **LARIAM® (mefloquine hydrochloride)**

more prevalent in elderly than in younger patients, the benefits of Lariam therapy should be weighed against the possibility of adverse cardiac effects in elderly patients.

### **ADVERSE REACTIONS**

#### **Clinical**

At the doses used for treatment of acute malaria infections, the symptoms possibly attributable to drug administration cannot be distinguished from those symptoms usually attributable to the disease itself.

Among subjects who received mefloquine for prophylaxis of malaria, the most frequently observed adverse experience was vomiting (3%). Dizziness, syncope, extrasystoles and other complaints affecting less than 1% were also reported.

Among subjects who received mefloquine for treatment, the most frequently observed adverse experiences included: dizziness, myalgia, nausea, fever, headache, vomiting, chills, diarrhea, skin rash, abdominal pain, fatigue, loss of appetite, and tinnitus. Those side effects occurring in less than 1% included bradycardia, hair loss, emotional problems, pruritus, asthenia, transient emotional disturbances and telogen effluvium (loss of resting hair). Seizures have also been reported.

Two serious adverse reactions were cardiopulmonary arrest in one patient shortly after ingesting a single prophylactic dose of mefloquine while concomitantly using propranolol (see **PRECAUTIONS: Drug Interactions**), and encephalopathy of unknown etiology during prophylactic mefloquine administration. The relationship of encephalopathy to drug administration could not be clearly established.

#### **Postmarketing**

Postmarketing surveillance indicates that the same kind of adverse experiences are reported during prophylaxis, as well as acute treatment.

The most frequently reported adverse events are nausea, vomiting, loose stools or diarrhea, abdominal pain, dizziness or vertigo, loss of balance, and neuropsychiatric events such as headache, somnolence, and sleep disorders (insomnia, abnormal dreams). These are usually mild and may decrease despite continued use.

Occasionally, more severe neuropsychiatric disorders have been reported such as: sensory and motor neuropathies (including paresthesia, tremor and ataxia), convulsions, agitation or restlessness, anxiety, depression, mood changes, panic attacks, forgetfulness, confusion, hallucinations, aggression, psychotic or paranoid reactions and encephalopathy. Rare cases of suicidal ideation and suicide have been reported though no relationship to drug administration has been confirmed.

Other infrequent adverse events include:

*Cardiovascular Disorders:* circulatory disturbances (hypotension, hypertension, flushing, syncope), chest pain, tachycardia or palpitation, bradycardia, irregular pulse, extrasystoles, A-V block, and other transient cardiac conduction alterations

## **LARIAM® (mefloquine hydrochloride)**

*Skin Disorders:* rash, exanthema, erythema, urticaria, pruritus, edema, hair loss, erythema multiforme, and Stevens-Johnson syndrome

*Musculoskeletal Disorders:* muscle weakness, muscle cramps, myalgia, and arthralgia

*Other Symptoms:* visual disturbances, vestibular disorders including tinnitus and hearing impairment, dyspnea, asthenia, malaise, fatigue, fever, sweating, chills, dyspepsia and loss of appetite

### **Laboratory**

The most frequently observed laboratory alterations which could be possibly attributable to drug administration were decreased hematocrit, transient elevation of transaminases, leukopenia and thrombocytopenia. These alterations were observed in patients with acute malaria who received treatment doses of the drug and were attributed to the disease itself.

During prophylactic administration of mefloquine to indigenous populations in malaria-endemic areas, the following occasional alterations in laboratory values were observed: transient elevation of transaminases, leukocytosis or thrombocytopenia.

Because of the long half-life of mefloquine, adverse reactions to Lariam may occur or persist up to several weeks after the last dose.

### **OVERDOSAGE**

In cases of overdosage with Lariam, the symptoms mentioned under **ADVERSE REACTIONS** may be more pronounced. The following procedure is recommended in case of overdosage: Induce vomiting or perform gastric lavage, as appropriate. Monitor cardiac function (if possible by ECG) and neuropsychiatric status for at least 24 hours. Provide symptomatic and intensive supportive treatment as required, particularly for cardiovascular disturbances.

### **DOSAGE AND ADMINISTRATION (see INDICATIONS AND USAGE)**

#### **Adult Patients**

Treatment of mild to moderate malaria in adults caused by *P. vivax* or mefloquine-susceptible strains of *P. falciparum*

Five tablets (1250 mg) mefloquine hydrochloride to be given as a single oral dose. The drug should not be taken on an empty stomach and should be administered with at least 8 oz (240 mL) of water.

If a full-treatment course with Lariam does not lead to improvement within 48 to 72 hours, Lariam should not be used for retreatment. An alternative therapy should be used. Similarly, if previous prophylaxis with mefloquine has failed, Lariam should not be used for curative treatment.

*Note:* Patients with acute *P. vivax* malaria, treated with Lariam, are at high risk of relapse because Lariam does not eliminate exoerythrocytic (hepatic phase) parasites. To avoid relapse after initial treatment of the acute infection with

## **LARIAM® (mefloquine hydrochloride)**

Lariam, patients should subsequently be treated with an 8-aminoquinoline derivative (eg, primaquine).

### **Malaria Prophylaxis**

One 250 mg Lariam tablet once weekly.

Prophylactic drug administration should begin 1 week before arrival in an endemic area. Subsequent weekly doses should be taken regularly, always on the same day of each week, preferably after the main meal. To reduce the risk of malaria after leaving an endemic area, prophylaxis must be continued for 4 additional weeks to ensure suppressive blood levels of the drug when merozoites emerge from the liver. Tablets should not be taken on an empty stomach and should be administered with at least 8 oz (240 mL) of water.

In certain cases, eg, when a traveler is taking other medication, it may be desirable to start prophylaxis 2 to 3 weeks prior to departure, in order to ensure that the combination of drugs is well tolerated (see **PRECAUTIONS: Drug Interactions**).

When prophylaxis with Lariam fails, physicians should carefully evaluate which antimalarial to use for therapy.

### **Pediatric Patients**

Treatment of mild to moderate malaria in pediatric patients caused by mefloquine-susceptible strains of *P. falciparum*

Twenty (20) to 25 mg/kg body weight. Splitting the total therapeutic dose into 2 doses taken 6 to 8 hours apart may reduce the occurrence or severity of adverse effects. Experience with Lariam in infants less than 3 months old or weighing less than 5 kg is limited. The drug should not be taken on an empty stomach and should be administered with ample water. The tablets may be crushed and suspended in a small amount of water, milk or other beverage for administration to small children and other persons unable to swallow them whole.

If a full-treatment course with Lariam does not lead to improvement within 48 to 72 hours, Lariam should not be used for retreatment. An alternative therapy should be used. Similarly, if previous prophylaxis with mefloquine has failed, Lariam should not be used for curative treatment.

In pediatric patients, the administration of Lariam for the treatment of malaria has been associated with early vomiting. In some cases, early vomiting has been cited as a possible cause of treatment failure (see **PRECAUTIONS**). If a significant loss of drug product is observed or suspected because of vomiting, a second full dose of Lariam should be administered to patients who vomit less than 30 minutes after receiving the drug. If vomiting occurs 30 to 60 minutes after a dose, an additional half-dose should be given. If vomiting recurs, the patient should be monitored closely and alternative malaria treatment considered if improvement is not observed within a reasonable period of time.

## **LARIAM® (mefloquine hydrochloride)**

The safety and effectiveness of Lariam to treat malaria in pediatric patients below the age of 6 months have not been established.

### **Malaria Prophylaxis**

The following doses have been extrapolated from the recommended adult dose. Neither the pharmacokinetics, nor the clinical efficacy of these doses have been determined in children owing to the difficulty of acquiring this information in pediatric subjects. The recommended prophylactic dose of Lariam is approximately 5 mg/kg body weight once weekly. One 250 mg Lariam tablet should be taken once weekly in pediatric patients weighing over 45 kg. In pediatric patients weighing less than 45 kg, the weekly dose decreases in proportion to body weight:

30 to 45 kg:	3/4 tablet
20 to 30 kg:	1/2 tablet
10 to 20 kg:	1/4 tablet
5 to 10 kg:	1/8 tablet*

\*Approximate tablet fraction based on a dosage of 5 mg/kg body weight. Exact doses for children weighing less than 10 kg may best be prepared and dispensed by pharmacists.

Experience with Lariam in infants less than 3 months old or weighing less than 5 kg is limited.

### **HOW SUPPLIED**

Lariam is available as scored, white, round tablets, containing 250 mg of mefloquine hydrochloride in unit-dose packages of 25 (NDC 0004-0172-02). Imprint on tablets: LARIAM 250 ROCHE

Tablets should be stored at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F).

### **ANIMAL TOXICOLOGY**

Ocular lesions were observed in rats fed mefloquine daily for 2 years. All surviving rats given 30 mg/kg/day had ocular lesions in both eyes characterized by retinal degeneration, opacity of the lens, and retinal edema. Similar but less severe lesions were observed in 80% of female and 22% of male rats fed 12.5 mg/kg/day for 2 years. At doses of 5 mg/kg/day, only corneal lesions were observed. They occurred in 9% of rats studied.

Revised: August 2003



## **LARIAM® (mefloquine hydrochloride)**

### **MEDICATION GUIDE**

**This Medication Guide is intended only for travelers who are taking Lariam to prevent malaria.** The information may not apply to patients who are sick with malaria and who are taking Lariam to treat malaria.

An information wallet card is provided at the end of this Medication Guide. Cut it out and carry it with you when you are taking Lariam.

This Medication Guide was revised in August 2003. Please read it before you start taking Lariam and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking with your prescriber (doctor or other health care provider) about Lariam and malaria prevention. Only you and your prescriber can decide if Lariam is right for you. If you cannot take Lariam, you may be able to take a different medicine to prevent malaria.

### **What is the most important information I should know about Lariam?**

**1. Take Lariam exactly as prescribed to prevent malaria.**

Malaria is an infection that can cause death and is spread to humans through mosquito bites. If you travel to parts of the world where the mosquitoes carry the malaria parasite, you must take a malaria prevention medicine. Lariam is one of a small number of medications approved to prevent and to treat malaria. If taken correctly, Lariam is effective at preventing malaria but, like all medications, it may produce side effects in some patients.

**2. Lariam can rarely cause serious mental problems in some patients.**

The most frequently reported side effects with Lariam, such as nausea, difficulty sleeping, and bad dreams are usually mild and do not cause people to stop taking the medicine. However, people taking Lariam occasionally experience severe anxiety, feelings that people are against them, hallucinations (seeing or hearing things that are not there, for example), depression, unusual behavior, or feeling disoriented. There have been reports that in some patients these side effects continue after Lariam is stopped. Some patients taking Lariam think about killing themselves, and there have been rare reports of suicides. It is not known whether Lariam was responsible for these suicides.

**3. You need to take malaria prevention medicine before you travel to a malaria area, while you are in a malaria area, and after you return from a malaria area.**

Medicines approved in the United States for malaria prevention include Lariam, doxycycline, atovaquone/proguanil, hydroxychloroquine, and chloroquine. Not all of these drugs work equally as well in all areas of the world where there is malaria. The chloroquines, for example, do not work in areas where the malaria parasite has developed resistance to chloroquine. Lariam may be effective against malaria that is resistant to chloroquine or other drugs. All drugs to treat malaria have side effects that are different for each one. For example, some may make your skin more sensitive to sunlight (Lariam does not do this). However, if you use Lariam to prevent malaria

## **LARIAM® (mefloquine hydrochloride)**

and you develop a sudden onset of anxiety, depression, restlessness, confusion (possible signs of more serious mental problems), or you develop other serious side effects, contact a doctor or other health care provider. It may be necessary to stop taking Lariam and use another malaria prevention medicine instead. If you can't get another medicine, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.

### **Who should not take Lariam?**

Do not take Lariam to **prevent** malaria if you

- **have depression or had depression recently**
- **have had recent mental illness or problems**, including anxiety disorder, schizophrenia (a severe type of mental illness), or psychosis (losing touch with reality)
- **have or had seizures (epilepsy or convulsions)**
- **are allergic to quinine or quinidine (medicines related to Lariam)**

**Tell your prescriber about all your medical conditions.** Lariam may not be right for you if you have certain conditions, especially the ones listed below:

- **Heart disease.** Lariam may not be right for you.
- **Pregnancy.** Tell your prescriber if you are pregnant or plan to become pregnant. It is dangerous for the mother and for the unborn baby (fetus) to get malaria during pregnancy. Therefore, ask your prescriber if you should take Lariam or another medicine to prevent malaria while you are pregnant.
- **Breast feeding.** Lariam can pass through your milk and may harm the baby. Therefore, ask your prescriber whether you will need to stop breast feeding or use another medicine.
- **Liver problems.**

**Tell your prescriber about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements.** Some medicines may give you a higher chance of having serious side effects from Lariam.

## **LARIAM® (mefloquine hydrochloride)**

### **How should I take Lariam?**

**Take Lariam exactly as prescribed. If you are an adult or pediatric patient weighing 45 kg (99 pounds) or less, your prescriber will tell you the correct dose based on your weight.**

To prevent malaria

- For adults and pediatric patients weighing over 45 kg, take 1 tablet of Lariam at least 1 week before you travel to a malaria area (or 2 to 3 weeks before you travel to a malaria area, if instructed by your prescriber). This starts the prevention and also helps you see how Lariam affects you and the other medicines you take. **Take 1 Lariam tablet once a week**, on the same day each week, while in a malaria area.
- **Continue taking Lariam for 4 weeks after returning from a malaria area.** If you cannot continue taking Lariam due to side effects or for other reasons, contact your prescriber.
- Take Lariam just after a meal and with at least 1 cup (8 ounces) of water.
- For children, Lariam can be given with water or crushed and mixed with water or sugar water. The prescriber will tell you the correct dose for children based on the child's weight.
- If you are told by a doctor or other health care provider to stop taking Lariam due to side effects or for other reasons, it will be necessary to take another malaria medicine. You must take **malaria prevention medicine before you travel to a malaria area, while you are in a malaria area, and after you return from a malaria area. If you don't have access to a doctor or other health care provider or to another medicine besides Lariam and have to stop taking it, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.**

### **What should I avoid while taking Lariam?**

- **Halofantrine (marketed under various brand names)**, a medicine used to treat malaria. Taking both of these medicines together can cause serious heart problems that can cause death.
- **Do not become pregnant.** Women should use effective birth control while taking Lariam.
- **Quinine, quinidine, or chloroquine (other medicines used to treat malaria).** Taking these medicines with Lariam could cause changes in your heart rate or increase the risk of seizures.

## **LARIAM® (mefloquine hydrochloride)**

### **In addition:**

- **Be careful driving or in other activities** needing alertness and careful movements (fine motor coordination). Lariam can cause dizziness or loss of balance, even after you stop taking it.
- **Be aware that certain vaccines may not work if given while you are taking Lariam.** Your prescriber may want you to finish taking your vaccines at least 3 days before starting Lariam.

### **What are the possible side effects of Lariam?**

Lariam, like all medicines, may cause side effects in some patients. The most frequently reported side effects with Lariam when used for prevention of malaria include nausea, vomiting, diarrhea, dizziness, difficulty sleeping, and bad dreams. These are usually mild and do not cause people to stop taking the medicine.

Lariam may cause serious mental problems in some patients. (See “What is the most important information I should know about Lariam?”).

Lariam may affect your liver and your eyes if you take it for a long time. Your prescriber will tell you if you should have your eyes and liver checked while taking Lariam.

### **What else should I know about preventing malaria?**

- **Find out whether you need malaria prevention.** Before you travel, talk with your prescriber about your travel plans to determine whether you need to take medicine to prevent malaria. Even in those countries where malaria is present, there may be areas of the country that are free of malaria. In general, malaria is more common in rural (country) areas than in big cities, and it is more common during rainy seasons, when mosquitoes are most common. You can get information about the areas of the world where malaria occurs from the Centers for Disease Control and Prevention (CDC) and from local authorities in the countries you visit. If possible, plan your travel to reduce the risk of malaria.
- **Take medicine to prevent malaria infection.** Without malaria prevention medicine, you have a higher risk of getting malaria. Malaria starts with flu-like symptoms, such as chills, fever, muscle pains, and headaches. However, malaria can make you very sick or cause death if you don't seek medical help immediately. These symptoms may disappear for a while, and you may think you are well. But, the symptoms return later and then it may be too late for successful treatment.

Malaria can cause confusion, coma, and seizures. It can cause kidney failure, breathing problems, and severe damage to red blood cells. However, malaria can be easily diagnosed with a blood test, and if caught in time, can be effectively treated.

**If you get flu-like symptoms (chills, fever, muscle pains, or headaches) after you return from a malaria area,** get medical help right away and tell your prescriber that you may have been exposed to malaria.

## **LARIAM® (mefloquine hydrochloride)**

People who have lived for many years in areas with malaria may have some immunity to malaria (they do not get it as easily) and may not take malaria prevention medicine. This does not mean that you don't need to take malaria prevention medicine.

- **Protect against mosquito bites.** Medicines do not always completely prevent your catching malaria from mosquito bites. So protect yourself very well against mosquitoes. Cover your skin with long sleeves and long pants, and use mosquito repellent and bednets while in malaria areas. If you are out in the bush, you may want to pre-wash your clothes with permethrin. This is a mosquito repellent that may be effective for weeks after use. Ask your prescriber for other ways to protect yourself.

### **General information about the safe and effective use of Lariam.**

Medicines are sometimes prescribed for conditions not listed in Medication Guides. If you have any concerns about Lariam, ask your prescriber. This Medication Guide contains certain important information for travelers visiting areas with malaria. Your prescriber or pharmacist can give you information about Lariam that was written for health care professionals. Do not use Lariam for a condition for which it was not prescribed. Do not share Lariam with other people.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: August 2003

## LARIAM® (mefloquine hydrochloride)

Information wallet card to carry when you are taking Lariam.



### Lariam® (mefloquine hydrochloride) Tablets

**You need to take malaria prevention medicine before you travel to a malaria area, while you are in a malaria area, and after you return from a malaria area.**

**If taken correctly, Lariam is effective at preventing malaria but, like all medications, it may produce side effects in some patients.**

**If you use Lariam to prevent malaria and you develop a sudden onset of anxiety, depression, restlessness, confusion (possible signs of more serious mental problems), or you develop other serious side effects, contact a doctor or other health care provider. It may be necessary to stop taking Lariam and use another malaria prevention medicine instead.**

Revised: August 2003

**Other medicines approved in the United States for malaria prevention include: doxycycline, atovaquone/proguanil, hydroxychloroquine, and chloroquine. Not all malaria medicines work equally well in malaria areas. The chloroquines, for example, do not work in many parts of the world. If you can't get another medicine, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.**

**Please read the Medication Guide for additional information on Lariam.**

Manufactured by:  
F. HOFFMANN-LA ROCHE LTD  
Basel, Switzerland

Distributed by:



Pharmaceuticals

Roche Laboratories Inc.  
340 Kingsland Street  
Nutley, New Jersey 07110-1199

27898582

Copyright © 1999-2003 by Roche Laboratories Inc. All rights reserved.



R<sub>x</sub> only

**MEDICATION GUIDE**  
**LARIAM<sup>®</sup> (LAH-ree-am)**  
**(mefloquine hydrochloride) Tablets**  
**to Prevent Malaria**

**This Medication Guide is intended only for travelers who are taking Lariam to prevent malaria.** The information may not apply to patients who are sick with malaria and who are taking Lariam to treat malaria.

An information wallet card is provided at the end of this Medication Guide. Cut it out and carry it with you when you are taking Lariam.

This Medication Guide was revised in August 2003. Please read it before you start taking Lariam and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking with your prescriber (doctor or other health care provider) about Lariam and malaria prevention. Only you and your prescriber can decide if Lariam is right for you. If you cannot take Lariam, you may be able to take a different medicine to prevent malaria.

**What is the most important information I should know about Lariam?**

**1. Take Lariam exactly as prescribed to prevent malaria.**

Malaria is an infection that can cause death and is spread to humans through mosquito bites. If you travel to parts of the world where the mosquitoes carry the malaria parasite, you must take a malaria prevention medicine. Lariam is one of a small number of medications approved to prevent and to treat malaria. If taken correctly, Lariam is effective at preventing malaria but, like all medications, it may produce side effects in some patients.

**2. Lariam can rarely cause serious mental problems in some patients.**

The most frequently reported side effects with Lariam, such as nausea, difficulty sleeping, and bad dreams are usually mild and do not cause people to stop taking the medicine. However, people taking Lariam occasionally experience severe anxiety, feelings that people are against them, hallucinations (seeing or hearing things that are not there, for example), depression, unusual behavior, or feeling disoriented. There have been reports that in some patients these side effects continue after Lariam is stopped. Some patients taking Lariam think about killing themselves, and there have been rare reports of suicides. It is not known whether Lariam was responsible for these suicides.

## **LARIAM® (mefloquine hydrochloride)**

### **3. You need to take malaria prevention medicine before you travel to a malaria area, while you are in a malaria area, and after you return from a malaria area.**

Medicines approved in the United States for malaria prevention include Lariam, doxycycline, atovaquone/proguanil, hydroxychloroquine, and chloroquine. Not all of these drugs work equally as well in all areas of the world where there is malaria. The chloroquines, for example, do not work in areas where the malaria parasite has developed resistance to chloroquine. Lariam may be effective against malaria that is resistant to chloroquine or other drugs. All drugs to treat malaria have side effects that are different for each one. For example, some may make your skin more sensitive to sunlight (Lariam does not do this). However, if you use Lariam to prevent malaria and you develop a sudden onset of anxiety, depression, restlessness, confusion (possible signs of more serious mental problems), or you develop other serious side effects, contact a doctor or other health care provider. It may be necessary to stop taking Lariam and use another malaria prevention medicine instead. If you can't get another medicine, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.

### **Who should not take Lariam?**

Do not take Lariam to **prevent** malaria if you

- **have depression or had depression recently**
- **have had recent mental illness or problems**, including anxiety disorder, schizophrenia (a severe type of mental illness), or psychosis (losing touch with reality)
- **have or had seizures (epilepsy or convulsions)**
- **are allergic to quinine or quinidine (medicines related to Lariam)**

**Tell your prescriber about all your medical conditions.** Lariam may not be right for you if you have certain conditions, especially the ones listed below:

- **Heart disease.** Lariam may not be right for you.
- **Pregnancy.** Tell your prescriber if you are pregnant or plan to become pregnant. It is dangerous for the mother and for the unborn baby (fetus) to get malaria during pregnancy. Therefore, ask your prescriber if you should take Lariam or another medicine to prevent malaria while you are pregnant.
- **Breast feeding.** Lariam can pass through your milk and may harm the baby. Therefore, ask your prescriber whether you will need to stop breast feeding or use another medicine.



## **LARIAM® (mefloquine hydrochloride)**

- **Liver problems.**

**Tell your prescriber about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements.** Some medicines may give you a higher chance of having serious side effects from Lariam.

### **How should I take Lariam?**

**Take Lariam exactly as prescribed. If you are an adult or pediatric patient weighing 45 kg (99 pounds) or less, your prescriber will tell you the correct dose based on your weight.**

To prevent malaria

- For adults and pediatric patients weighing over 45 kg, take 1 tablet of Lariam at least 1 week before you travel to a malaria area (or 2 to 3 weeks before you travel to a malaria area, if instructed by your prescriber). This starts the prevention and also helps you see how Lariam affects you and the other medicines you take. **Take 1 Lariam tablet once a week, on the same day each week, while in a malaria area.**
- **Continue taking Lariam for 4 weeks after returning from a malaria area.** If you cannot continue taking Lariam due to side effects or for other reasons, contact your prescriber.
- Take Lariam just after a meal and with at least 1 cup (8 ounces) of water.
- For children, Lariam can be given with water or crushed and mixed with water or sugar water. The prescriber will tell you the correct dose for children based on the child's weight.
- If you are told by a doctor or other health care provider to stop taking Lariam due to side effects or for other reasons, it will be necessary to take another malaria medicine. **You must take malaria prevention medicine before you travel to a malaria area, while you are in a malaria area, and after you return from a malaria area. If you don't have access to a doctor or other health care provider or to another medicine besides Lariam and have to stop taking it, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.**

### **What should I avoid while taking Lariam?**

- **Halofantrine (marketed under various brand names),** a medicine used to treat malaria. Taking both of these medicines together can cause serious heart problems that can cause death.
- **Do not become pregnant.** Women should use effective birth control while taking Lariam.

## **LARIAM® (mefloquine hydrochloride)**

- **Quinine, quinidine, or chloroquine (other medicines used to treat malaria).** Taking these medicines with Lariam could cause changes in your heart rate or increase the risk of seizures.

### **In addition:**

- **Be careful driving or in other activities** needing alertness and careful movements (fine motor coordination). Lariam can cause dizziness or loss of balance, even after you stop taking it.
- **Be aware that certain vaccines may not work if given while you are taking Lariam.** Your prescriber may want you to finish taking your vaccines at least 3 days before starting Lariam.

### **What are the possible side effects of Lariam?**

Lariam, like all medicines, may cause side effects in some patients. The most frequently reported side effects with Lariam when used for prevention of malaria include nausea, vomiting, diarrhea, dizziness, difficulty sleeping, and bad dreams. These are usually mild and do not cause people to stop taking the medicine.

Lariam may cause serious mental problems in some patients. (See “What is the most important information I should know about Lariam?”).

Lariam may affect your liver and your eyes if you take it for a long time. Your prescriber will tell you if you should have your eyes and liver checked while taking Lariam.

### **What else should I know about preventing malaria?**

- **Find out whether you need malaria prevention.** Before you travel, talk with your prescriber about your travel plans to determine whether you need to take medicine to prevent malaria. Even in those countries where malaria is present, there may be areas of the country that are free of malaria. In general, malaria is more common in rural (country) areas than in big cities, and it is more common during rainy seasons, when mosquitoes are most common. You can get information about the areas of the world where malaria occurs from the Centers for Disease Control and Prevention (CDC) and from local authorities in the countries you visit. If possible, plan your travel to reduce the risk of malaria.
- **Take medicine to prevent malaria infection.** Without malaria prevention medicine, you have a higher risk of getting malaria. Malaria starts with flu-like symptoms, such as chills, fever, muscle pains, and headaches. However, malaria can make you very sick or cause death if you don't seek medical help immediately. These symptoms may disappear for a while, and you may think you are well. But, the symptoms return later and then it may be too late for successful treatment.

Malaria can cause confusion, coma, and seizures. It can cause kidney failure, breathing problems, and severe damage to red blood cells. However, malaria can be easily diagnosed with a blood test, and if caught in time, can be effectively treated.

## **LARIAM® (mefloquine hydrochloride)**

**If you get flu-like symptoms (chills, fever, muscle pains, or headaches) after you return from a malaria area, get medical help right away and tell your prescriber that you may have been exposed to malaria.**

People who have lived for many years in areas with malaria may have some immunity to malaria (they do not get it as easily) and may not take malaria prevention medicine. This does not mean that you don't need to take malaria prevention medicine.

- **Protect against mosquito bites.** Medicines do not always completely prevent your catching malaria from mosquito bites. So protect yourself very well against mosquitoes. Cover your skin with long sleeves and long pants, and use mosquito repellent and bednets while in malaria areas. If you are out in the bush, you may want to pre-wash your clothes with permethrin. This is a mosquito repellent that may be effective for weeks after use. Ask your prescriber for other ways to protect yourself.

### **General information about the safe and effective use of Lariam.**

Medicines are sometimes prescribed for conditions not listed in Medication Guides. If you have any concerns about Lariam, ask your prescriber. This Medication Guide contains certain important information for travelers visiting areas with malaria. Your prescriber or pharmacist can give you information about Lariam that was written for health care professionals. Do not use Lariam for a condition for which it was not prescribed. Do not share Lariam with other people.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Manufactured by:  
F. HOFFMANN-LA ROCHE LTD  
Basel, Switzerland

Distributed by:



**Pharmaceuticals**

Roche Laboratories Inc.  
340 Kingsland Street  
Nutley, New Jersey 07110-1199

27898583

Revised: August 2003

Copyright © 2003 by Roche Laboratories Inc. All rights reserved.

## **LARIAM® (mefloquine hydrochloride)**

**Information wallet card to carry when you are taking Lariam.**



### **Lariam® (mefloquine hydrochloride) Tablets**

**You need to take malaria prevention medicine before you travel to a malaria area, while you are in a malaria area, and after you return from a malaria area.**

**If taken correctly, Lariam is effective at preventing malaria but, like all medications, it may produce side effects in some patients.**

**If you use Lariam to prevent malaria and you develop a sudden onset of anxiety, depression, restlessness, confusion (possible signs of more serious mental problems), or you develop other serious side effects, contact a doctor or other health care provider. It may be necessary to stop taking Lariam and use another malaria prevention medicine instead.**

Revised: August 2003

**Other medicines approved in the United States for malaria prevention include: doxycycline, atovaquone/proguanil, hydroxychloroquine, and chloroquine. Not all malaria medicines work equally well in malaria areas. The chloroquines, for example, do not work in many parts of the world. If you can't get another medicine, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.**

**Please read the Medication Guide for additional information on Lariam.**

**CENTER FOR DRUG EVALUATION AND  
RESEARCH**

*APPLICATION NUMBER:*

**NDA 019591/S-022**

**OTHER REVIEW(S)**

**Labeling, and Clinical Review of Supplemental Labeling Revision (SLR):**

**Sponsor:** Hoffman LaRoche, Inc.

**Product:** Lariam® (mefloquine hydrochloride) Tablets, 250 mg

**Materials Reviewed:**

<u>SLR</u>	<u>Date submitted</u>	<u>Date received</u>	<u>Date completed</u>
022	July 23, 2003	July 25, 2003	August 18 , 2003
<u>Amendment</u>			
022	August 15, 2003	August 15, 2003	August 18, 2003

- Approved package insert for NDA 19-591 dated January 10, 2003
- FDA approval letter for NDA 19-591/S-019 (Lariam Medication Guide) dated July 9, 2003

**Background:**

NDA 19-591 (Tablets) was originally approved on May 2, 1989. The last approved labeling change for the Lariam® package insert occurred on January 10, 2003 when Geriatric Use information was added. A new Lariam Medication Guide was approved by the Division on July 9, 2003.

Supplement-022 provides for the addition of:

1. The approved Lariam Medication Guide to the end of the Lariam package insert.
2. Information to the **PRECAUTIONS** section, **Information for Patients** subsection of the Lariam package insert concerning the Medication Guide.
3. Information concerning the new Medication Guide to the Lariam carton labeling.

In an e-mail message dated July 28, Dr. Gene Holbert, Chemistry Reviewer stated that the company's proposal to add the sentence "PHARMACIST: Provide a Medication Guide to each patient when dispensing Lariam." to the front of the Lariam carton was acceptable.

(b) (4)

The Division proposed the following alternative statement:

"Medication Guide: As required by law, a Lariam Medication Guide is supplied to patients when Lariam is dispensed. Patients should be instructed to read the MedGuide when Lariam is received. The complete text of the MedGuide is reprinted at the end of this document."

This proposal was sent to the company in an e-mail message dated July 29, 2003, and the company was asked to submit a revised Lariam package insert with this statement included.

On July 31, 2003 Dr. Sandra Kweder, Deputy OND Director notified the Division that following the Lariam medication guide approval on July 9, 2003, DDMAC proposed that several revisions should be made since they considered a few of the comments to be "promotional". Dr. Kweder also stated that ODS had proposed a change in the medication guide that the Division had not incorporated in the approved version but that she agreed with. Dr. Kweder requested that the Division inform the company that revisions would be necessary. The company was called on August 1, 2003, and was advised that additional changes for the medication guide would be forthcoming from the Division. The following changes were sent to the company via e-mail on August 4, 2003:

~~Strikethrough~~=deleted text

Double underline=added text

- If taken correctly, Lariam is ~~highly~~ effective at preventing malaria but, like all medications, it may produce side effects in some patients.
- ~~It has been reported that sometimes,~~ In some patients, these side effects continue after Lariam is stopped. Some patients taking Lariam think about killing themselves, and there have been rare reports of suicides. ~~We do not know if~~ It is not known whether Lariam was responsible for these suicides.
- **You need to take malaria prevention medicine before you travel to a malaria area, while you are in a malaria area, and after you return from a malaria area.**

Medicines approved in the United States for malaria prevention include Lariam, doxycycline, atovaquone/proguanil (Malarone), hydroxychloroquine, and chloroquine. Not all of these drugs work equally as well in all areas of the world where there is malaria. The chloroquines, for example, do not work in areas where the malaria parasite has developed resistance to chloroquine. Lariam may be effective against malaria that is resistant to chloroquine or other drugs. All drugs to treat malaria have side effects that are different for each one. For example, some may make your skin more sensitive to sunlight (Lariam does not do this). Lariam has the advantage of once weekly dosing, it can be used in children, it does not sensitize people to sunlight and it may be effective against malaria that is resistant to other drugs. However if you use Lariam to prevent malaria and you develop a sudden onset of anxiety, depression, restlessness, confusion (possible signs of more serious mental problems), or you develop other

serious side effects, contact a doctor or other health care provider, ~~as it~~ It may be necessary to stop taking Lariam and use another malaria prevention medicine instead. If you can't get another medicine, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.

- **Take Lariam exactly as prescribed. If you are an adult under 45 kg (99 pounds), your doctor will tell you the correct dose based on your weight.**
- For most adults, Take 1 tablet of Lariam at least 1 week before you travel to a malaria area (or 2 to 3 weeks before you travel to a malaria area, if instructed by your prescriber).

(b) (4)





On August 15, 2003 the company submitted a revised label for Lariam that included all of the FDA proposed and agreed-upon changes to the package insert and the medication guide.

**Electronic Labeling Comparison:**

The approved package insert for Lariam dated January 10, 2003 was electronically compared to the proposed package insert dated August 15, 2003. The changes were as follows:

~~Strikethrough~~=deleted text  
Double underline=added text

**1. PRECAUTIONS**

- In the **Information for Patients** subsection, the following sentences now appear first:

Medication Guide: As required by law, a Lariam Medication Guide is supplied to patients when Lariam is dispensed. Patients should be instructed to read the MedGuide when Lariam is received. The complete text of the Medication Guide is reprinted at the end of this document.

2. The Lariam Medication Guide approved on July 9, 2003 (with the additional agreed upon proposed labeling changes discussed with the company via e-mail on August 4-6, 2003 as follows) has been added to the end of the package insert:

**MEDICATION GUIDE**  
**LARIAM® (LAH-ree-am)**  
**(mefloquine hydrochloride) Tablets**  
**to Prevent Malaria**

**This Medication Guide is intended only for travelers who are taking Lariam to prevent malaria.** The information may not apply to patients who are sick with malaria and who are taking Lariam to treat malaria.

An information wallet card is provided at the end of this Medication Guide. Cut it out and carry it with you when you are taking Lariam.

This Medication Guide was ~~issued on [Month/Year]~~ revised in August 2003. Please read it before you start taking Lariam and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking with your prescriber (doctor or other health care provider) about Lariam and malaria prevention. Only you and your prescriber can decide if Lariam is right for you. If you cannot take Lariam, you may be able to take a different medicine to prevent malaria.

### ***What is the most important information I should know about Lariam?***

1. **Take Lariam exactly as prescribed to prevent malaria.**

Malaria is an infection that can cause death and is spread to humans through mosquito bites. If you travel to parts of the world where the mosquitoes carry the malaria parasite, you must take a malaria prevention medicine. Lariam is one of a small number of medications approved to prevent and to treat malaria. If taken correctly, Lariam is highly effective at preventing malaria but, like all medications, it may produce side effects in some patients.

2. **Lariam can rarely cause serious mental problems in some patients.**

The most frequently reported side effects with Lariam, such as nausea, difficulty sleeping, and bad dreams are usually mild and do not cause people to stop taking the medicine. However, people taking Lariam occasionally experience severe anxiety, feelings that people are against them, hallucinations (seeing or hearing things that are not there, for example), depression, unusual behavior, or feeling disoriented. ~~It has been reported that sometimes, in some patients,~~ There have been reports that in some patients these side effects continue after Lariam is stopped. Some patients taking Lariam think about killing themselves, and there have been rare reports of suicides. ~~We do not know if~~ It is not known whether Lariam was responsible for these suicides.

3. **You need to take malaria prevention medicine before you travel to a malaria area, while you are in a malaria area, and after you return from a malaria area.**

Medicines approved in the United States for malaria prevention include Lariam, doxycycline, atovaquone/proguanil, hydroxychloroquine, and chloroquine. Not all of these drugs work equally as well in all areas of the world where there is malaria. The chloroquines, for example, do not work in areas where the malaria parasite has developed resistance to chloroquine. Lariam may be effective against malaria that is resistant to chloroquine or other drugs. All drugs to treat malaria have side effects that are different for each one. For example, some may make your skin more sensitive to sunlight (Lariam does not do this). Lariam has the advantage of once weekly dosing, it can be used in children, it does not sensitize people to sunlight and it may be effective against malaria that is resistant to other drugs. However, if you use Lariam to prevent malaria and you develop a sudden onset of anxiety, depression, restlessness, confusion (possible signs of more serious mental problems), or you develop other serious side effects, contact a doctor or other health care provider, ~~as it~~ It may be necessary to stop taking Lariam and use another malaria prevention medicine instead. If you can't get another medicine, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.

### ***Who should not take Lariam?***

Do not take Lariam to **prevent** malaria if you

- **have depression or had depression recently**

- **have had recent mental illness or problems**, including anxiety disorder, schizophrenia (a severe type of mental illness), or psychosis (losing touch with reality)
- **have or had seizures (epilepsy or convulsions)**
- **are allergic to quinine or quinidine (medicines related to Lariam)**

**Tell your prescriber about all your medical conditions.** Lariam may not be right for you if you have certain conditions, especially the ones listed below:

- **Heart disease.** Lariam may not be right for you.
- **Pregnancy.** Tell your prescriber if you are pregnant or plan to become pregnant. It is dangerous for the mother and for the unborn baby (fetus) to get malaria during pregnancy. Therefore, ask your prescriber if you should take Lariam or another medicine to prevent malaria while you are pregnant.
- **Breast feeding.** Lariam can pass through your milk and may harm the baby. Therefore, ask your prescriber whether you will need to stop breast feeding or use another medicine.
- **Liver problems.**

**Tell your prescriber about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements.** Some medicines may give you a higher chance of having serious side effects from Lariam.

### ***How should I take Lariam?***

**Take Lariam exactly as prescribed. If you are an adult or pediatric patient weighing 45 kg (99 pounds) or less, your prescriber will tell you the correct dose based on your weight.**

To prevent malaria

- For adults and pediatric patients weighing over 45 kg, Take 1 tablet of Lariam at least 1 week before you travel to a malaria area (or 2 to 3 weeks before you travel to a malaria area, if instructed by your prescriber). This starts the prevention and also helps you see how Lariam affects you and the other medicines you take. **Take 1 Lariam tablet once a week**, on the same day each week, while in a malaria area.
- **Continue taking Lariam for 4 weeks after returning from a malaria area.** If you cannot continue taking Lariam due to side effects or for other reasons, contact your prescriber.
- Take Lariam just after a meal and with at least 1 cup (8 ounces) of water.

- For children, Lariam can be given with water or crushed and mixed with water or sugar water. The prescriber will tell you the correct dose for children based on the child's weight.
- If you are told by a doctor or other health care provider to stop taking Lariam due to side effects or for other reasons, it will be necessary to take another malaria medicine. You must take **malaria prevention medicine before you travel to a malaria area, while you are in a malaria area, and after you return from a malaria area. If you don't have access to a doctor or other health care provider or to another medicine besides Lariam and have to stop taking it, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.**

### ***What should I avoid while taking Lariam?***

- **Halofantrine (marketed under various brand names)**, a medicine used to treat malaria. Taking both of these medicines together can cause serious heart problems that can cause death.
- **Do not become pregnant.** Women should use effective birth control while taking Lariam.
- **Quinine, quinidine, or chloroquine (other medicines used to treat malaria).** Taking these medicines with Lariam could cause changes in your heart rate or increase the risk of seizures.

### **In addition:**

- **Be careful driving or in other activities** needing alertness and careful movements (fine motor coordination). Lariam can cause dizziness or loss of balance, even after you stop taking it.
- **Be aware that certain vaccines may not work if given while you are taking Lariam.** Your prescriber may want you to finish taking your vaccines at least 3 days before starting Lariam.

### ***What are the possible side effects of Lariam?***

Lariam, like all medicines, may cause side effects in some patients. The most frequently reported side effects with Lariam when used for prevention of malaria include nausea, vomiting, diarrhea, dizziness, difficulty sleeping, and bad dreams. These are usually mild and do not cause people to stop taking the medicine.

Lariam may cause serious mental problems in some patients. (See "What is the most important information I should know about Lariam?").

Lariam may affect your liver and your eyes if you take it for a long time. Your prescriber will tell you if you should have your eyes and liver checked while taking Lariam.

***What else should I know about preventing malaria?***

- **Find out whether you need malaria prevention.** Before you travel, talk with your prescriber about your travel plans to determine whether you need to take medicine to prevent malaria. Even in those countries where malaria is present, there may be areas of the country that are free of malaria. In general, malaria is more common in rural (country) areas than in big cities, and it is more common during rainy seasons, when mosquitoes are most common. You can get information about the areas of the world where malaria occurs from the Centers for Disease Control and Prevention (CDC) and from local authorities in the countries you visit. If possible, plan your travel to reduce the risk of malaria.
- **Take medicine to prevent malaria infection.** Without malaria prevention medicine, you have a higher risk of getting malaria. Malaria starts with flu-like symptoms, such as chills, fever, muscle pains, and headaches. However, malaria can make you very sick or cause death if you don't seek medical help immediately. These symptoms may disappear for a while, and you may think you are well. But, the symptoms return later and then it may be too late for successful treatment.

Malaria can cause confusion, coma, and seizures. It can cause kidney failure, breathing problems, and severe damage to red blood cells. However, malaria can be easily diagnosed with a blood test, and if caught in time, can be effectively treated.

**If you get flu-like symptoms (chills, fever, muscle pains, or headaches) after you return from a malaria area,** get medical help right away and tell your prescriber that you may have been exposed to malaria.

People who have lived for many years in areas with malaria may have some immunity to malaria (they do not get it as easily) and may not take malaria prevention medicine. This does not mean that you don't need to take malaria prevention medicine.

- **Protect against mosquito bites.** Medicines do not always completely prevent your catching malaria from mosquito bites. So protect yourself very well against mosquitoes. Cover your skin with long sleeves and long pants, and use mosquito repellent and bednets while in malaria areas. If you are out in the bush, you may want to pre-wash your clothes with permethrin. This is a mosquito repellent that may be effective for weeks after use. Ask your prescriber for other ways to protect yourself.

***General information about the safe and effective use of Lariam.***

Medicines are sometimes prescribed for conditions not listed in Medication Guides. If you have any concerns about Lariam, ask your prescriber. This Medication Guide contains certain important information for travelers visiting areas with malaria. Your prescriber or pharmacist can give you information about Lariam that was written for health care professionals. Do not use Lariam for a condition for which it was not prescribed. Do not share Lariam with other people.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Manufactured by:  
F. HOFFMANN-LA ROCHE LTD  
Basel, Switzerland

Distributed by:



**Pharmaceuticals**

Roche Laboratories Inc.  
340 Kingsland Street  
Nutley, New Jersey 07110-1199

27898583

Revised: August 2003

Copyright © 2003 by Roche Laboratories Inc. All rights reserved.

**Information wallet card to carry when you are taking Lariam.**



<b>Lariam® (mefloquine hydrochloride) Tablets</b>	
<p><b>You need to take malaria prevention medicine before you travel to a malaria area, while you are in a malaria area, and after you return from a malaria area.</b></p> <p><b>If taken correctly, Lariam is highly effective at preventing malaria but, like all medications, it may produce side effects in some patients.</b></p> <p><b>If you use Lariam to prevent malaria and you develop a sudden onset of anxiety, depression, restlessness, confusion (possible signs of more serious mental problems), or you develop other serious side effects, contact a doctor or other health care provider as it. <u>It</u> may be necessary to stop taking Lariam and use another malaria prevention medicine instead.</b></p> <p>Issued: Month Year</p> <p><u>Revised: August 2003</u></p>	<p><b>Other medicines approved in the United States for malaria prevention include: doxycycline, atovaquone/proguanil (<del>Malarone</del>), hydroxychloroquine, and chloroquine. Not all malaria medicines work equally well in malaria areas. The chloroquines, for example, do not work in many parts of the world. If you can't get another medicine, leave the malaria area. However, be aware that leaving the malaria area may not protect you from getting malaria. You still need to take a malaria prevention medicine.</b></p> <p><b>Please read the Medication Guide for additional information on Lariam.</b></p>

The carton labeling now includes the following statement (b) (4) located next to the dosage on the front of the carton:

PHARMACIST: Provide a Medication Guide to each patient when dispensing Lariam.

**Conclusions/Recommendations:**

These labeling changes are acceptable. An approval letter should be sent advising the applicant that this supplemental NDA submission is approved.

Robin Anderson, R.N., M.B.A.  
Labeling Reviewer

Leonard Sacks, M.B.B.Ch  
Medical Officer

cc:

HFD-590/MO/L. Sacks  
HFD-590/MedTL/R. Roca  
HFD-590/DivDir/S. Gitterman for R. Albrecht  
  
HFD-590/PM/J. Saliba

Concurrence:

HFD-590/MO/L. Sacks 8/20/03  
HFD-590/MedTL/R. Roca 8/20/03  
HFD-590/DivDir/S. Gitterman for R.  
Albrecht



-----  
**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**  
-----

/s/

-----  
Robin Anderson  
8/20/03 05:42:04 PM  
INTERDISCIPLINARY

Please sign for Dr. Renata Albrecht.

Steven Gitterman  
8/20/03 10:22:00 PM  
MEDICAL OFFICER  
for Renata Albrecht