

CENTER FOR DRUG EVALUATION AND RESEARCH

Approval Package for:

APPLICATION NUMBER:

020687Orig1s025

Trade Name: Mifeprex Tablets 200 mg

Generic or Proper Name: Mifepristone

Sponsor: Danco Laboratories, LLC

Approval Date: January 3, 2023

Indication: For modification to the approved single, shared system (SSS) risk evaluation and mitigation strategy (REMS) for mifepristone 200 mg tablets, in a regimen with misoprostol, for the medical termination of intrauterine pregnancy through 70 days gestation, as well as corresponding labeling revisions to the prescribing information and the Medication Guide to align with the modification to the Mifepristone REMS Program.

CENTER FOR DRUG EVALUATION AND RESEARCH

020687Orig1s025

CONTENTS

Reviews / Information Included in this NDA Review.

Approval Letter	X
Other Action Letters	
Labeling	X
REMS	X
Summary Review	X
Officer/Employee List	
Office Director Memo	
Cross Discipline Team Leader Review	
Clinical Review(s)	
Product Quality Review(s)	
Non-Clinical Review(s)	
Statistical Review(s)	
Clinical Microbiology / Virology Review(s)	
Clinical Pharmacology Review(s)	
Other Reviews	X
Risk Assessment and Risk Mitigation Review(s)	X
Proprietary Name Review(s)	
Administrative/Correspondence Document(s)	X

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

020687Orig1s025

APPROVAL LETTER



NDA 020687/S-025

SUPPLEMENT APPROVAL

Danco Laboratories, LLC

(b) (4), (b) (6)

P.O. Box 4816
New York, NY 10185

Dear (b) (4), (b) (6) :

Please refer to your supplemental new drug application (sNDA) dated and received June 22, 2022, and your amendments, submitted under section 505(b) of the Federal Food, Drug, and Cosmetic Act (FDCA) for Mifeprex (mifepristone) Tablets, 200 mg.

This Prior Approval sNDA provides for modification to the approved single, shared system (SSS) risk evaluation and mitigation strategy (REMS) for mifepristone 200 mg tablets, in a regimen with misoprostol, for the medical termination of intrauterine pregnancy through 70 days gestation, as well as corresponding labeling revisions to the prescribing information and the Medication Guide to align with the modification to the Mifepristone REMS Program. This SSS REMS is known as the Mifepristone REMS Program.

APPROVAL & LABELING

We have completed our review of the supplemental application, as amended. It is approved effective the date of this letter.

CONTENT OF LABELING

As soon as possible, but no later than 14 days from the date of this letter, submit the content of labeling [21 CFR 314.50(l)] in structured product labeling (SPL) format using the FDA automated drug registration and listing system (eLIST), as described at FDA.gov.¹ Content of labeling must be identical to the enclosed labeling (text for the Prescribing Information and Medication Guide), with the addition of any labeling changes in pending "Changes Being Effectuated" (CBE) supplements, as well as annual reportable changes not included in the enclosed labeling.

¹ <http://www.fda.gov/ForIndustry/DataStandards/StructuredProductLabeling/default.htm>

Information on submitting SPL files using eList may be found in the guidance for industry *SPL Standard for Content of Labeling Technical Qs and As*.²

The SPL will be accessible from publicly available labeling repositories.

Also within 14 days, amend all pending supplemental applications that include labeling changes for this NDA, including CBE supplements for which FDA has not yet issued an action letter, with the content of labeling [21 CFR 314.50(l)(1)(i)] in Microsoft Word format, that includes the changes approved in this supplemental application, as well as annual reportable changes. To facilitate review of your submission(s), provide a highlighted or marked-up copy that shows all changes, as well as a clean Microsoft Word version. The marked-up copy should provide appropriate annotations, including supplement number(s) and annual report date(s).

We request that the labeling approved today be available on your website within 10 days of receipt of this letter.

REQUIRED PEDIATRIC ASSESSMENTS

Under the Pediatric Research Equity Act (PREA) (21 U.S.C. 355c), all applications for new active ingredients (which includes new salts and new fixed combinations), new indications, new dosage forms, new dosing regimens, or new routes of administration are required to contain an assessment of the safety and effectiveness of the product for the claimed indication in pediatric patients unless this requirement is waived, deferred, or inapplicable.

Because none of these criteria apply to your application, you are exempt from this requirement.

RISK EVALUATION AND MITIGATION STRATEGY (REMS) REQUIREMENTS

The Mifepristone REMS Program, of which Mifeprex is a member, was originally approved on April 11, 2019, and the most recent REMS modification was approved on May 14, 2021. The Mifepristone REMS Program consists of elements to assure safe use, an implementation system, and a timetable for submission of assessments of the REMS.

In order to ensure the benefits of Mifeprex outweigh its risks and to minimize burden on the healthcare delivery system of complying with the REMS, we determined that you were required to make the REMS modifications outlined in our REMS Modification

² We update guidances periodically. For the most recent version of a guidance, check the FDA Guidance Documents Database <https://www.fda.gov/RegulatoryInformation/Guidances/default.htm>.

Notification letter dated December 16, 2021. In addition the following modifications were communicated during the course of the review:

- Revisions to the REMS goal to align with the updated REMS requirements.
- Replacing serial number with recording of NDC and lot number of mifepristone dispensed.
- Additional edits for clarification and consistency in the REMS Document and REMS materials (*Prescriber Agreement Forms*, *Patient Agreement Form*, and *Pharmacy Agreement Forms*).

Your proposed modified REMS, received on June 22, 2022, amended and appended to this letter, is approved. The modified REMS consists of the elements to assure safe use, implementation system, and a timetable for submission of assessments of the REMS.

The modification of the approved REMS must be fully implemented within 120 calendar days of this letter.

This shared system REMS, known as the Mifepristone REMS Program, currently includes those products listed on the FDA REMS website³.

Other products may be added in the future if additional NDAs or ANDAs are approved.

The timetable for submission of assessments of the REMS must be revised to one year from the date of the approval of the modified SSS REMS (1/3/2023) and annually thereafter.

The revised REMS assessment plan must include, but is not limited to, the following:

Program Implementation and Operations

1. REMS Certification Statistics

a. Prescribers

- i. Number of certified prescribers who have certified with the Sponsor's distributor(s) and number who have submitted *Prescriber Agreement Forms* to Certified Pharmacies
- ii. Number and percentage of newly certified prescribers
- iii. Number and percentage of active certified prescribers (i.e., who ordered mifepristone or submitted a prescription during the reporting period)

b. Pharmacies

³ <https://www.accessdata.fda.gov/scripts/cder/remis/index.cfm>

- i. Number of certified pharmacies
 - ii. Number and percentage of newly certified pharmacies
 - iii. Number and percentage of active certified pharmacies (i.e., that dispensed mifepristone during the reporting period)
 - c. Wholesalers/Distributors
 - i. Number of authorized wholesalers/distributors
 - ii. Number and percentage of newly authorized wholesalers/distributors
 - iii. Number and percentage of active authorized wholesalers/distributors (i.e. that shipped mifepristone during the reporting period)
2. Utilization Data
- a. Total number of tablets shipped by wholesalers/distributors, stratified by Certified Prescriber or Certified Pharmacy location
 - b. Number of prescriptions dispensed from pharmacies
3. REMS Compliance Data
- a. Audits: Summary of audit activities for each stakeholder (i.e., certified pharmacies and wholesalers/distributors) including but not limited to:
 - i. A copy of the final audit plan for each stakeholder type (provide for the current reporting period)
 - ii. The number of audits expected, and the number of audits performed
 - iii. The number and type of deficiencies noted
 - iv. For those with deficiencies noted, report the corrective and preventive actions (CAPAs) required, if any, to address the deficiencies, including the status (e.g., completed, not completed, in progress) (provide for the current reporting period)
 - v. For any stakeholders that did not complete the CAPA within the timeframe specified in the audit plan, describe actions taken (provide for the current reporting period)
 - vi. A summary report of all resulting changes to processes and procedures necessary to ensure compliance with the REMS requirements (provide for the current reporting period)
 - b. A summary report of non-compliance, associated corrective action plans (CAPAs), and the status of CAPAs including but not limited to:
 - i. A copy of the final non-compliance plans for Pharmacies and Distributors (provide for the current reporting period)
 - ii. For each instance of noncompliance below (iii-v), report the following information (provide for the current reporting period):
 - 1. A unique, anonymized ID for the stakeholder(s) associated with the non-compliance event to enable tracking over time
 - 2. The source of the non-compliance data (e.g., self-reported, audit, other)
 - 3. A root cause analysis of the non-compliance

4. Actions to prevent future occurrences and outcomes of such actions
- iii. Prescriber compliance
 1. Number and percentage of certified prescribers who became decertified as a result of non-compliance
 - Provide a summary of reasons for decertification (provide for the current reporting period)
 2. Summary and analysis of any program deviations and corrective actions taken (provide for the current reporting period)
- iv. Pharmacy compliance
 1. Number and percentage of prescriptions dispensed that were written by prescriber(s) who did not submit a Prescriber Agreement to the dispensing Certified Pharmacy
 2. Number and percentage of mifepristone tablets dispensed by non-certified pharmacies
 3. Number and percentage of pharmacies that became decertified as a result of non-compliance
 - Provide a summary of reasons for decertification (provide for the current reporting period)
 4. An assessment of prescription delivery timelines, including percentage delivered more than four days after receipt of the prescription, duration and causes for delay. A proposal for this assessment will be submitted within 60 days of the approval of the REMS Modification.
 5. Summary and analysis of any program deviations and corrective actions taken (provide for the current reporting period)
- v. Wholesaler/distributor compliance
 1. Number of healthcare providers who successfully ordered mifepristone who were not certified
 2. Number of non-certified pharmacies that successfully ordered mifepristone
 3. Number of shipments sent to non-certified prescriber receiving locations
 4. Number of shipments sent to non-certified pharmacy receiving locations
 5. Summary and analysis of any program deviations and corrective actions taken (provide for the current reporting period)

Overall Assessment of REMS Effectiveness

The requirements for assessments of an approved REMS under section 505-1(g)(3) include with respect to each goal included in the strategy, an assessment of the extent to which the approved strategy, including each element of the strategy, is meeting the goal or whether one or more such goals or such elements should be modified.

We remind you that in addition to the REMS assessments submitted according to the timetable in the approved REMS, you must include an adequate rationale to support a

proposed REMS modification for the addition, modification, or removal of any goal or element of the REMS, as described in section 505-1(g)(4) of the FDCA.

We also remind you that you must submit a REMS assessment when you submit a supplemental application for a new indication for use, as described in section 505-1(g)(2)(A) of the FDCA. This assessment should include:

- a) An evaluation of how the benefit-risk profile will or will not change with the new indication;
- b) A determination of the implications of a change in the benefit-risk profile for the current REMS;
- c) *If the new indication for use introduces unexpected risks:* A description of those risks and an evaluation of whether those risks can be appropriately managed with the currently approved REMS.
- d) *If a REMS assessment was submitted in the 18 months prior to submission of the supplemental application for a new indication for use:* A statement about whether the REMS was meeting its goals at the time of that last assessment and if any modifications of the REMS have been proposed since that assessment.
- e) *If a REMS assessment has not been submitted in the 18 months prior to submission of the supplemental application for a new indication for use:* Provision of as many of the currently listed assessment plan items as is feasible.
- f) *If you propose a REMS modification based on a change in the benefit-risk profile or because of the new indication of use, submit an adequate rationale to support the modification, including:* Provision of the reason(s) why the proposed REMS modification is necessary, the potential effect on the serious risk(s) for which the REMS was required, on patient access to the drug, and/or on the burden on the health care delivery system; and other appropriate evidence or data to support the proposed change. Additionally, include any changes to the assessment plan necessary to assess the proposed modified REMS. *If you are not proposing REMS modifications,* provide a rationale for why the REMS does not need to be modified.

If the assessment instruments and methodology for your REMS assessments are not included in the REMS supporting document, or if you propose changes to the submitted assessment instruments or methodology, you should update the REMS supporting document to include specific assessment instrument and methodology information at least 90 days before the assessments will be conducted. Updates to the REMS supporting document may be included in a new document that references previous REMS supporting document submission(s) for unchanged portions. Alternatively,

U.S. Food and Drug Administration
Silver Spring, MD 20993
www.fda.gov

updates may be made by modifying the complete previous REMS supporting document, with all changes marked and highlighted.

Prominently identify the submission containing the assessment instruments and methodology with the following wording in bold capital letters at the top of the first page of the submission:

NDA 020687 REMS ASSESSMENT METHODOLOGY

(insert concise description of content in bold capital letters, e.g.,

ASSESSMENT METHODOLOGY, PROTOCOL, SURVEY METHODOLOGIES, AUDIT PLAN, DRUG USE STUDY)

An authorized generic drug under this NDA must have an approved REMS prior to marketing. Should you decide to market, sell, or distribute an authorized generic drug under this NDA, contact us to discuss what will be required in the authorized generic drug REMS submission.

We remind you that section 505-1(f)(8) of FDCA prohibits holders of an approved covered application with elements to assure safe use from using any element to block or delay approval of an application under section 505(b)(2) or (j). A violation of this provision in 505-1(f) could result in enforcement action.

Prominently identify any submission containing the REMS assessments or proposed modifications of the REMS with the following wording in bold capital letters at the top of the first page of the submission as appropriate:

NDA 020687 REMS ASSESSMENT

or

**NEW SUPPLEMENT FOR NDA 020687/S-000
CHANGES BEING EFFECTED IN 30 DAYS
PROPOSED MINOR REMS MODIFICATION**

or

**NEW SUPPLEMENT FOR NDA 020687/S-000
PRIOR APPROVAL SUPPLEMENT
PROPOSED MAJOR REMS MODIFICATION**

or

**NEW SUPPLEMENT FOR NDA 020687/S-000/
PRIOR APPROVAL SUPPLEMENT**

**PROPOSED REMS MODIFICATIONS DUE TO SAFETY LABELING
CHANGES SUBMITTED IN SUPPLEMENT XXX**

or

**NEW SUPPLEMENT (NEW INDICATION FOR USE)
FOR NDA 020687/S-000
REMS ASSESSMENT
PROPOSED REMS MODIFICATION (if included)**

Should you choose to submit a REMS revision, prominently identify the submission containing the REMS revisions with the following wording in bold capital letters at the top of the first page of the submission:

REMS REVISIONS FOR NDA 020687

To facilitate review of your submission, we request that you submit your proposed modified REMS and other REMS-related materials in Microsoft Word format. If certain documents, such as enrollment forms, or website screenshots are only in PDF format, they may be submitted as such, but Word format is preferred.

SUBMISSION OF REMS DOCUMENT IN SPL FORMAT

FDA can accept the REMS document in Structured Product Labeling (SPL) format. If you intend to submit the REMS document in SPL format, as soon as possible, but no later than 14 days from the date of this letter, submit the REMS document in SPL format using the FDA automated drug registration and listing system (eLIST).

For more information on submitting REMS in SPL format, please email FDAREMSwebsite@fda.hhs.gov

PATENT LISTING REQUIREMENTS

Pursuant to 21 CFR 314.53(d)(2) and 314.70(f), certain changes to an approved NDA submitted in a supplement require you to submit patent information for listing in the Orange Book upon approval of the supplement. You must submit the patent information required by 21 CFR 314.53(d)(2)(i)(A) through (C) and 314.53(d)(2)(ii)(A) and (C), as applicable, to FDA on Form FDA 3542 within 30 days after the date of approval of the supplement for the patent information to be timely filed (see 21 CFR 314.53(c)(2)(ii)). You also must ensure that any changes to your approved NDA that require the submission of a request to remove patent information from the Orange Book are submitted to FDA at the time of approval of the supplement pursuant to 21 CFR 314.53(d)(2)(ii)(B) and 314.53(f)(2)(iv).

REPORTING REQUIREMENTS

We remind you that you must comply with reporting requirements for an approved NDA (21 CFR 314.80 and 314.81).

If you have any questions, call  (b) (6)

Sincerely,

{See appended electronic signature page}

 (b) (6)

Center for Drug Evaluation and Research

ENCLOSURE(S):

- Content of Labeling
 - o Prescribing Information
 - o Medication Guide
 - o REMS Document
 - o Prescriber Agreement
 - o Patient Agreement Form
 - o Pharmacy Agreement Form

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

(b) (6)

01/03/2023 05:35:41 PM

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

020687Orig1s025

LABELING

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use MIFEPREX safely and effectively. See full prescribing information for MIFEPREX.

MIFEPREX® (mifepristone) tablets, for oral use
Initial U.S. Approval: 2000

WARNING: SERIOUS AND SOMETIMES FATAL INFECTIONS OR BLEEDING

See full prescribing information for complete boxed warning. Serious and sometimes fatal infections and bleeding occur very rarely following spontaneous, surgical, and medical abortions, including following MIFEPREX use.

- Atypical Presentation of Infection. Patients with serious bacterial infections and sepsis can present without fever, bacteremia or significant findings on pelvic examination. A high index of suspicion is needed to rule out serious infection and sepsis. (5.1)
- Bleeding. Prolonged heavy bleeding may be a sign of incomplete abortion or other complications and prompt medical or surgical intervention may be needed. (5.2)

MIFEPREX is only available through a restricted program called the mifepristone REMS Program (5.3).

Before prescribing MIFEPREX, inform the patient about these risks. Ensure the patient knows whom to call and what to do if they experience sustained fever, severe abdominal pain, prolonged heavy bleeding, or syncope, or if they experience abdominal pain or discomfort or general malaise for more than 24 hours after taking misoprostol.

INDICATIONS AND USAGE

MIFEPREX is a progestin antagonist indicated, in a regimen with misoprostol, for the medical termination of intrauterine pregnancy through 70 days gestation. (1)

DOSAGE AND ADMINISTRATION

- 200 mg MIFEPREX on Day 1, followed 24-48 hours after MIFEPREX dosing by 800 mcg buccal misoprostol. (2.1)
- Instruct the patient what to do if significant adverse reactions occur. (2.2)
- Follow-up is needed to confirm complete termination of pregnancy. (2.3)

DOSAGE FORMS AND STRENGTHS

Tablets containing 200 mg of mifepristone each, supplied as 1 tablet on one blister card (3)

CONTRAINDICATIONS

- Confirmed/suspected ectopic pregnancy or undiagnosed adnexal mass (4)
- Chronic adrenal failure (4)
- Concurrent long-term corticosteroid therapy (4)
- History of allergy to mifepristone, misoprostol, or other prostaglandins (4)
- Hemorrhagic disorders or concurrent anticoagulant therapy (4)
- Inherited porphyria (4)
- Intrauterine device (IUD) in place (4)

WARNINGS AND PRECAUTIONS

- Ectopic pregnancy: Exclude before treatment. (5.4)
- Rhesus immunization: Prevention needed as for surgical abortion. (5.5)

ADVERSE REACTIONS

Most common adverse reactions (>15%) are nausea, weakness, fever/chills, vomiting, headache, diarrhea, and dizziness. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Danco Laboratories, LLC at 1-877-432-7596 or medicaldirector@earlyoptionpill.com or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- CYP3A4 inducers can lower mifepristone concentrations. (7.1)
- CYP3A4 inhibitors can increase mifepristone concentrations. Use with caution. (7.2)
- CYP3A4 substrate concentrations can be increased. Caution with coadministration of substrates with narrow therapeutic margin. (7.3)

USE IN SPECIFIC POPULATIONS

- Pregnancy: Risk of fetal malformations in ongoing pregnancy if not terminated is unknown. (8.1)

See 17 for PATIENT COUNSELING INFORMATION, Medication Guide.

Revised: 01/2023

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: SERIOUS AND SOMETIMES FATAL INFECTIONS OR BLEEDING

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosing Regimen
- 2.2 Patient Management Following Misoprostol Administration
- 2.3 Post-treatment Assessment: Day 7 to 14
- 2.4 Contact for Consultation

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Infections and Sepsis
- 5.2 Uterine Bleeding
- 5.3 Mifepristone REMS Program
- 5.4 Ectopic Pregnancy
- 5.5 Rhesus Immunization

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Drugs that May Reduce MIFEPREX Exposure (Effect of CYP 3A4 Inducers on MIFEPREX)

- 7.2 Drugs that May Increase Exposure (Effect of CYP 3A4 Inhibitors on MIFEPREX)

- 7.3 Effects of MIFEPREX on Other Drugs (Effect of MIFEPREX on CYP 3A4 Substrates)

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: SERIOUS AND SOMETIMES FATAL INFECTIONS OR BLEEDING

Serious and sometimes fatal infections and bleeding occur very rarely following spontaneous, surgical, and medical abortions, including following MIFEPREX use. No causal relationship between the use of MIFEPREX and misoprostol and these events has been established.

- **Atypical Presentation of Infection.** Patients with serious bacterial infections (e.g., *Clostridium sordellii*) and sepsis can present without fever, bacteremia, or significant findings on pelvic examination following an abortion. Very rarely, deaths have been reported in patients who presented without fever, with or without abdominal pain, but with leukocytosis with a marked left shift, tachycardia, hemoconcentration, and general malaise. A high index of suspicion is needed to rule out serious infection and sepsis [see *Warnings and Precautions (5.1)*].
- **Bleeding.** Prolonged heavy bleeding may be a sign of incomplete abortion or other complications and prompt medical or surgical intervention may be needed. Advise patients to seek immediate medical attention if they experience prolonged heavy vaginal bleeding [see *Warnings and Precautions (5.2)*].

Because of the risks of serious complications described above, MIFEPREX is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called the mifepristone REMS Program [see *Warnings and Precautions (5.3)*].

Before prescribing MIFEPREX, inform the patient about the risk of these serious events. Ensure that the patient knows whom to call and what to do, including going to an Emergency Room if none of the provided contacts are reachable, if they experience sustained fever, severe abdominal pain, prolonged heavy bleeding, or syncope, or if they experience abdominal pain or discomfort, or general malaise (including weakness, nausea, vomiting, or diarrhea) for more than 24 hours after taking misoprostol.

1 INDICATIONS AND USAGE

MIFEPREX is indicated, in a regimen with misoprostol, for the medical termination of intrauterine pregnancy through 70 days gestation.

2 DOSAGE AND ADMINISTRATION

2.1 Dosing Regimen

For purposes of this treatment, pregnancy is dated from the first day of the last menstrual period. The duration of pregnancy may be determined from menstrual history and clinical examination. Assess the pregnancy by ultrasonographic scan if the duration of pregnancy is uncertain or if ectopic pregnancy is suspected.


Remove any intrauterine device ("IUD") before treatment with MIFEPREX begins [see *Contraindications (4)*].

The dosing regimen for MIFEPREX and misoprostol is:

- MIFEPREX 200 mg orally + misoprostol 800 mcg buccally
 - *Day One:* MIFEPREX Administration
One 200 mg tablet of MIFEPREX is taken in a single oral dose.
 - *Day Two or Three:* Misoprostol Administration (minimum 24-hour interval between MIFEPREX and misoprostol)
Four 200 mcg tablets (total dose 800 mcg) of misoprostol are taken by the buccal route.

Tell the patient to place two 200 mcg misoprostol tablets in each cheek pouch (the area between the cheek and gums) for 30 minutes and then swallow any remnants with water or another liquid (see Figure 1).

Figure 1



2 pills between cheek and gum on left side + 2 pills between cheek and gum on right side

Patients taking MIFEPREX must take misoprostol within 24 to 48 hours after taking MIFEPREX. The effectiveness of the regimen may be lower if misoprostol is administered less than 24 hours or more than 48 hours after mifepristone administration.

Because most women will expel the pregnancy within 2 to 24 hours of taking misoprostol [see *Clinical Studies (14)*], discuss with the patient an appropriate location for them to be when taking the misoprostol, taking into account that expulsion could begin within 2 hours of administration.

2.2 Patient Management Following Misoprostol Administration

During the period immediately following the administration of misoprostol, the patient may need medication for cramps or gastrointestinal symptoms [see *Adverse Reactions (6)*].

Give the patient:

- Instructions on what to do if significant discomfort, excessive vaginal bleeding or other adverse reactions occur
- A phone number to call if the patient has questions following the administration of the misoprostol
- The name and phone number of the healthcare provider who will be handling emergencies.

2.3 Post-treatment Assessment: Day 7 to 14

Patients should follow-up with their healthcare provider approximately 7 to 14 days after the administration of MIFEPREX. This assessment is very important to confirm that complete termination of pregnancy has occurred and to evaluate the degree of bleeding. Termination can be confirmed by medical history, clinical examination, human Chorionic Gonadotropin (hCG) testing, or ultrasonographic scan. Lack of bleeding following treatment usually indicates failure; however, prolonged or heavy bleeding is not proof of a complete abortion.

The existence of debris in the uterus (e.g., if seen on ultrasonography) following the treatment procedure will not necessarily require surgery for its removal.

Patients should expect to experience vaginal bleeding or spotting for an average of 9 to 16 days. Women report experiencing heavy bleeding for a median duration of 2 days. Up to 8% of women may experience some type of bleeding for more than 30 days. Persistence of heavy or moderate vaginal bleeding at the time of follow-up, however, could indicate an incomplete abortion.

If complete expulsion has not occurred, but the pregnancy is not ongoing, patients may be treated with another dose of misoprostol 800 mcg buccally. There have been rare reports of uterine rupture in women who took MIFEPREX and misoprostol, including women with prior uterine rupture or uterine scar and women who received multiple doses of misoprostol within 24 hours. Patients who choose to use a repeat dose of misoprostol should have a follow-up visit with their healthcare provider in approximately 7 days to assess for complete termination.

Surgical evacuation is recommended to manage ongoing pregnancies after medical abortion [see *Use in Specific Populations (8.1)*]. Advise the patient whether you will provide such care or will refer them to another provider as part of counseling prior to prescribing MIFEPREX.

2.4 Contact for Consultation

For consultation 24 hours a day, 7 days a week with an expert in mifepristone, call Danco Laboratories at 1-877-4 Early Option (1-877-432-7596).

3 DOSAGE FORMS AND STRENGTHS

Tablets containing 200 mg of mifepristone each, supplied as 1 tablet on one blister card. MIFEPREX tablets are light yellow, cylindrical, and bi-convex tablets, approximately 11 mm in diameter and imprinted on one side with "MF."

4 CONTRAINDICATIONS

- Administration of MIFEPREX and misoprostol for the termination of pregnancy (the "treatment procedure") is contraindicated in patients with any of the following conditions:
 - Confirmed or suspected ectopic pregnancy or undiagnosed adnexal mass (the treatment procedure will not be effective to terminate an ectopic pregnancy) [see *Warnings and Precautions (5.4)*]
 - Chronic adrenal failure (risk of acute adrenal insufficiency)
 - Concurrent long-term corticosteroid therapy (risk of acute adrenal insufficiency)
 - History of allergy to mifepristone, misoprostol, or other prostaglandins (allergic reactions including anaphylaxis, angioedema, rash, hives, and itching have been reported [see *Adverse Reactions (6.2)*])
 - Hemorrhagic disorders or concurrent anticoagulant therapy (risk of heavy bleeding)

- Inherited porphyrias (risk of worsening or of precipitation of attacks)
- Use of MIFEPREX and misoprostol for termination of intrauterine pregnancy is contraindicated in patients with an intrauterine device (“IUD”) in place (the IUD might interfere with pregnancy termination). If the IUD is removed, MIFEPREX may be used.

5 WARNINGS AND PRECAUTIONS

5.1 Infection and Sepsis

As with other types of abortion, cases of serious bacterial infection, including very rare cases of fatal septic shock, have been reported following the use of MIFEPREX [see *Boxed Warning*]. Healthcare providers evaluating a patient who is undergoing a medical abortion should be alert to the possibility of this rare event. A sustained (> 4 hours) fever of 100.4°F or higher, severe abdominal pain, or pelvic tenderness in the days after a medical abortion may be an indication of infection.

A high index of suspicion is needed to rule out sepsis (e.g., from *Clostridium sordellii*) if a patient reports abdominal pain or discomfort or general malaise (including weakness, nausea, vomiting, or diarrhea) more than 24 hours after taking misoprostol. Very rarely, deaths have been reported in patients who presented without fever, with or without abdominal pain, but with leukocytosis with a marked left shift, tachycardia, hemoconcentration, and general malaise. No causal relationship between MIFEPREX and misoprostol use and an increased risk of infection or death has been established. *Clostridium sordellii* infections have also been reported very rarely following childbirth (vaginal delivery and caesarian section), and in other gynecologic and non-gynecologic conditions.

5.2 Uterine Bleeding

Uterine bleeding occurs in almost all patients during a medical abortion. Prolonged heavy bleeding (soaking through two thick full-size sanitary pads per hour for two consecutive hours) may be a sign of incomplete abortion or other complications, and prompt medical or surgical intervention may be needed to prevent the development of hypovolemic shock. Counsel patients to seek immediate medical attention if they experience prolonged heavy vaginal bleeding following a medical abortion [see *Boxed Warning*].

Women should expect to experience vaginal bleeding or spotting for an average of 9 to 16 days. Women report experiencing heavy bleeding for a median duration of 2 days. Up to 8% of all subjects may experience some type of bleeding for 30 days or more. In general, the duration of bleeding and spotting increased as the duration of the pregnancy increased.

Decreases in hemoglobin concentration, hematocrit, and red blood cell count may occur in patients who bleed heavily.

Excessive uterine bleeding usually requires treatment by uterotonics, vasoconstrictor drugs, surgical uterine evacuation, administration of saline infusions, and/or blood transfusions. Based on data from several large clinical trials, vasoconstrictor drugs were used in 4.3% of all subjects, there was a decrease in hemoglobin of more than 2 g/dL in 5.5% of subjects, and blood transfusions were administered to ≤ 0.1% of subjects. Because heavy bleeding requiring surgical uterine evacuation occurs in about 1% of patients, special care should be given to patients with hemostatic disorders, hypocoagulability, or severe anemia.

5.3 Mifepristone REMS Program

MIFEPREX is available only through a restricted program under a REMS called the mifepristone REMS Program, because of the risks of serious complications [see *Warnings and Precautions* (5.1, 5.2)].

Notable requirements of the mifepristone REMS Program include the following:

- Prescribers must be certified with the program by completing the Prescriber Agreement Form.
- Patients must sign a Patient Agreement Form.
- MIFEPREX must only be dispensed to patients by or under the supervision of a certified prescriber, or by certified pharmacies on prescriptions issued by certified prescribers.

Further information is available at 1-877-4 Early Option (1-877-432-7596).

5.4 Ectopic Pregnancy

MIFEPREX is contraindicated in patients with a confirmed or suspected ectopic pregnancy because MIFEPREX is not effective for terminating ectopic pregnancies [see *Contraindications* (4)]. Healthcare providers should remain alert to the possibility that a patient who is undergoing a medical abortion could have an undiagnosed ectopic pregnancy because some of the expected symptoms experienced with a medical abortion (abdominal pain, uterine bleeding) may be similar to those of a ruptured ectopic pregnancy. The presence of an ectopic pregnancy may have been missed even if the patient underwent ultrasonography prior to being prescribed MIFEPREX.

Patients who became pregnant with an IUD in place should be assessed for ectopic pregnancy.

5.5 Rhesus Immunization

The use of MIFEPREX is assumed to require the same preventive measures as those taken prior to and during surgical abortion to prevent rhesus immunization.

6 ADVERSE REACTIONS

The following adverse reactions are described in greater detail in other sections:

- Infection and sepsis [see *Warnings and Precautions* (5.1)]
- Uterine bleeding [see *Warnings and Precautions* (5.2)]

6.1 Clinical Trials Experience

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

Information presented on common adverse reactions relies solely on data from U.S. studies, because rates reported in non-U.S. studies were markedly lower and are not likely generalizable to the U.S. population. In three U.S. clinical studies totaling 1,248 women through 70 days gestation who used mifepristone 200 mg orally followed 24-48 hours later by misoprostol 800 mcg buccally, women reported adverse reactions in diaries and in interviews at the follow-up visit. These studies enrolled generally healthy women of reproductive age without contraindications to mifepristone or misoprostol use according to the MIFEPREX product label. Gestational age was assessed prior to study enrollment using the date of the woman's last menstrual period, clinical evaluation, and/or ultrasound examination.

About 85% of patients report at least one adverse reaction following administration of MIFEPREX and misoprostol, and many can be expected to report more than one such reaction. The most commonly reported adverse reactions (>15%) were nausea, weakness, fever/chills, vomiting, headache, diarrhea, and dizziness (see Table 1). The frequency of adverse reactions varies between studies and may be dependent on many factors, including the patient population and gestational age.

Abdominal pain/cramping is expected in all medical abortion patients and its incidence is not reported in clinical studies. Treatment with MIFEPREX and misoprostol is designed to induce uterine bleeding and cramping to cause termination of an intrauterine pregnancy. Uterine bleeding and cramping are expected consequences of the action of MIFEPREX and misoprostol as used in the treatment procedure. Most patients can expect bleeding more heavily than they do during a heavy menstrual period [see *Warnings and Precautions* (5.2)].

Table 1 lists the adverse reactions reported in U.S. clinical studies with incidence >15% of women.

Table 1
Adverse Reactions Reported in Women Following Administration of Mifepristone (oral) and Misoprostol (buccal) in U.S. Clinical Studies

Adverse Reaction	# U.S. studies	Number of Evaluable Women	Range of frequency (%)	Upper Gestational Age of Studies Reporting Outcome
Nausea	3	1,248	51-75%	70 days
Weakness	2	630	55-58%	63 days
Fever/chills	1	414	48%	63 days
Vomiting	3	1,248	37-48%	70 days
Headache	2	630	41-44%	63 days
Diarrhea	3	1,248	18-43%	70 days
Dizziness	2	630	39-41%	63 days

One study provided gestational-age stratified adverse reaction rates for women who were 57-63 and 64-70 days; there was little difference in frequency of the reported common adverse reactions by gestational age.

Information on serious adverse reactions was reported in six U.S. and four non-U.S. clinical studies, totaling 30,966 women through 70 days gestation who used mifepristone 200 mg orally followed 24-48 hours later by misoprostol 800 mcg buccally. Serious adverse reaction rates were similar between U.S. and non-U.S. studies, so rates from both U.S. and non-U.S. studies are presented. In the U.S. studies, one studied women through 56 days gestation, four through 63 days gestation, and one through 70 days gestation, while in the non-U.S. studies, two studied women through 63 days gestation, and two through 70 days gestation. Serious adverse reactions were reported in <0.5% of women. Information from the U.S. and non-U.S. studies is presented in Table 2.

Table 2
Serious Adverse Reactions Reported in Women Following Administration of Mifepristone (oral) and Misoprostol (buccal) in U.S. and Non-U.S. Clinical Studies

Adverse Reaction	U.S.			Non-U.S.		
	# of studies	Number of Evaluable Women	Range of frequency (%)	# of studies	Number of Evaluable Women	Range of frequency (%)
Transfusion	4	17,774	0.03-0.5%	3	12,134	0-0.1%
Sepsis	1	629	0.2%	1	11,155	<0.01%*
ER visit	2	1,043	2.9-4.6%	1	95	0
Hospitalization Related to Medical Abortion	3	14,339	0.04-0.6%	3	1,286	0-0.7%
Infection without sepsis	1	216	0	1	11,155	0.2%
Hemorrhage	NR	NR	NR	1	11,155	0.1%

NR= Not reported

* This outcome represents a single patient who experienced death related to sepsis.

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of MIFEPREX and misoprostol. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Infections and infestations: post-abortion infection (including endometritis, endomyometritis, parametritis, pelvic infection, pelvic inflammatory disease, salpingitis)

Blood and the lymphatic system disorders: anemia

Immune system disorders: allergic reaction (including anaphylaxis, angioedema, hives, rash, itching)

Psychiatric disorders: anxiety

Cardiac disorders: tachycardia (including racing pulse, heart palpitations, heart pounding)

Vascular disorders: syncope, fainting, loss of consciousness, hypotension (including orthostatic), light-headedness

Respiratory, thoracic and mediastinal disorders: shortness of breath

Gastrointestinal disorders: dyspepsia

Musculoskeletal, connective tissue and bone disorders: back pain, leg pain

Reproductive system and breast disorders: uterine rupture, ruptured ectopic pregnancy, hematometra, leukorrhea

General disorders and administration site conditions: pain

7 DRUG INTERACTIONS

7.1 Drugs that May Reduce MIFEPREX Exposure (Effect of CYP 3A4 Inducers on MIFEPREX)

CYP450 3A4 is primarily responsible for the metabolism of mifepristone. CYP3A4 inducers such as rifampin, dexamethasone, St. John's Wort, and certain anticonvulsants (such as phenytoin, phenobarbital, carbamazepine) may induce mifepristone metabolism (lowering serum concentrations of mifepristone). Whether this action has an impact on the efficacy of the dose

regimen is unknown. Refer to the follow-up assessment [see *Dosage and Administration* (2.3)] to verify that treatment has been successful.

7.2 Drugs that May Increase MIFEPREX Exposure (Effect of CYP 3A4 Inhibitors on MIFEPREX)

Although specific drug or food interactions with mifepristone have not been studied, on the basis of this drug's metabolism by CYP 3A4, it is possible that ketoconazole, itraconazole, erythromycin, and grapefruit juice may inhibit its metabolism (increasing serum concentrations of mifepristone). MIFEPREX should be used with caution in patients currently or recently treated with CYP 3A4 inhibitors.

7.3 Effects of MIFEPREX on Other Drugs (Effect of MIFEPREX on CYP 3A4 Substrates)

Based on *in vitro* inhibition information, coadministration of mifepristone may lead to an increase in serum concentrations of drugs that are CYP 3A4 substrates. Due to the slow elimination of mifepristone from the body, such interaction may be observed for a prolonged period after its administration. Therefore, caution should be exercised when mifepristone is administered with drugs that are CYP 3A4 substrates and have narrow therapeutic range.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

MIFEPREX is indicated, in a regimen with misoprostol, for the medical termination of intrauterine pregnancy through 70 days gestation. Risks to pregnant patients are discussed throughout the labeling.

Refer to misoprostol labeling for risks to pregnant patients with the use of misoprostol.

The risk of adverse developmental outcomes with a continued pregnancy after a failed pregnancy termination with MIFEPREX in a regimen with misoprostol is unknown; however, the process of a failed pregnancy termination could disrupt normal embryo-fetal development and result in adverse developmental effects. Birth defects have been reported with a continued pregnancy after a failed pregnancy termination with MIFEPREX in a regimen with misoprostol. In animal reproduction studies, increased fetal losses were observed in mice, rats, and rabbits and skull deformities were observed in rabbits with administration of mifepristone at doses lower than the human exposure level based on body surface area.

Data

Animal Data

In teratology studies in mice, rats and rabbits at doses of 0.25 to 4.0 mg/kg (less than 1/100 to approximately 1/3 the human exposure based on body surface area), because of the antiprogestational activity of mifepristone, fetal losses were much higher than in control animals. Skull deformities were detected in rabbit studies at approximately 1/6 the human exposure, although no teratogenic effects of mifepristone have been observed to date in rats or mice. These deformities were most likely due to the mechanical effects of uterine contractions resulting from inhibition of progesterone action.

8.2 Lactation

MIFEPREX is present in human milk. Limited data demonstrate undetectable to low levels of the drug in human milk with the relative (weight-adjusted) infant dose 0.5% or less as compared to maternal dosing. There is no information on the effects of MIFEPREX in a regimen with

misoprostol in a breastfed infant or on milk production. Refer to misoprostol labeling for lactation information with the use of misoprostol. The developmental and health benefits of breast-feeding should be considered along with any potential adverse effects on the breast-fed child from MIFEPREX in a regimen with misoprostol.

8.4 Pediatric Use

Safety and efficacy of MIFEPREX have been established in pregnant females. Data from a clinical study of MIFEPREX that included a subset of 322 females under age 17 demonstrated a safety and efficacy profile similar to that observed in adults.


10 OVERDOSAGE

No serious adverse reactions were reported in tolerance studies in healthy non-pregnant female and healthy male subjects where mifepristone was administered in single doses greater than 1800 mg (ninefold the recommended dose for medical abortion). If a patient ingests a massive overdose, the patient should be observed closely for signs of adrenal failure.

11 DESCRIPTION

MIFEPREX tablets each contain 200 mg of mifepristone, a synthetic steroid with antiprogestational effects. The tablets are light yellow in color, cylindrical, and bi-convex, and are intended for oral administration only. The tablets include the inactive ingredients colloidal silica anhydrous, corn starch, povidone, microcrystalline cellulose, and magnesium stearate.

Mifepristone is a substituted 19-nor steroid compound chemically designated as 11 β -[p-(Dimethylamino)phenyl]-17 β -hydroxy-17-(1-propynyl)estra-4,9-dien-3-one. Its empirical formula is C₂₉H₃₅NO₂. Its structural formula is:



The compound is a yellow powder with a molecular weight of 429.6 and a melting point of 192-196°C. It is very soluble in methanol, chloroform and acetone and poorly soluble in water, hexane and isopropyl ether.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The anti-progestational activity of mifepristone results from competitive interaction with progesterone at progesterone-receptor sites. Based on studies with various oral doses in several animal species (mouse, rat, rabbit, and monkey), the compound inhibits the activity of endogenous or exogenous progesterone, resulting in effects on the uterus and cervix that, when combined with misoprostol, result in termination of an intrauterine pregnancy.

During pregnancy, the compound sensitizes the myometrium to the contraction-inducing activity

of prostaglandins.

12.2 Pharmacodynamics

Use of MIFEPREX in a regimen with misoprostol disrupts pregnancy by causing decidual necrosis, myometrial contractions, and cervical softening, leading to the expulsion of the products of conception.

Doses of 1 mg/kg or greater of mifepristone have been shown to antagonize the endometrial and myometrial effects of progesterone in women.

Antiglucocorticoid and antiandrogenic activity: Mifepristone also exhibits antiglucocorticoid and weak antiandrogenic activity. The activity of the glucocorticoid dexamethasone in rats was inhibited following doses of 10 to 25 mg/kg of mifepristone. Doses of 4.5 mg/kg or greater in human beings resulted in a compensatory elevation of adrenocorticotrophic hormone (ACTH) and cortisol. Antiandrogenic activity was observed in rats following repeated administration of doses from 10 to 100 mg/kg.

12.3 Pharmacokinetics

Mifepristone is rapidly absorbed after oral ingestion with non-linear pharmacokinetics for C_{max} after single oral doses of 200 mg and 600 mg in healthy subjects.

Absorption

The absolute bioavailability of a 20 mg mifepristone oral dose in females of childbearing age is 69%. Following oral administration of a single dose of 600 mg, mifepristone is rapidly absorbed, with a peak plasma concentration of 1.98 ± 1.0 mg/L occurring approximately 90 minutes after ingestion.

Following oral administration of a single dose of 200 mg in healthy men (n=8), mean C_{max} was 1.77 ± 0.7 mg/L occurring approximately 45 minutes after ingestion. Mean AUC_{0-∞} was 25.8 ± 6.2 mg*hr/L.

Distribution

Mifepristone is 98% bound to plasma proteins, albumin, and α_1 -acid glycoprotein. Binding to the latter protein is saturable, and the drug displays nonlinear kinetics with respect to plasma concentration and clearance.

Elimination

Following a distribution phase, elimination of mifepristone is slow at first (50% eliminated between 12 and 72 hours) and then becomes more rapid with a terminal elimination half-life of 18 hours.

Metabolism

Metabolism of mifepristone is primarily via pathways involving N-demethylation and terminal hydroxylation of the 17-propynyl chain. *In vitro* studies have shown that CYP450 3A4 is primarily responsible for the metabolism. The three major metabolites identified in humans are: (1) RU 42 633, the most widely found in plasma, is the N-monodemethylated metabolite; (2) RU 42 848, which results from the loss of two methyl groups from the 4-dimethylaminophenyl in position 11β; and (3) RU 42 698, which results from terminal hydroxylation of the 17-propynyl chain.

Excretion

By 11 days after a 600 mg dose of tritiated compound, 83% of the drug has been accounted for by the feces and 9% by the urine. Serum concentrations are undetectable by 11 days.

Specific Populations

The effects of age, hepatic disease and renal disease on the safety, efficacy and pharmacokinetics of mifepristone have not been investigated.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

No long-term studies to evaluate the carcinogenic potential of mifepristone have been performed.

Mutagenesis

Results from studies conducted *in vitro* and in animals have revealed no genotoxic potential for mifepristone. Among the tests carried out were: Ames test with and without metabolic activation; gene conversion test in *Saccharomyces cerevisiae* D4 cells; forward mutation in *Schizosaccharomyces pombe* P1 cells; induction of unscheduled DNA synthesis in cultured HeLa cells; induction of chromosome aberrations in CHO cells; *in vitro* test for gene mutation in V79 Chinese hamster lung cells; and micronucleus test in mice.

Impairment of Fertility

In rats, administration of 0.3 mg/kg mifepristone per day caused severe disruption of the estrus cycles for the three weeks of the treatment period. Following resumption of the estrus cycle, animals were mated and no effects on reproductive performance were observed.

14 CLINICAL STUDIES

Safety and efficacy data from clinical studies of mifepristone 200 mg orally followed 24-48 hours later by misoprostol 800 mcg buccally through 70 days gestation are reported below. Success was defined as the complete expulsion of the products of conception without the need for surgical intervention. The overall rates of success and failure, shown by reason for failure based on 22 worldwide clinical studies (including 7 U.S. studies) appear in Table 3.

The demographics of women who participated in the U.S. clinical studies varied depending on study location and represent the racial and ethnic variety of American females. Females of all reproductive ages were represented, including females less than 18 and more than 40 years of age; most were 27 years or younger.

Table 3
Outcome Following Treatment with Mifepristone (oral) and Misoprostol (buccal)
Through 70 Days Gestation

	U.S. Trials	Non-U.S. Trials
N	16,794	18,425
Complete Medical Abortion	97.4%	96.2%
Surgical Intervention*	2.6%	3.8%
Ongoing Pregnancy**	0.7%	0.9%
* Reasons for surgical intervention include ongoing pregnancy, medical necessity, persistent or heavy bleeding after treatment, patient request, or incomplete expulsion. ** Ongoing pregnancy is a subcategory of surgical intervention, indicating the percent of women who have surgical intervention due to an ongoing pregnancy.		

The results for clinical studies that reported outcomes, including failure rates for ongoing pregnancy, by gestational age are presented in Table 4.

Table 4
Outcome by Gestational Age Following Treatment with Mifepristone and
Misoprostol (buccal) for U.S. and Non-U.S. Clinical Studies

	≤49 days			50-56 days			57-63 days			64-70 days		
	N	%	Number of Evaluable Studies	N	%	Number of Evaluable Studies	N	%	Number of Evaluable Studies	N	%	Number of Evaluable Studies
Complete medical abortion	12,046	98.1	10	3,941	96.8	7	2,294	94.7	9	479	92.7	4
Surgical intervention for ongoing pregnancy	10,272	0.3	6	3,788	0.8	6	2,211	2	8	453	3.1	3

One clinical study asked subjects through 70 days gestation to estimate when they expelled the pregnancy, with 70% providing data. Of these, 23-38% reported expulsion within 3 hours and over 90% within 24 hours of using misoprostol.

16 HOW SUPPLIED/STORAGE AND HANDLING

is only available through a restricted program called the Mifepristone REMS Program [see *Warnings and Precautions* (5.3)].

MIFEPREX is supplied as light yellow, cylindrical, and bi-convex tablets imprinted on one side with "MF." Each tablet contains 200 mg of mifepristone. One tablet is individually blistered on one blister card that is packaged in an individual package (National Drug Code 64875-001-01).

Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F) [see USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide), included with each package of MIFEPREX. Additional copies of the Medication Guide are available by contacting Danco Laboratories at 1-877-4 Early Option (1-877-432-7596) or from www.earlyoptionpill.com.

Serious Infections and Bleeding

- Inform the patient that uterine bleeding and uterine cramping will occur [see *Warnings and Precautions* (5.2)].
- Advise the patient that serious and sometimes fatal infections and bleeding can occur very rarely [see *Warnings and Precautions* (5.1, 5.2)].
- MIFEPREX is only available through a restricted program called the Mifepristone REMS Program [see *Warnings and Precautions* (5.3)]. Under the mifepristone REMS Program:
 - o Patients must sign a Patient Agreement Form.
 - o MIFEPREX is only dispensed by or under the supervision of certified prescribers or by certified pharmacies on prescriptions issued by certified prescribers.

Provider Contacts and Actions in Case of Complications

- Ensure that the patient knows whom to call and what to do, including going to an Emergency Room if none of the provided contacts are reachable, or if the patient experiences complications including prolonged heavy bleeding, severe abdominal pain, or sustained fever [see *Boxed Warning*].
-

Compliance with Treatment Schedule and Follow-up Assessment

- Advise the patient that it is necessary to complete the treatment schedule, including a follow-up assessment approximately 7 to 14 days after taking MIFEPREX [see *Dosage and Administration* (2.3)].
- Explain that
 - o prolonged heavy vaginal bleeding is not proof of a complete abortion,
 - o if the treatment fails and the pregnancy continues, the risk of fetal malformation is unknown,
 - o it is recommended that ongoing pregnancy be managed by surgical termination [see *Dosage and Administration* (2.3)]. Advise the patient whether you will provide such care or will refer them to another provider.

Subsequent Fertility

- Inform the patient that another pregnancy can occur following medical abortion and before resumption of normal menses.
- Inform the patient that contraception can be initiated as soon as pregnancy expulsion has been confirmed, or before resuming sexual intercourse.

MIFEPREX is a registered trademark of Danco Laboratories, LLC.

Manufactured for:
Danco Laboratories, LLC
P.O. Box 4816
New York, NY 10185
1-877-4 Early Option (1-877-432-7596)
www.earlyoptionpill.com

01/2023

MEDICATION GUIDE

Mifeprex (MIF-eh-prex) (mifepristone tablets, for oral use)

Read this information carefully before taking Mifeprex and misoprostol. It will help you understand how the treatment works. This Medication Guide does not take the place of talking with your healthcare provider.

What is the most important information I should know about Mifeprex?

What symptoms should I be concerned with? Although cramping and bleeding are an expected part of ending a pregnancy, rarely, serious and potentially life-threatening bleeding, infections, or other problems can occur following a miscarriage, surgical abortion, medical abortion, or childbirth. Seeking medical attention as soon as possible is needed in these circumstances. Serious infection has resulted in death in a very small number of cases. There is no information that use of Mifeprex and misoprostol caused these deaths. If you have any questions, concerns, or problems, or if you are worried about any side effects or symptoms, you should contact your healthcare provider. You can write down your healthcare provider's telephone number here _____.

Be sure to contact your healthcare provider promptly if you have any of the following:

- **Heavy Bleeding.** Contact your healthcare provider right away if you bleed enough to soak through two thick full-size sanitary pads per hour for two consecutive hours or if you are concerned about heavy bleeding. In about 1 out of 100 women, bleeding can be so heavy that it requires a surgical procedure (surgical aspiration or D&C).
- **Abdominal Pain or "Feeling Sick."** If you have abdominal pain or discomfort, or you are "feeling sick," including weakness, nausea, vomiting, or diarrhea, with or without fever, more than 24 hours after taking misoprostol, you should contact your healthcare provider without delay. These symptoms may be a sign of a serious infection or another problem (including an ectopic pregnancy, a pregnancy outside the womb).
- **Fever.** In the days after treatment, if you have a fever of 100.4°F or higher that lasts for more than 4 hours, you should contact your healthcare provider right away. Fever may be a symptom of a serious infection or another problem.

If you cannot reach your healthcare provider, go to the nearest hospital emergency room.

What to do if you are still pregnant after Mifeprex with misoprostol treatment. If you are still pregnant, your healthcare provider will talk with you about a surgical procedure to end your pregnancy. In many cases, this surgical procedure can be done in the office/clinic. The chance of birth defects if the pregnancy is not ended is unknown.

Talk with your healthcare provider. Before you take Mifeprex, you should read this Medication Guide and you and your healthcare provider should discuss the benefits and risks of your using Mifeprex.

What is Mifeprex?

Mifeprex is used in a regimen with another prescription medicine called misoprostol, to end an early pregnancy. Early pregnancy means it is 70 days (10 weeks) or less since your last menstrual period began. Mifeprex is not approved for ending pregnancies that are further along. Mifeprex blocks a hormone needed for your pregnancy to continue. When you use Mifeprex on Day 1, you also need to take another medicine called misoprostol 24 to 48 hours after you take Mifeprex, to cause the pregnancy to be passed from your uterus.

The pregnancy is likely to be passed from your uterus within 2 to 24 hours after taking Mifeprex and misoprostol. When the pregnancy is passed from the uterus, you will have bleeding and cramping that will likely be heavier than your usual period. About 2 to 7 out of 100 women taking Mifeprex will need a surgical procedure because the pregnancy did not completely pass from the uterus or to stop bleeding.

Who should not take Mifeprex?

Some patients should not take Mifeprex. Do not take Mifeprex if you:

- Have a pregnancy that is more than 70 days (10 weeks). Your healthcare provider may do a clinical examination, an ultrasound examination, or other testing to determine how far along you are in pregnancy.
- Are using an IUD (intrauterine device or system). It must be taken out before you take Mifeprex.
- Have been told by your healthcare provider that you have a pregnancy outside the uterus (ectopic pregnancy).
- Have problems with your adrenal glands (chronic adrenal failure).
- Take a medicine to thin your blood.
- Have a bleeding problem.
- Have porphyria.
- Take certain steroid medicines.
- Are allergic to mifepristone, misoprostol, or medicines that contain misoprostol, such as Cytotec or Arthrotec.

Ask your healthcare provider if you are not sure about all your medical conditions before taking this medicine to find out if you can take Mifeprex.

What should I tell my healthcare provider before taking Mifeprex?

Before you take Mifeprex, tell your healthcare provider if you:

- cannot follow-up within approximately 7 to 14 days of your first visit
 - are breastfeeding. Mifeprex can pass into your breast milk. The effect of the Mifeprex and misoprostol regimen on the breastfed infant or on milk production is unknown.
 - are taking medicines, including prescription and over-the-counter medicines, vitamins, and herbal supplements.
- Mifeprex and certain other medicines may affect each other if they are used together. This can cause side effects.

How should I take Mifeprex?

- Mifeprex will be given to you by a healthcare provider or pharmacy.
- You and your healthcare provider will plan the most appropriate location for you to take the misoprostol, because it may cause bleeding, cramps, nausea, diarrhea, and other symptoms that usually begin within 2 to 24 hours after taking it.
- Most women will pass the pregnancy within 2 to 24 hours after taking the misoprostol tablets.

Follow the instruction below on how to take Mifeprex and misoprostol:**Mifeprex (1 tablet) orally + misoprostol (4 tablets) buccally****Day 1:**

- Take 1 Mifeprex tablet by mouth.

24 to 48 hours after taking Mifeprex:

- Take 4 misoprostol tablets by placing 2 tablets in each cheek pouch (the area between your teeth and cheek - see Figure A) for 30 minutes and then swallow anything left over with a drink of water or another liquid.
- The medicines may not work as well if you take misoprostol sooner than 24 hours after Mifeprex or later than 48 hours after Mifeprex.
- Misoprostol often causes cramps, nausea, diarrhea, and other symptoms. Your healthcare provider may send you home with medicines for these symptoms.



Figure A (2 tablets between your left cheek and gum and 2 tablets between your right cheek and gum).

Follow-up Assessment at Day 7 to 14:

- This follow-up assessment is very important. You must follow-up with your healthcare provider about 7 to 14 days after you have taken Mifeprex to be sure you are well and that you have had bleeding and the pregnancy has passed from your uterus.
- Your healthcare provider will assess whether your pregnancy has passed from your uterus. If your pregnancy continues, the chance that there may be birth defects is unknown. If you are still pregnant, your healthcare provider will talk with you about a surgical procedure to end your pregnancy.
- If your pregnancy has ended, but has not yet completely passed from your uterus, your provider will talk with you about other choices you have, including waiting, taking another dose of misoprostol, or having a surgical procedure to empty your uterus.

When should I begin birth control?

You can become pregnant again right after your pregnancy ends. If you do not want to become pregnant again, start using birth control as soon as your pregnancy ends or before you start having sexual intercourse again.

What should I avoid while taking Mifeprex and misoprostol?

Do not take any other prescription or over-the-counter medicines (including herbal medicines or supplements) at any time during the treatment period without first asking your healthcare provider about them because they may interfere with the treatment. Ask your healthcare provider about what medicines you can take for pain and other side effects.

What are the possible side effects of Mifeprex and misoprostol?

Mifeprex may cause serious side effects. See “What is the most important information I should know about Mifeprex?”

Cramping and bleeding. Cramping and vaginal bleeding are expected with this treatment. Usually, these symptoms mean that the treatment is working. But sometimes you can get cramping and bleeding and still be pregnant. This is why you must follow-up with your healthcare provider approximately 7 to 14 days after taking Mifeprex. See “How should I take Mifeprex?” for more information on your follow-up assessment. If you are not already bleeding after taking Mifeprex, you probably will begin to bleed once you take misoprostol, the medicine you take 24 to 48 hours after Mifeprex. Bleeding or spotting can be expected for an average of 9 to 16 days and may last for up to 30 days. Your bleeding may be similar to, or greater than, a normal heavy period. You may see blood clots and tissue. This is an expected part of passing the pregnancy.

The most common side effects of Mifeprex treatment include: nausea, weakness, fever/chills, vomiting, headache, diarrhea and dizziness. Your provider will tell you how to manage any pain or other side effects. These are not all the possible side effects of Mifeprex.

Call your healthcare provider for medical advice about any side effects that bother you or do not go away. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of Mifeprex.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. This Medication Guide summarizes the most important information about Mifeprex. If you would like more information, talk with your healthcare provider. You may ask your healthcare provider for information about Mifeprex that is written for healthcare professionals.

For more information about Mifeprex, go to www.earlyoptionpill.com or call 1-877-4 Early Option (1-877-432-7596).

Manufactured for: *Danco Laboratories, LLC*
P.O. Box 4816
New York, NY 10185
1-877-4 Early Option (1-877-432-7596) www.earlyoptionpill.com

This Medication Guide has been approved by the U.S. Food and Drug Administration. Approval 01/2023

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

020687Orig1s025

REMS

Initial Shared System REMS approval: 04/2019

Most Recent Modification: 01/2023

Mifepristone Tablets, 200 mg

Progestin Antagonist

RISK EVALUATION AND MITIGATION STRATEGY (REMS) SINGLE SHARED SYSTEM FOR MIFEPRISTONE 200 MG

I. GOAL

The goal of the REMS for mifepristone is to mitigate the risk of serious complications associated with mifepristone by:

- a) Requiring healthcare providers who prescribe mifepristone to be certified in the Mifepristone REMS Program.
- b) Ensuring that mifepristone is only dispensed by or under the supervision of certified prescribers, or by certified pharmacies on prescriptions issued by certified prescribers.
- c) Informing patients about the risk of serious complications associated with mifepristone.

II. REMS ELEMENTS

A. Elements to Assure Safe Use

1. Healthcare providers who prescribe mifepristone must be specially certified.
 - a. To become specially certified to prescribe mifepristone, healthcare providers must:
 - i. Review the Prescribing Information for mifepristone.
 - ii. Complete a *Prescriber Agreement Form*. By signing¹ a *Prescriber Agreement Form*, prescribers agree that:
 - 1) They have the following qualifications:
 - a) Ability to assess the duration of pregnancy accurately
 - b) Ability to diagnose ectopic pregnancies
 - c) Ability to provide surgical intervention in cases of incomplete abortion or severe bleeding, or to have made plans to provide such care through others, and ability to assure patient access to medical facilities equipped to provide blood transfusions and resuscitation, if necessary
 - 2) They will follow the guidelines for use of mifepristone (see b.i-vii below).
 - b. As a condition of certification, prescribers must follow the guidelines for use of mifepristone described below:
 - i. Ensure that the *Patient Agreement Form* is reviewed with the patient and the risks of the mifepristone treatment regimen are fully explained. Ensure any questions the patient may have prior to receiving mifepristone are answered.
 - ii. Ensure that the healthcare provider and patient sign the *Patient Agreement Form*.

¹ In this REMS, the terms “sign” and “signature” include electronic signatures.

- iii. Ensure that the patient is provided with a copy of the *Patient Agreement Form* and Medication Guide.
 - iv. Ensure that the signed *Patient Agreement Form* is placed in the patient's medical record.
 - v. Ensure that any deaths are reported to the Mifepristone Sponsor that provided the mifepristone, identifying the patient by a non-identifiable reference and including the NDC and lot number from the package of mifepristone that was dispensed to the patient.
 - vi. If mifepristone will be dispensed by a certified pharmacy:
 - 1) Provide the certified pharmacy a signed *Prescriber Agreement Form*.
 - 2) Assess appropriateness of dispensing mifepristone when contacted by a certified pharmacy about patients who will receive mifepristone more than 4 calendar days after the prescription was received by the certified pharmacy.
 - 3) Obtain the NDC and lot number of the package of mifepristone the patient received in the event the prescriber becomes aware of the death of the patient.
 - vii. The certified prescriber who dispenses mifepristone or who supervises the dispensing of mifepristone must:
 - 1) Provide an authorized distributor with a signed *Prescriber Agreement Form*.
 - 2) Ensure that the NDC and lot number from each package of mifepristone dispensed are recorded in the patient's record.
 - 3) Ensure that healthcare providers under their supervision follow guidelines i.-v.
 - c. Mifepristone Sponsors must:
 - i. Ensure that healthcare providers who prescribe their mifepristone are specially certified in accordance with the requirements described above and de-certify healthcare providers who do not maintain compliance with certification requirements.
 - ii. Ensure prescribers previously certified in the Mifepristone REMS Program complete the new *Prescriber Agreement Form*:
 - 1) Within 120 days after approval of this modification, for those previously certified prescribers submitting prescriptions to certified pharmacies.
 - 2) Within one year after approval of this modification, if previously certified and ordering from an authorized distributor.
 - iii. Ensure that healthcare providers can complete the certification process by email or fax to an authorized distributor and/or certified pharmacy.
 - iv. Provide the Prescribing Information and their *Prescriber Agreement Form* to healthcare providers who inquire about how to become certified.
 - v. Ensure annually with each certified prescriber that their locations for receiving mifepristone are up to date.
- The following materials are part of the Mifepristone REMS Program:
- *Prescriber Agreement Form for Danco Laboratories, LLC*
 - *Prescriber Agreement Form for GenBioPro, Inc.*
 - *Patient Agreement Form*

2. Pharmacies that dispense mifepristone must be specially certified
 - a. To become specially certified to dispense mifepristone, pharmacies must:
 - i. Be able to receive *Prescriber Agreement Forms* by email and fax.
 - ii. Be able to ship mifepristone using a shipping service that provides tracking information.
 - iii. Designate an authorized representative to carry out the certification process on behalf of the pharmacy.
 - iv. Ensure the authorized representative oversees implementation and compliance with the Mifepristone REMS Program by doing the following:
 - 1) Review the Prescribing Information for mifepristone.
 - 2) Complete a *Pharmacy Agreement Form*. By signing a *Pharmacy Agreement Form*, the authorized representative agrees that the pharmacy will put processes and procedures in place to ensure the following requirements are completed:
 - a) Verify that the prescriber is certified by confirming their completed *Prescriber Agreement Form* was received with the prescription or is on file with the pharmacy.
 - b) Dispense mifepristone such that it is delivered to the patient within 4 calendar days of the date the pharmacy receives the prescription, except as provided in c) below.
 - c) Confirm with the prescriber the appropriateness of dispensing mifepristone for patients who will receive the drug more than 4 calendar days after the date the pharmacy receives the prescription and document the prescriber's decision.
 - d) Record in the patient's record the NDC and lot number from each package of mifepristone dispensed.
 - e) Track and verify receipt of each shipment of mifepristone.
 - f) Dispense mifepristone in its package as supplied by the Mifepristone Sponsor.
 - g) Report any patient deaths to the prescriber, including the NDC and lot number from the package of mifepristone dispensed to the patient, and remind the prescriber of their obligation to report the deaths to the Mifepristone Sponsor that provided the mifepristone. Notify the Mifepristone Sponsor that provided the dispensed mifepristone that the pharmacy submitted a report of death to the prescriber, including the name and contact information for the prescriber and the NDC and lot number of the dispensed product.
 - h) Not distribute, transfer, loan or sell mifepristone except to certified prescribers or other locations of the pharmacy.
 - i) Maintain records of *Prescriber Agreement Forms*.
 - j) Maintain records of dispensing and shipping.
 - k) Maintain records of all processes and procedures including compliance with those processes and procedures.
 - l) Maintain the identity of the patient and prescriber as confidential, including limiting access to patient and prescriber identity only to those personnel necessary to dispense mifepristone in accordance with the Mifepristone REMS Program requirements, or as necessary for payment and/or insurance purposes.
 - m) Train all relevant staff on the Mifepristone REMS Program requirements.

- n) Comply with audits carried out by the Mifepristone Sponsors or a third party acting on behalf of the Mifepristone Sponsors to ensure that all processes and procedures are in place and are being followed.
- b. Mifepristone Sponsors must:
 - i. Ensure that pharmacies are specially certified in accordance with the requirements described above and de-certify pharmacies that do not maintain compliance with certification requirements.
 - ii. Ensure that pharmacies can complete the certification process by email and fax to an authorized distributor.
 - i. Verify annually that the name and contact information for the pharmacy's authorized representative corresponds to that of the current designated authorized representative for the certified pharmacy, and if different, require the pharmacy to recertify with the new authorized representative.

The following materials are part of the Mifepristone REMS Program:

- *Pharmacy Agreement Form for Danco Laboratories, LLC*
 - *Pharmacy Agreement Form for GenBioPro, Inc.*
- 3. Mifepristone must be dispensed to patients with evidence or other documentation of safe use conditions as ensured by the certified prescriber in signing the *Prescriber Agreement Form*.
 - a. The patient must sign a *Patient Agreement Form* indicating that the patient has:
 - i. Received, read and been provided a copy of the *Patient Agreement Form*.
 - ii. Received counseling from the healthcare provider regarding the risk of serious complications associated with mifepristone.

B. Implementation System

- 1. Mifepristone Sponsors must ensure that their mifepristone is only distributed to certified prescribers and certified pharmacies by:
 - a. Ensuring that distributors who distribute their mifepristone comply with the program requirements for distributors.
 - i. The distributors must put processes and procedures in place to:
 - 1) Complete the certification process upon receipt of a *Prescriber Agreement Form* or *Pharmacy Agreement Form*.
 - 2) Notify healthcare providers and pharmacies when they have been certified by the Mifepristone REMS Program.
 - 3) Ship mifepristone only to certified pharmacies or locations identified by certified prescribers.
 - 4) Not ship mifepristone to pharmacies or prescribers who become de-certified from the Mifepristone REMS Program.
 - 5) Provide the Prescribing Information and their Prescriber Agreement Form to healthcare providers who (1) attempt to order mifepristone and are not yet certified, or (2) inquire about how to become certified.
 - ii. Put processes and procedures in place to maintain a distribution system that is secure,

confidential and follows all processes and procedures, including those for storage, handling, shipping, tracking package serial numbers, NDC and lot numbers, proof of delivery and controlled returns of mifepristone.

- iii. Train all relevant staff on the Mifepristone REMS Program requirements.
 - iv. Comply with audits by Mifepristone Sponsors or a third party acting on behalf of Mifepristone Sponsors to ensure that all processes and procedures are in place and are being followed for the Mifepristone REMS Program. In addition, distributors must maintain appropriate documentation and make it available for audits.
- b. Ensuring that distributors maintain secure and confidential distribution records of all shipments of mifepristone.
2. Mifepristone Sponsors must monitor their distribution data to ensure compliance with the Mifepristone REMS Program.
 3. Mifepristone Sponsors must ensure that adequate records are maintained to demonstrate that the Mifepristone REMS Program requirements have been met, including, but not limited to records of mifepristone distribution; certification of prescribers and pharmacies; and audits of pharmacies and distributors. These records must be readily available for FDA inspections.
 4. Mifepristone Sponsors must audit their new distributors within 90 calendar days and annually thereafter after the distributor is authorized to ensure that all processes and procedures are in place and functioning to support the requirements of the Mifepristone REMS Program. Mifepristone Sponsors will take steps to address their distributor compliance if noncompliance is identified.
 5. Mifepristone Sponsors must audit their certified pharmacies within 180 calendar days after the pharmacy places its first order of mifepristone, and annually thereafter audit certified pharmacies that have ordered mifepristone in the previous 12 months, to ensure that all processes and procedures are in place and functioning to support the requirements of the Mifepristone REMS Program. Mifepristone Sponsors will take steps to address their pharmacy compliance if noncompliance is identified.
 6. Mifepristone Sponsors must take reasonable steps to improve implementation of and compliance with the requirements of the Mifepristone REMS Program based on monitoring and assessment of the Mifepristone REMS Program.
 7. Mifepristone Sponsors must report to FDA any death associated with mifepristone whether or not considered drug-related, as soon as possible but no later than 15 calendar days from the initial receipt of the information by the Mifepristone Sponsor. This requirement does not affect the sponsors' other reporting and follow-up requirements under FDA regulations.

C. Timetable for Submission of Assessments

The NDA Sponsor must submit REMS assessments to FDA one year from the date of the approval of the modified REMS (1/3/2023) and annually thereafter. To facilitate inclusion of as much information as possible while allowing reasonable time to prepare the submission, the reporting interval covered by each assessment should conclude no earlier than 90 calendar days before the submission date for that assessment. The NDA Sponsor must submit each assessment so that it will be received by the FDA on or before the due date.

MIFEPREX® (Mifepristone) Tablets, 200 mg

PRESCRIBER AGREEMENT FORM

Mifeprex* (Mifepristone) Tablets, 200 mg, is indicated, in a regimen with misoprostol, for the medical termination of intrauterine pregnancy through 70 days gestation. Please see Prescribing Information and Medication Guide for complete safety information.

To **become a certified prescriber**, you must:

- **If you submit Mifeprex prescriptions for dispensing from certified pharmacies:**
 - o Submit this form to each certified pharmacy to which you intend to submit Mifeprex prescriptions. The form must be received by the certified pharmacy before any prescriptions are dispensed by that pharmacy.
- **If you order Mifeprex for dispensing by you or healthcare providers under your supervision:**
 - o Submit this form to the distributor. This form must be received by the distributor before the first order will be shipped to the healthcare setting.
 - o Healthcare settings, such as medical offices, clinics, and hospitals, where Mifeprex will be dispensed by or under the supervision of a certified prescriber in the Mifepristone REMS Program do not require pharmacy certification.

Prescriber Agreement: By signing this form, you agree that you meet the qualifications below and will follow the guidelines for use. You are responsible for overseeing implementation and compliance with the Mifepristone REMS Program. You also understand that if the guidelines below are not followed, the distributor may stop shipping mifepristone to the locations that you identify and certified pharmacies may stop accepting your mifepristone prescriptions.

Mifepristone must be provided by or under the supervision of a certified prescriber who meets the following qualifications:

- Ability to assess the duration of pregnancy accurately.
- Ability to diagnose ectopic pregnancies.
- Ability to provide surgical intervention in cases of incomplete abortion or severe bleeding, or have made plans to provide such care through others, and be able to assure patient access to medical facilities equipped to provide blood transfusions and resuscitation, if necessary.
- Has read and understood the Prescribing Information for mifepristone. The Prescribing Information is available by calling 1-877-4 EARLY OPTION (1-877-432-7596 toll-free), or by visiting www.earlyoptionpill.com.

In addition to meeting these qualifications, you also agree to follow these guidelines for use:

- Ensure that the *Patient Agreement Form* is reviewed with the patient and the risks of the mifepristone treatment regimen are fully explained. Ensure any questions the patient may have prior to receiving mifepristone are answered.
- Ensure the healthcare provider and patient sign the *Patient Agreement Form*.
- Ensure that the patient is provided with a copy of the *Patient Agreement Form* and Medication Guide.
- Ensure that the signed *Patient Agreement Form* is placed in the patient's medical record.
- Ensure that any deaths of patients who received Mifeprex are reported to Danco Laboratories, LLC, identifying the patient by a non-identifiable reference and including the NDC and lot number from the package of Mifeprex that was dispensed to the patient.



*MIFEPREX is a registered trademark of Danco Laboratories, LLC
P.O. Box 4816-New York, NY 10185

1-877-4-EARLY-OPTION (1-877-432-7596) www.earlyoptionpill.com

Ensure that healthcare providers under your supervision follow the guidelines listed above.

- If Mifeprex will be dispensed through a certified pharmacy:
 - o Assess appropriateness of dispensing Mifeprex when contacted by a certified pharmacy about patients who will receive Mifeprex more than 4 calendar days after the prescription was received by the certified pharmacy.
 - o Obtain the NDC and lot number of the package of Mifeprex the patient received in the event the prescriber becomes aware of the death of a patient.
- If Mifeprex will be dispensed by you or by healthcare providers under your supervision:
 - o Ensure the NDC and lot number from each package of Mifeprex are recorded in the patient's record.

I understand that a certified pharmacy may dispense mifepristone made by a different manufacturer than that stated on this Prescriber Agreement Form.

Print Name: _____ Title: _____

Signature: _____ Date: _____

Medical License # _____ State _____

NPI # _____

Practice Setting Address: _____

Return completed form to Mifeprex@dancodistributor.com or fax to 1-866-227-3343.

Approved 01/2023 [Doc control ID]



*MIFEPREX is a registered trademark of Danco Laboratories, LLC
P.O. Box 4816-New York, NY 10185
1-877-4-EARLY-OPTION (1-877-432-7596) www.earlyoptionpill.com

PRESCRIBER AGREEMENT FORM

Mifepristone Tablets, 200 mg

Mifepristone Tablets, 200 mg, is indicated, in a regimen with misoprostol, for the medical termination of intrauterine pregnancy through 70 days gestation. Please see Prescribing Information and Medication Guide for complete safety information.

To **become a certified prescriber**, you must:

- **If you submit mifepristone prescriptions for dispensing from certified pharmacies:**
 - o Submit this form to each certified pharmacy to which you intend to submit mifepristone prescriptions. The form must be received by the certified pharmacy before any prescriptions are dispensed by that pharmacy.
- **If you order mifepristone for dispensing by you or healthcare providers under your supervision:**
 - o Submit this form to the distributor. This form must be received by the distributor before the first order will be shipped to the healthcare setting.
 - o Healthcare settings, such as medical offices, clinics, and hospitals, where mifepristone will be dispensed by or under the supervision of a certified prescriber in the Mifepristone REMS Program do not require pharmacy certification.

Prescriber Agreement: By signing this form, you agree that you meet the qualifications below and will follow the guidelines for use. You are responsible for overseeing implementation and compliance with the Mifepristone REMS Program. You also understand that if the guidelines below are not followed, the distributor may stop shipping mifepristone to the locations that you identify and certified pharmacies may stop accepting your mifepristone prescriptions.

Mifepristone must be provided by or under the supervision of a certified prescriber who meets the following qualifications:

- Ability to assess the duration of pregnancy accurately.
- Ability to diagnose ectopic pregnancies.
- Ability to provide surgical intervention in cases of incomplete abortion or severe bleeding, or have made plans to provide such care through others, and be able to assure patient access to medical facilities equipped to provide blood transfusions and resuscitation, if necessary.
- Has read and understood the Prescribing Information for mifepristone. The Prescribing Information is available by calling 1-855-MIFE-INFO (1-855—643-3463 toll-free), or by visiting www.MifeInfo.com.

In addition to meeting these qualifications, you also agree to follow these guidelines for use:

- Ensure that the *Patient Agreement Form* is reviewed with the patient and the risks of the mifepristone treatment regimen are fully explained. Ensure any questions the patient may have prior to receiving mifepristone are answered.
- Ensure the healthcare provider and patient sign the *Patient Agreement Form*.
- Ensure that the patient is provided with a copy of the *Patient Agreement Form* and Medication Guide.
- Ensure that the signed *Patient Agreement Form* is placed in the patient's medical record.
- Ensure that any deaths of patients who received mifepristone are reported to GenBioPro, Inc. that provided the mifepristone, identifying the patient by a non-identifiable reference and including the NDC and lot number from the package of mifepristone that was dispensed to the patient.

Ensure that healthcare providers under your supervision follow the guidelines listed above.



GenBioPro Inc. - PO Box 32011 - Las Vegas, NV 89103
1-855-MIFE-INFO (1-855-643-3463) - www.MifeInfo.com

- If mifepristone will be dispensed through a certified pharmacy:
 - o Assess appropriateness of dispensing mifepristone when contacted by a certified pharmacy about patients who will receive mifepristone more than 4 calendar days after the prescription was received by the certified pharmacy.
 - o Obtain the NDC and lot number of the package of mifepristone the patient received in the event the prescriber becomes aware of the death of a patient.
- If mifepristone will be dispensed by you or by healthcare providers under your supervision:
 - o Ensure the NDC and lot number from each package of mifepristone are recorded in the patient's record.

I understand that a certified pharmacy may dispense mifepristone made by a different manufacturer than that stated on this Prescriber Agreement Form.

Print Name: _____ Title: _____

Signature: _____ Date: _____

Medical License # _____ State _____

NPI # _____

Practice Setting Address: _____

Return completed form to RxAgreements@GenBioPro.com or fax to 1-877-239-8036

Approved 01/2023 [Doc control ID]

PATIENT AGREEMENT FORM

Mifepristone Tablets, 200 mg

Healthcare Providers: *Counsel the patient on the risks of mifepristone. Both you and the patient must provide a written or electronic signature on this form.*

Patient Agreement:

1. I have decided to take mifepristone and misoprostol to end my pregnancy and will follow my healthcare provider's advice about when to take each drug and what to do in an emergency.
2. I understand:
 - a. I will take mifepristone on Day 1.
 - b. I will take the misoprostol tablets 24 to 48 hours after I take mifepristone.
3. My healthcare provider has talked with me about the risks, including:
 - heavy bleeding
 - infection
4. I will contact the clinic/office/provider right away if in the days after treatment I have:
 - a fever of 100.4°F or higher that lasts for more than four hours
 - heavy bleeding (soaking through two thick full-size sanitary pads per hour for two hours in a row)
 - severe stomach area (abdominal) pain or discomfort, or I am “feeling sick,” including weakness, nausea, vomiting, or diarrhea, more than 24 hours after taking misoprostol
— these symptoms may be a sign of a serious infection or another problem (including an ectopic pregnancy, a pregnancy outside the womb).

My healthcare provider has told me that these symptoms listed above could require emergency care. If I cannot reach the clinic/office/provider right away, my healthcare provider has told me who to call and what to do.
5. I should follow up with my healthcare provider about 7 to 14 days after I take mifepristone to be sure that my pregnancy has ended and that I am well.
6. I know that, in some cases, the treatment will not work. This happens in about 2 to 7 out of 100 women who use this treatment. If my pregnancy continues after treatment with mifepristone and misoprostol, I will talk with my provider about a surgical procedure to end my pregnancy.
7. If I need a surgical procedure because the medicines did not end my pregnancy or to stop heavy bleeding, my healthcare provider has told me whether they will do the procedure or refer me to another healthcare provider who will.
8. I have the MEDICATION GUIDE for mifepristone.
9. My healthcare provider has answered all my questions.

Patient Signature: _____ **Patient Name (print):** _____ **Date:** _____

Provider Signature: _____ **Provider Name (print):** _____ **Date:** _____

Patient Agreement Forms may be provided, completed, signed, and transmitted in paper or electronically.

01/2023

MIFEPREX®(Mifepristone) Tablets, 200mg

PHARMACY AGREEMENT FORM

Pharmacies must designate an authorized representative to carry out the certification process and oversee implementation and compliance with the Mifepristone REMS Program on behalf of the pharmacy.

Healthcare settings, such as medical offices, clinics, and hospitals, where mifepristone will be dispensed by or under the supervision of a certified prescriber in the Mifepristone REMS Program do not require pharmacy certification.

By signing this form, as the Authorized Representative I certify that:

- Each location of my pharmacy that will dispense Mifeprex is able to receive *Prescriber Agreement Forms* by email and fax.
- Each location of my pharmacy that will dispense Mifeprex is able to ship Mifeprex using a shipping service that provides tracking information.
- I have read and understood the Prescribing Information for Mifeprex. The Prescribing Information is available by calling 1-877-4 EARLY OPTION (1-877-432-7596 toll-free) or online at www.earlyoptionpill.com; and
- Each location of my pharmacy that will dispense Mifeprex will put processes and procedures in place to ensure the following requirements are completed. I also understand that if my pharmacy does not complete these requirements, the distributor may stop accepting Mifeprex orders.
 - o Verify that the prescriber is certified in the Mifepristone REMS Program by confirming their completed *Prescriber Agreement Form* was received with the prescription or is on file with your pharmacy.
 - o Dispense Mifeprex such that it is delivered to the patient within 4 calendar days of the date the pharmacy receives the prescription, except as provided in the following bullet.
 - o Confirm with the prescriber the appropriateness of dispensing Mifeprex for patients who will receive the drug more than 4 calendar days after the date the pharmacy receives the prescription and document the prescriber's decision.
 - o Record in the patient's record the NDC and lot number from each package of Mifeprex dispensed.
 - o Track and verify receipt of each shipment of Mifeprex.
 - o Dispense mifepristone in its package as supplied by Danco Laboratories, LLC.
 - o Report any patient deaths to the prescriber, including the NDC and lot number from the package of Mifeprex dispensed to the patient, and remind the prescriber of their obligation to report the deaths to Danco Laboratories, LLC. Notify Danco that your pharmacy submitted a report of death to the prescriber, including the name and contact information for the prescriber and the NDC and lot number of the dispensed product.
 - o Not distribute, transfer, loan or sell mifepristone except to certified prescribers or other locations of the pharmacy.
 - o Maintain records of *Prescriber Agreement Forms*, dispensing and shipping, and all processes and procedures including compliance with those processes and procedures.
 - o Maintain the identity of Mifeprex patients and prescribers as confidential and protected from disclosure except to the extent necessary for dispensing under this REMS or as necessary for payment and/or insurance.
 - o Train all relevant staff on the Mifepristone REMS Program requirements.
 - o Comply with audits carried out by the Mifepristone Sponsors or a third party acting on behalf of the Mifepristone Sponsors to ensure that all processes and procedures are in place and are being followed.

Any new authorized representative must complete and submit the *Pharmacy Agreement Form*.

Authorized Representative Name: _____ Title: _____



*MIFEPREX is a registered trademark of Danco Laboratories, LLC

P.O. Box 4816-New York, NY 10185

1-877-4-EARLY-OPTION (1-877-432-7596) www.earlyoptionpill.com

Signature: _____ Date: _____

Email: _____ Phone: _____ Preferred __ email __ phone

Pharmacy Name: _____

Pharmacy Address: _____

Return completed form to Mifeprex@dancodistributor.com or fax to 1-866-227-3343.



*MIFEPREX is a registered trademark of Danco Laboratories, LLC
P.O. Box 4816-New York, NY 10185
1-877-4-EARLY-OPTION (1-877-432-7596) www.earlyoptionpill.com

PHARMACY AGREEMENT FORM

Mifepristone Tablets, 200 mg

Pharmacies must designate an authorized representative to carry out the certification process and oversee implementation and compliance with the Mifepristone REMS Program on behalf of the pharmacy.

Healthcare settings, such as medical offices, clinics, and hospitals, where mifepristone will be dispensed by or under the supervision of a certified prescriber in the Mifepristone REMS Program do not require pharmacy certification.

By signing this form, as the Authorized Representative I certify that:

- Each location of my pharmacy that will dispense mifepristone is able to receive *Prescriber Agreement Forms* by email and fax.
- Each location of my pharmacy that will dispense mifepristone is able to ship mifepristone using a shipping service that provides tracking information.
- I have read and understood the Prescribing Information for mifepristone. The Prescribing Information is available by calling 1-855-MIFE-INFO (1-855-643-3463 toll-free) or online at www.MifeInfo.com; and
- Each location of my pharmacy that will dispense mifepristone will put processes and procedures in place to ensure the following requirements are completed. I also understand that if my pharmacy does not complete these requirements, the distributor may stop accepting mifepristone orders.
 - o Verify that the prescriber is certified in the Mifepristone REMS Program by confirming their completed *Prescriber Agreement Form* was received with the prescription or is on file with your pharmacy.
 - o Dispense mifepristone such that it is delivered to the patient within 4 calendar days of the date the pharmacy receives the prescription, except as provided in the following bullet.
 - o Confirm with the prescriber the appropriateness of dispensing mifepristone for patients who will receive the drug more than 4 calendar days after the date the pharmacy receives the prescription and document the prescriber's decision.
 - o Record in the patient's record the NDC and lot number from each package of mifepristone dispensed.
 - o Track and verify receipt of each shipment of mifepristone.
 - o Dispense mifepristone in its package as supplied by GenBioPro, Inc.
 - o Report any patient deaths to the prescriber, including the NDC and lot number from the package of mifepristone dispensed to the patient, and remind the prescriber of their obligation to report the deaths to GenBioPro, Inc. Notify GenBioPro that your pharmacy submitted a report of death to the prescriber, including the name and contact information for the prescriber and the NDC and lot number of the dispensed product.
 - o Not distribute, transfer, loan or sell mifepristone except to certified prescribers or other locations of the pharmacy.
 - o Maintain records of *Prescriber Agreement Forms*, dispensing and shipping, all processes and procedures including compliance with those processes and procedures.
 - o Maintain the identity of mifepristone patients and prescribers as confidential and protected from disclosure except to the extent necessary for dispensing under this REMS or as necessary for payment and/or insurance purposes.
 - o Train all relevant staff on the Mifepristone REMS Program requirements.
 - o Comply with audits carried out by the Mifepristone Sponsors or a third party acting on behalf of the Mifepristone Sponsors to ensure that all processes and procedures are in place and are being followed.

Any new authorized representative must complete and submit the *Pharmacy Agreement Form*.

Authorized Representative Name: _____ Title: _____

Signature: _____ Date: _____

Email: _____ Phone: _____ Preferred ___ email ___ phone

Pharmacy Name: _____

Pharmacy Address: _____

Return completed form to RxAgreements@GenBioPro.com or fax to 1-877-239-8036.



GenBioPro Inc. - PO Box 32011 - Las Vegas, NV 89103
1-855-MIFE-INFO (1-855-643-3463) - www.MifeInfo.com

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

020687Orig1s025

SUMMARY REVIEW

(b) (6) and (b) (6)
(b) (6)
(b) (6)
Center for Drug Evaluation and Research (CDER)

Application Type	NDA and ANDA
Application Number	NDA 020687 and ANDA 091178
Supplement Number, Date Received	NDA Supplement-025 and ANDA Supplement-004 received June 22, 2022 (sequences 18 and 87 respectively) and amended October 19, 2022 (sequences 22 and 91 respectively), November 30, 2022 (sequences 24 and 92 respectively), December 9, 2022 (sequences 25 and 93 respectively) and December 16, 2022 (sequences 26 and 95 respectively). This supplement is on a 180-Day clock.
Targeted Action Date	December 19, 2022
(b) (6) #	2022-1169
Reviewer Names	(b) (6) (b) (6) (b) (6)
(b) (6)	(b) (6) (b) (6)
(b) (6)	(b) (6)
(b) (6)	(b) (6)
(b) (6)	(b) (6)
Review Completion Date	January 3, 2023
Subject	Review of proposed Major REMS Modification
Established Name	Mifepristone REMS
Name of Sponsor	Danco Laboratories, LLC and GenBioPro, Inc.
Therapeutic Class	Progestin antagonist
Formulation	Oral tablet

TABLE OF CONTENTS

EXECUTIVE SUMMARY	3
1. Introduction	4
2. Background	4
2.1. Product Information and REMS Information	4
2.2. Regulatory History	6
3. Review of Proposed REMS Modification	8
3.1. REMS Goal	8
3.2. REMS Document	8
3.3. REMS Requirements	9
3.3.1. Addition and Removal of ETASU	9
3.3.2. REMS Participant Requirements and Materials	9
3.3.2.1. Prescriber Requirements	9
3.3.2.2. Patient Requirements	11
3.3.2.3. Pharmacy Requirements	12
3.3.2.4. Distributor Requirements	15
3.3.3. REMS Sponsor Requirements	15
3.3.3.1. Sponsor Requirements to Support Prescriber Certification	15
3.3.3.2. Sponsor Requirements to Support Pharmacy Certification	16
3.3.3.3. Sponsor Implementation Requirements	16
3.4. REMS Assessment Timetable	16
4. Supporting Document	17
5. REMS Assessment Plan	17
6. Discussion	19
7. Conclusions and Recommendations	21
8. References	21
9. Appendices	22

EXECUTIVE SUMMARY

This is a review of the proposed modification to the single, shared system Risk Evaluation and Mitigation Strategy (REMS) for mifepristone 200 mg (hereafter referred to as the Mifepristone REMS Program) submitted by Danco Laboratories, LLC (Danco) for new drug application (NDA) 020687 and by GenBioPro, Inc. (GBP) for abbreviated new drug application (ANDA) 091178. The Sponsors submitted proposed modification to the Mifepristone REMS Program on June 22, 2022, and amended their submissions on October 19, 2022 (Danco), October 20, 2022 (GBP), November 30, 2022 (both), December 9, 2022 (both) and December 16, 2022 (both).

The Mifepristone REMS Program was originally approved on April 11, 2019, to mitigate the risk of serious complications associated with mifepristone 200 mg. The most recent REMS modification was approved on May 14, 2021.^a The Mifepristone REMS Program consists of elements to assure safe use (ETASU) A, C and D, an implementation system, and a timetable for submission of assessments of the REMS.

The Sponsors submitted the proposed modification to the REMS in response to the Agency's REMS Modification Notification letters dated December 16, 2021, which required removal of the requirement that mifepristone be dispensed only in certain healthcare settings, specifically clinics, medical offices, and hospitals (i.e., the "in-person dispensing requirement") and the addition of certification of pharmacies that dispense the drug.

In addition, the following were addressed during the course of the review:

- revisions to the REMS goal to align with the updated REMS requirements.
- replacing serial number with recording of NDC and lot number of mifepristone dispensed.
- additional edits for clarification and consistency in the REMS Document and REMS materials (*Prescriber Agreement Forms, Patient Agreement Form, and Pharmacy Agreement Forms*).

The review team finds the proposed modification to the Mifepristone REMS Program last submitted on December 16, 2022, to be acceptable and recommends approval of the REMS modification. The proposed REMS modification includes changes to the REMS goal, additional REMS requirements for prescribers to incorporate dispensing from certified pharmacies and new REMS requirements for pharmacy certification.

The proposed goal of the modified REMS for mifepristone 200 mg is to mitigate the risk of serious complications associated with mifepristone by:

- a) Requiring healthcare providers who prescribe mifepristone to be certified in the Mifepristone REMS Program.
- b) Ensuring that mifepristone is only dispensed by or under the supervision of certified prescribers, or by certified pharmacies on prescriptions issued by certified prescribers.
- c) Informing patients about the risk of serious complications associated with mifepristone.

^a The May 14, 2021 REMS modification approved the inclusion of gender neutral language in the Patient Agreement Form as well as corresponding minor changes to the REMS document to be consistent with the changes made to the Patient Agreement Form.

The timetable for submission of assessments of the REMS was modified to one year from the date of the approval of the modified REMS and annually thereafter. The assessment plan was revised to align with the changes to the REMS and capture additional metrics for drug utilization and REMS operations.

The modified REMS includes ETASU A, B and D, an implementation system, and a timetable for submission of assessments of the REMS. Mifepristone will no longer be required to be dispensed only in certain healthcare settings, specifically clinics, medical offices, and hospitals (referred to as the “in-person dispensing requirement” for brevity) and will be able to be dispensed from certified pharmacies.

1. Introduction

This review evaluates the proposed modification to the single, shared system Risk Evaluation and Mitigation Strategy (REMS) for mifepristone 200 mg (hereafter referred to as the Mifepristone REMS Program) submitted by Danco Laboratories, LLC (Danco) for new drug application (NDA) 020687 and by GenBioPro, Inc. (GBP) for abbreviated new drug application (ANDA) 091178.

The Sponsors initially submitted proposed modification to the Mifepristone REMS Program on June 22, 2022, in response to the Agency’s REMS Modification Notification letters issued on December 16, 2021, to Danco and GBP, requiring the following modification to minimize the burden on the healthcare delivery system of complying with the REMS and to ensure that the benefits of the drug outweigh the risks:

- removal of the requirement that mifepristone be dispensed only in certain healthcare settings, specifically clinics, medical offices, and hospitals (i.e., the “in-person dispensing requirement”)
- addition of certification of pharmacies that dispense the drug

Per the Agency’s December 16, 2021, REMS Modification Notification letters, the proposed REMS was required to include the following ETASU to mitigate the risk of serious complications associated with mifepristone, including at least the following:

- healthcare providers have particular experience or training, or are specially certified
- pharmacies, practitioners, or health care settings that dispense the drug are specially certified
- the drug is dispensed to patients with evidence or other documentation of safe use conditions

The REMS was also required to include an implementation system and timetable for submission of assessments.

2. Background

2.1. Product Information and REMS Information

Mifepristone is a progestin antagonist indicated, in a regimen with misoprostol, for the medical termination of intrauterine pregnancy (IUP) through 70 days gestation. Mifepristone is available as 200 mg tablets for oral use.

Mifeprex (mifepristone) was approved on September 28, 2000, with a restricted distribution program under 21 CFR 314.520 (subpart H)^b to ensure that the benefits of the drug outweighed

^b NDA approval letter Mifeprex (NDA 020687) dated September 28, 2000.

the risk of serious complications associated with mifepristone when used for medical abortion.^c Mifeprex was deemed to have in effect an approved REMS under section 505-1 of the Federal Food, Drug, and Cosmetic Act with the passage of the Food and Drug Administration Amendments Act of 2007 (FDAAA), and the Mifeprex REMS was approved on June 8, 2011.

On March 29, 2016, FDA approved an efficacy supplement for Mifeprex, which included changes in the dose of Mifeprex and the dosing regimen for taking Mifeprex and misoprostol, as well as a modification of the gestational age up to which Mifeprex has been shown to be safe and effective and a modification to the process for follow-up after administration of the drug. FDA also approved modification to the Mifeprex REMS that reflected the changes approved in the efficacy supplement.¹⁻⁵ On April 11, 2019, FDA approved ANDA 091178 and the Mifepristone REMS Program.⁶⁻⁷ The Mifepristone REMS Program is a single, shared system REMS that includes NDA 020687 and ANDA 091178. The goal of the approved Mifepristone REMS Program is to mitigate the risk of serious complications associated with mifepristone by:

- a) Requiring healthcare providers who prescribe mifepristone to be certified in the Mifepristone REMS Program (under ETASU A).
- b) Ensuring that mifepristone is only dispensed in certain healthcare settings by or under the supervision of a certified prescriber (under ETASU C).
- c) Informing patients about the risk of serious complications associated with mifepristone (under ETASU D).

The Mifepristone REMS Program was last modified and approved in 2021 to revise the *Patient Agreement Form* to include gender-neutral language; however, the goal of the Mifepristone REMS Program has not changed since the initial approval in 2019.

Under ETASU A, to become specially certified to prescribe mifepristone, a healthcare provider must review the prescribing information, complete and sign the *Prescriber Agreement Form*, and agree to follow the guidelines for use of mifepristone. Under ETASU C, in the Mifepristone REMS Program as approved prior to today's action, mifepristone was required to be dispensed to patients only in certain healthcare settings, specifically clinics, medical offices, and hospitals, by or under the supervision of a certified prescriber. Under ETASU D, mifepristone must be dispensed to patients with evidence or other documentation of safe use conditions (i.e., the patient must sign a *Patient Agreement Form*). The approved Mifepristone REMS Program includes an implementation system, and a timetable for assessments (one year from the date of the initial approval of the REMS on April 11, 2019, and every three years thereafter).

In April 2021, FDA communicated its intent to exercise enforcement discretion during the COVID-19 public health emergency (PHE) regarding the in-person dispensing requirement in the Mifepristone REMS Program. Specifically, FDA communicated that provided all other requirements of the Mifepristone REMS Program are met, the Agency intended to exercise enforcement discretion with respect to the in-person dispensing requirement of the Mifepristone REMS Program, including any in-person requirements that may be related to the *Patient Agreement Form*, during the COVID-19 PHE. This determination, which FDA made on April 12, 2021, was effective immediately. We also note that from July 13, 2020, to January 12, 2021, per a court order, FDA was enjoined from enforcing the in-person dispensing requirement of the Mifepristone REMS Program.⁸

^c Mifepristone is also approved in approximately 80 other countries.
https://gynuity.org/assets/resources/biblio_ref_lst_mife_en.pdf

Further, and as we also communicated on April 12, 2021, to the extent all of the other requirements of the Mifepristone REMS Program are met, the Agency intended to exercise enforcement discretion during the COVID-19 PHE with respect to the dispensing of Mifeprex or the approved generic version of Mifeprex, Mifepristone Tablets, 200 mg, through the mail, either by or under the supervision of a certified prescriber, or through a mail-order pharmacy when such dispensing is done under the supervision of a certified prescriber.

2.2. Regulatory History

The following is a summary of the regulatory history relevant to this review:

- 04/11/2019: Approval of the Mifepristone REMS Program, a single, shared system REMS that includes NDA 020687 and ANDA 091178.
- 04/12/2021: The Agency issued a General Advice letter to both the NDA and ANDA Applicants, explaining that FDA intended to exercise enforcement discretion during the COVID-19 PHE with respect to the in-person dispensing requirement in the Mifepristone REMS Program, including any in-person requirements that may be related to the Patient Agreement Form.
- 05/07/2021: The Agency stated that it would be reviewing the elements of the Mifepristone REMS Program in accordance with section 505-1 of the FD&C Act.
- 12/16/2021: The Agency completed its review of the Mifepristone REMS Program and determined, among other things, that the REMS must be modified to remove the in-person dispensing requirement and add pharmacy certification.⁹
- 12/16/2021: REMS Modification Notification letters were sent to both Sponsors stating that the approved Mifepristone REMS Program must be modified to minimize the burden on the healthcare system of complying with the REMS and ensure that the benefits of the drug outweigh the risks.
- 04/08/2022: Final written responses to a Type A meeting request were provided to Danco, the point of contact for the Mifepristone REMS Program. The questions pertained to the 12/16/2021 REMS Modification Notification letter requirements.
- 04/13/2022: The Sponsors requested an extension to 6/30/2022, to submit a proposed REMS modification in response to the Agency's 12/16/2021 REMS Modification Notification letters.
- 04/15/2022: The Agency granted the Sponsors' request for an extension to submit a proposed REMS modification and conveyed that the modification must be submitted no later than 06/30/2022.¹⁰
- 06/22/2022: Danco and GBP submitted a proposed REMS modification to their respective applications in response to the 12/16/2021 REMS Modification Notification letters.
- 07/22/2022: An Information Request was sent to the Sponsors requesting clarification of the proposed prescriber and dispenser requirements and additional rationale to support their proposal.
- 08/26/2022: Sponsors submitted responses to 07/22/2022 Information Request.
- 09/19/2022: Teleconference was held between Agency and Sponsors where the Agency communicated the REMS requirements that are necessary to support the addition of pharmacy

certification. The Agency proposed focusing on the pharmacy settings where a closed system^d REMS could be implemented using the existing email and facsimile based system, (b) (4), as the best strategy for an approvable modification by the goal date.

- 09/22/2022: An Information Request was sent to Sponsors requesting confirmation that the Sponsors agree with the pharmacy distribution approach outlined in the 09/19/2022 teleconference so that the Agency's feedback could be appropriately tailored.
- 09/23/2022: The Sponsors confirmed via email that they were willing to pursue (b) (4), as discussed in the 09/19/2022 teleconference. The Sponsors also requested a teleconference to discuss the current modification (b) (4).
- 09/27/2022: Comments from the 09/19/2022 teleconference sent to Sponsors with additional comments and requests regarding what will be necessary for pharmacy certification.
- 09/29/2022: An Information request was sent to the Sponsors asking for agenda items, questions, and a request to walk through their proposed system for pharmacy certification, including dispensing through mail-order or specialty pharmacies, at the 10/06/2022 scheduled teleconference.
- 10/04/2022: Sponsors emailed that they will focus the 10/06/2022 teleconference on the 09/27/2022 Agency comments and their mail order and specialty pharmacy distribution model.
- 10/06/2022: Teleconference was held between Agency and Sponsors where Sponsors outlined their proposal for pharmacy certification, including dispensing through mail order and specialty pharmacies, as well as their concerns with certain requirements and general timelines.
- 10/19/2022: Danco submitted a REMS amendment to their pending sNDA, which included a REMS document and REMS materials. They did not submit a REMS Supporting Document.
- 10/20/2022: GBP submitted a REMS amendment to their pending sANDA, which included a REMS document and REMS materials. They did not submit a REMS Supporting Document.
- 10/25/2022: Teleconference was held between Agency and Sponsors to discuss the *Patient Agreement Form* and timing related to shipping a mifepristone prescription from a certified pharmacy to the patient.
- 11/23/2022: An Information Request was sent to Sponsors with comments on their proposed REMS Document, submitted on 10/19/2022 (Danco) and 10/20/2022 (GBP).
- 11/30/2022: Danco and GBP submitted REMS amendments, which included the REMS Document, to their respective pending supplemental applications.
- 12/01/2022: Teleconference was held between Agency and Sponsors to discuss the REMS Document.
- 12/05/2022: An Information Request was sent to Sponsors with comments on their proposed REMS Document submitted on 11/30/2022 and discussed at the teleconference on 12/01/2022, and REMS materials submitted to their applications on 10/19/2022 and 10/20/2022.

^d "Closed system" in this case refers to a system where prescribers, pharmacies, and distributors are certified or authorized in the REMS and the certification of the stakeholder must be verified prior to distribution or dispensing, as per the REMS.

- 12/07/2022: Teleconference was held between Agency and Sponsors to discuss the REMS Document and REMS materials the Agency sent to the Sponsors on 12/05/22.
- 12/08/2022: Danco and GBP submitted REMS amendments, including the REMS Document, *Prescriber Agreement Form*, *Pharmacy Agreement Form*, *Patient Agreement Form* and REMS Supporting Document, to their respective pending applications.
- 12/09/2022: An Information Request was sent to Sponsors with the Agency's comments on the REMS assessment plan.
- 12/14/2022: An Information Request was sent to Sponsors with the Agency's comments on the REMS Document, *Prescriber Agreement Form*, *Pharmacy Agreement Form*, and REMS Supporting Document.
- 12/15/2022: Two teleconferences were held between Agency and Sponsors to discuss the proposed REMS Document and REMS materials the Agency sent to the Sponsors on 12/14/22.
- 12/16/2022: Sponsors submitted a REMS amendment to their respective applications.

3. Review of Proposed REMS Modification

(b) (6) has discussed the Sponsors' proposed modification with the review team, which includes members of the (b) (6) and the (b) (6); hereafter referred to as the review team. This review includes their input and concurrence with the analysis and proposed changes to the Mifepristone REMS Program.

3.1. REMS Goal

The Sponsors proposed modification to the goal for the Mifepristone REMS Program to add that mifepristone can also be dispensed from certified pharmacies on prescriptions issued by certified prescribers. The proposed REMS goal is:

The goal of the REMS for mifepristone is to mitigate the risk of serious complications associated with mifepristone by:

- a) Requiring healthcare providers who prescribe mifepristone to be certified in the Mifepristone REMS Program.
- b) Ensuring that mifepristone is only dispensed by or under the supervision of certified prescribers, or by certified pharmacies on prescriptions issued by certified prescribers.
- c) Informing patients about the risk of serious complications associated with mifepristone.

Reviewer Comment: *We agree with the Sponsors' proposal.*

3.2. REMS Document

The proposed REMS Document is not in the format as outlined in the 2017 Draft Guidance for Industry, Format and Content of a REMS Document.¹¹

Reviewer Comment: To avoid the misperception that this REMS modification is making major changes to the REMS document that go beyond our December 16, 2021, determination that the REMS must be modified to remove the in-person dispensing requirement and add pharmacy certification, CDER staff and management discussed whether to change the format of the REMS document to that described in the 2017 draft guidance.¹¹ After internal discussion, CDER staff and management aligned not to transition the REMS document at this time to the format described in the 2017 draft guidance.

3.3. REMS Requirements

3.3.1. Addition and Removal of ETASU

The December 16, 2021, REMS Modification Notification letters specified that the ETASU must be modified to minimize the burden on the healthcare delivery system of complying with the REMS and to ensure the benefits of the drug outweigh the risks by:

- Removing the requirement that mifepristone be dispensed only in certain healthcare settings, specifically clinics, medical offices and hospitals (i.e., the “in-person dispensing requirement”), and;
- Adding a requirement that pharmacies that dispense the drug be specially certified.

The Sponsors proposed changes to the REMS as reflected in the subsections below.

3.3.2. REMS Participant Requirements and Materials

3.3.2.1. Prescriber Requirements

Consistent with the approved Mifepristone REMS Program prescribers must be specially certified. To become specially certified to prescribe mifepristone, healthcare providers who prescribe must review the Prescribing Information for mifepristone and complete the *Prescriber Agreement Form*. In signing the *Prescriber Agreement Form*, prescribers agree they meet certain qualifications and will follow the guidelines for use of mifepristone. The guidelines for use include ensuring i) that the *Patient Agreement Form* is reviewed with the patient and the risks of the mifepristone treatment regimen are fully explained; ii) that the healthcare provider (HCP) and the patient sign the *Patient Agreement Form*, iii) the patient receives a copy of the *Patient Agreement Form* and Medication Guide, iv) the *Patient Agreement Form* is placed in the patient’s medical record; v) that any patient deaths are reported to the Mifepristone Sponsor that provided the mifepristone, identifying the patient by a non-identifiable reference and including the NDC and lot number from the package of mifepristone that was dispensed to the patient. The language on the guidelines for use was revised from the Mifepristone REMS Program approved in 2021 to clarify that, if the certified prescriber supervises the dispensing of mifepristone, they must ensure the guidelines for use of mifepristone are followed by those under their supervision. This clarification reflects the ongoing implementation of the approved Mifepristone REMS Program. For example, consistent with the approved REMS, the *Patient Agreement Form* does not require the certified prescriber’s signature, but rather the signature of the healthcare provider counseling the patient on the risks of mifepristone. Additional changes were made globally to provide consistency and clarity of the requirements for certified prescribers and healthcare providers who complete tasks under the supervision of certified prescribers.

A certified prescriber may submit the *Prescriber Agreement Form* to an authorized distributor if the certified prescriber wishes to dispense or supervise the dispensing of mifepristone; this is consistent with the current requirements of the Mifepristone REMS Program. Additional requirements were

added to incorporate mifepristone dispensing by a certified pharmacy. If a healthcare provider wishes to prescribe mifepristone by sending a prescription to a certified pharmacy for dispensing, the healthcare provider must become certified by providing the pharmacy a *Prescriber Agreement Form* signed by the provider. A certified prescriber must also assess the appropriateness of dispensing mifepristone when contacted by a certified pharmacy about patients who will receive mifepristone more than four calendar days after the prescription was received by the certified pharmacy.

The NDC and lot number of the dispensed drug will be recorded in the patient's record when mifepristone is dispensed by or under the supervision of a certified prescriber, replacing the requirement that serial numbers from each package of mifepristone be recorded in the patient's record. If prescribers become aware of the death of a patient for whom the mifepristone was dispensed from a certified pharmacy, the prescribers will be required to obtain the NDC and lot number of the package of mifepristone the patient received from the pharmacy.

The following materials support prescriber requirements:

- *Prescriber Agreement Form* for Danco Laboratories, LLC
- *Prescriber Agreement Form* for GenBioPro, Inc.
- *Patient Agreement Form*

Reviewer Comment: *We agree with the Sponsors' proposal.*

Although certain activities (review of the Patient Agreement Form with patients and answering any questions about treatment, signing, providing a copy to the patient and retaining the Patient Agreement Form, providing a copy of the Medication Guide, and ensuring any deaths are reported to the Mifepristone Sponsor, recording the NDC and lot number from drug dispensed from the certified prescriber or those under their supervision) may be conducted by healthcare providers under the supervision of a certified prescriber, the certified prescriber remains responsible for ensuring compliance with the requirements of the Mifepristone REMS Program. We agree with the additional language to further clarify that the certified prescriber must ensure the guidelines for use of mifepristone are followed.

As proposed, certified prescribers may either, 1) continue to submit the Prescriber Agreement Form to an authorized distributor if the certified prescriber is dispensing or supervising the dispensing of the drug (as already required in the REMS), or 2) if the drug will be dispensed from a certified pharmacy, submit the Prescriber Agreement Form to the certified pharmacy that will dispense the drug (as proposed in the modification). Regarding #2, the pharmacy can only fill prescriptions written by a certified prescriber.

Based on our review of the proposed changes, the review team finds it acceptable for prescribers to submit their Prescriber Agreement Form directly to the certified pharmacy. Although certified prescribers still have the option of in-person dispensing of the drug, not all prescribers may want to stock mifepristone. Typically due to the number of drugs that are available and the expense associated with stocking prescription medications intended for outpatient use, most prescribers do not stock many medications, if they stock medications at all.

The proposal to submit a Prescriber Agreement Form to a certified pharmacy provides another option for dispensing mifepristone. The burden of providing the Prescriber Agreement Form prior to or when the prescription is provided to a certified pharmacy does not create unreasonable burden for prescribers. The burden of prescriber certification has been minimized to the extent possible. The Prescriber Agreement Form is designed to require minimal time to complete and requires that the prescriber submit it to the authorized distributor once, and if the prescriber chooses to use a certified pharmacy to dispense mifepristone, they will need to submit the form to the certified pharmacy.

There is an additional requirement added for certified pharmacies and certified prescribers in the event that a patient will not receive their medication from the certified pharmacy within four calendar days of the pharmacy's receipt of the prescription (for example, if the medication is not in stock). In this circumstance, the pharmacy will be required to contact the certified prescriber to make them aware of the delay and will be required to obtain from the prescriber confirmation that it is appropriate to dispense mifepristone to the patient even though they will receive mifepristone more than four calendar days after the prescription was received by the certified pharmacy. This confirmation is intended to ensure timeliness of delivery in light of the labeled indication and gestational age. Additional details and rationale on the pharmacy requirements to dispense and ship drug in a timely manner are described in section 3.3.2.3.

If a certified prescriber becomes aware of a patient death that occurs subsequent to the use of mifepristone dispensed from a pharmacy, the certified prescriber must obtain the NDC and lot number of the package of mifepristone the patient received from the pharmacy. This information will be reported to the appropriate Mifepristone Sponsor in the same manner prescribers have done previously. This additional requirement to obtain the NDC and lot number from the pharmacy is needed to ensure consistent adverse event reporting when mifepristone is dispensed from a certified pharmacy.

Prescriber Agreement Form

The Sponsors' proposed changes to the *Prescriber Agreement Form* aligned with those described above. The proposed *Prescriber Agreement Form* explains the two methods of certification which are: 1) submitting the form to the authorized distributor and 2) submitting the form to the dispensing certified pharmacy. Further clarification was added that healthcare settings, such as medical offices, clinics, and hospitals, where mifepristone will be dispensed by or under the supervision of a certified prescriber in the Mifepristone REMS Program do not require pharmacy certification. The statement that certified prescribers are responsible for overseeing implementation and compliance with the REMS Program was also added. The following statement was added to the form: "I understand that the pharmacy may dispense mifepristone made by a different manufacturer than that stated on the Prescriber Agreement Form." The account set up information was removed and replaced with prescriber information response fields.

Reviewer Comment: *We agree with the Sponsors' proposal. Changes in the above prescriber requirements were incorporated in the Prescriber Agreement Form.*

3.3.2.2. Patient Requirements

The *Patient Agreement Form* was updated to clarify that the signatures may be written or electronic, to reorganize the risk information about ectopic pregnancy, and to remove the statement that the Medication Guide will be taken to an emergency room or provided to a healthcare provider who did not prescribe mifepristone so that it is known that the patient had a medical abortion with mifepristone.

The following materials support patient requirements:

- *Patient Agreement Form*

Reviewer Comment: *We agree with the Sponsors' proposal.*

The Patient Agreement Form continues to be an important part of standardizing the medication information on the use of mifepristone that prescribers communicate to their patients, and also provides the information in a brief and understandable format for patients. The requirement to counsel the

patient, to provide the patient with the Patient Agreement Form, and to have the healthcare provider and patient sign the Patient Agreement Form, ensures that each provider, including new providers, informs each patient of the appropriate use of mifepristone, risks associated with treatment, and what to do if the patient experiences symptoms that may require emergency care. The form is signed by the patient and the provider and placed in the patient's medical record, and a copy is provided to the patient, to document the patient's acknowledgment of receiving the information from the prescriber. The Agency agrees that the further clarification that signatures can be written or electronic is appropriate for the continued use of the form.

The reference to ectopic pregnancy has been reorganized in the document since it is not a risk of the drug. The signs and symptoms of an untreated ectopic pregnancy that may persist after mifepristone use have been clarified in the section of the form that explains the signs and symptoms of potential problems that may occur after mifepristone use.

The review team agrees with removing the patient's agreement to take the Medication Guide with them if they visit an emergency room or HCP who did not give them mifepristone so the emergency room or HCP will understand that the patient is having a medical abortion. Although this statement has been in the Medication Guide for a number of years, upon further consideration, the Agency has concluded that patients seeking emergency medical care are not likely to carry a Medication Guide with them, the Medication Guide is readily available online, and information about medical conditions and previous treatments can be obtained at the point of care.

3.3.2.3. Pharmacy Requirements

The Sponsors proposed that certified pharmacies, in addition to certified prescribers and HCPs under the supervision of certified prescribers, can dispense mifepristone. In order for a pharmacy to become certified, the pharmacy must designate an authorized representative to carry out the certification process and oversee implementation and compliance with the Mifepristone REMS Program on behalf of the pharmacy. The Authorized Representative must certify that they have read and understood the Prescribing Information for mifepristone. Each location of the pharmacy must be able to receive *Prescriber Agreement Forms* by email and fax and be able to ship mifepristone using a shipping service that provides tracking information.

Additionally, each dispensing pharmacy location must put processes and procedures in place to fulfill the REMS requirements. Certified pharmacies must verify prescriber certification by confirming they have obtained a copy of the prescriber's signed *Prescriber Agreement Form* before dispensing. Certified pharmacies must dispense mifepristone such that it is received by the patient within four days from the day of prescription receipt by the pharmacy. If the pharmacy will not be able to deliver mifepristone to the patient within four days of receipt of the prescription, the pharmacy must contact the prescriber to confirm the appropriateness of dispensing mifepristone and document the certified prescriber's decision. The pharmacy must also record the NDC and lot number from each package of mifepristone dispensed in the patient's record, track and verify receipt of each shipment of mifepristone, dispense mifepristone in its original package, and only distribute, transfer, loan, or sell mifepristone to certified prescribers or between locations of the certified pharmacy. The pharmacy must also report any patient deaths to the prescriber, including the NDC and lot number from the package dispensed to the patient, and remind the prescriber of their obligation under the REMS to report patient deaths to the Sponsor that supplied the mifepristone; the certified pharmacy also must notify the Sponsor that supplied the mifepristone that the pharmacy submitted a report of a patient death to the prescriber and include the name and contact information for the prescriber as well as the NDC and lot number of the dispensed

product. Record-keeping requirements of the pharmacy include records of *Prescriber Agreement Forms*, mifepristone dispensing and shipping, and all processes and procedures and compliance with those processes and procedures. Pharmacies must train all relevant staff and participate in compliance audits. Pharmacies must also maintain the identity of patients and providers as confidential, including limiting access to patient and provider identity only to those personnel necessary to dispense mifepristone in accordance with the Mifepristone REMS Program requirements, or as necessary for payment and/or insurance purposes. The requirement that mifepristone not be dispensed from retail pharmacies was removed.

The following materials support pharmacy requirements:

- *Pharmacy Agreement Form* for Danco Laboratories, LLC
- *Pharmacy Agreement Form* for GenBioPro, Inc.

Reviewer Comment: *We agree with the Sponsors' proposal. The Mifepristone REMS Program continues to require that mifepristone be prescribed only by certified prescribers. With the removal of the in-person dispensing requirement, however, mifepristone can be dispensed from a pharmacy, provided the product is prescribed by a certified prescriber and all other requirements of the REMS are met. Given this modification to the dispensing requirements in the REMS, it is necessary to add a requirement for certification of pharmacies. Adding the pharmacy certification requirement incorporates pharmacies into the REMS, ensures that pharmacies are aware of and agree to follow applicable REMS requirements, and ensures that mifepristone is only dispensed pursuant to prescriptions that are written by certified prescribers. Without pharmacy certification, a pharmacy might dispense product that was not prescribed by a certified prescriber. Adding pharmacy certification ensures that the prescriber is certified prior to dispensing the product to a patient; certified prescribers, in turn, have agreed to meet all the conditions of the REMS, including ensuring that the Patient Agreement Form is completed. In addition, wholesalers and distributors can only ship to certified pharmacies. Based on our review and our consideration of the distribution model implemented by the Sponsors during the periods when the in-person dispensing requirement was not being enforced, as well as REMS assessment data and published literature, we conclude that provided all other requirements of the REMS are met, the REMS program, with the removal of the in-person dispensing requirement and the addition of a requirement for pharmacy certification, will continue to ensure the benefits of mifepristone for medical abortion outweigh the risks while minimizing the burden imposed by the REMS on healthcare providers and patients.*

The requirement to maintain confidentiality, including limiting access to patient and provider identity only to those personnel necessary for dispensing under the Mifepristone REMS Program or as necessary for payment and/or insurance purposes, is included to avoid unduly burdening patient access.

The Sponsors proposed inclusion of this requirement because of concerns that patients may be reluctant or unwilling to seek to obtain mifepristone from pharmacies if they are concerned that confidentiality of their medical information could be compromised, potentially exposing them to intimidation, threats, or acts of violence by individuals opposed to the use of mifepristone for medical abortion.^e Further, unwillingness on the part of prescribers to participate in the Mifepristone REMS Program on the basis of

^e See e.g., 2020 Violence and Disruption Statistics, National Abortion Federation (Dec. 16, 2021), <https://prochoice.org/national-abortion-federation-releases-2020-violence-disruption-statistics/>; Amanda Musa, CNN, *Wyoming Authorities Search for a Suspect Believed to Have Set an Abortion Clinic on Fire*, CNN WIRE (June 10, 2022), <https://abc17news.com/news/2022/06/10/wyoming-authorities-search-for-a-suspect-believed-to-have-set-an-abortion-clinic-on-fire/>.

similar confidentiality concerns may unduly burden patient access by limiting the number of prescribers who are willing to send prescriptions to certified pharmacies. Addition of this requirement protects patient access by requiring the pharmacy to put processes and procedures in place to limit access to confidential information to only those individuals who are essential for dispensing mifepristone under the Mifepristone REMS Program or as necessary for payment or insurance purposes. Inclusion of this requirement for certified pharmacies is consistent with the requirement in the current Mifepristone REMS Program, that distributors maintain secure and confidential records.

Reference to mifepristone not being available in retail pharmacies was removed from the REMS. There is no single definition of the term "retail pharmacy" and therefore the scope of the exclusion in the REMS was not well defined. Including a restriction in the Mifepristone REMS Program that retail pharmacies cannot participate in the REMS may unintentionally prohibit the participation of mail order and specialty pharmacies that could, under one or more definitions, also be considered a "retail pharmacy."

After reconsideration of the term, "retail," the Agency concluded that a more appropriate approach was to articulate the specific requirements that would be necessary for pharmacy certification. As modified, the REMS will not preclude the participation of any pharmacy that meets the certification requirements. However, we acknowledge that the provision in the REMS related to pharmacies' verification of prescriber enrollment will likely limit the types of pharmacies that will choose to certify in the REMS. The REMS requires that pharmacies dispense mifepristone only after verifying that the prescriber is certified. The REMS further requires that pharmacies be able to receive the Prescriber Agreement Forms by email and fax.

(b) (4)

The pharmacy certification requirements include that the drug reach patients within four days of the certified pharmacy receiving the prescription. During the course of the review, the review team concluded that requiring medication delivery to the patient within four days of the pharmacy's receipt of a prescription is acceptable based on the labeled indication and literature,¹³ while taking into account practical shipping considerations (e.g., shipping over weekends and holidays). For patients who will not receive the drug within four calendar days of the date the pharmacy receives the prescription, the pharmacy must notify the certified prescriber and the certified prescriber must determine if it is still appropriate for the certified pharmacy to dispense the drug. The pharmacy must document the certified prescriber's decision. A prescriber's confirmation that it is appropriate to dispense mifepristone when it will not be delivered to the patient within the allotted four days is intended to ensure timeliness of delivery in light of the labeled indication and gestational age.

Pharmacy Agreement Form

The proposed *Pharmacy Agreement Form* is a new form and is the means by which a pharmacy becomes certified to dispense mifepristone. The form, which is submitted by an authorized representative on behalf of a pharmacy seeking certification, outlines all requirements proposed above. Clarification is included in the form that healthcare settings, such as medical offices, clinics, and hospitals, where mifepristone will be dispensed by or under the supervision of a certified prescriber in the Mifepristone REMS Program, do not require pharmacy certification. Any new authorized representative must complete and submit the *Pharmacy Agreement Form*. Spaces for specific authorized representative information and pharmacy name and address are included. The completed form can be submitted by email or fax to the authorized distributor.

Reviewer Comment: *We agree with the Sponsors' proposal. The Pharmacy Agreement Form aligns with the pharmacy requirements discussed above.*

3.3.2.4. Distributor Requirements

The Sponsors proposed that the distributors' processes and procedures in the approved Mifepristone REMS Program be updated to ensure that mifepristone is only shipped to clinics, medical offices and hospitals identified by certified prescribers and to certified pharmacies. Distributors will continue to complete the certification process for any *Prescriber Agreement Forms* they receive and also will complete the certification process for pharmacies upon receipt of a *Pharmacy Agreement Form*, including notifying pharmacies when they become certified. FDA was removed as a potential auditor for distributors.

Reviewer Comment: *We agree with the Sponsors' proposal. At this time, FDA does not audit distributors directly, it carries out inspections of Sponsors to monitor industry compliance with REMS requirements.*

3.3.3. REMS Sponsor Requirements

3.3.3.1. Sponsor Requirements to Support Prescriber Certification

The Sponsors proposed additions to this section of the REMS document, including that Sponsors will ensure prescribers can complete the certification process by email or fax to an authorized distributor and/or certified pharmacy, and that Sponsors will ensure annually with each certified prescriber that their locations for receiving mifepristone are up to date. Sponsors will also ensure prescribers previously certified in the Mifepristone REMS Program complete the new *Prescriber Agreement Form*: (1) within 120 days after approval of this modification, for those previously certified prescribers submitting prescriptions to certified pharmacies, or (2) within one year after approval of this modification, if previously certified and ordering from an authorized distributor.

Reviewer Comment: *We agree with the Sponsors' proposal. The requirement to confirm that the locations associated with the certified prescriber are current is parallel to the pharmacy requirement that the authorized representative's contact information is up to date. In determining the pharmacy requirement, which is necessary to ensure program compliance and is consistent with other approved REMS that include pharmacy certification, the Agency also concluded that a parallel requirement for certified prescribers should be added.*

With respect to recertification, it is important that active certified prescribers are informed of and agree to new REMS requirements to ensure the continued safe use of mifepristone. There is minimal burden to recertification and the timelines allow sufficient time to accomplish recertification.

3.3.3.2. Sponsor Requirements to Support Pharmacy Certification

The Sponsors proposed the addition of Sponsor requirements to support pharmacy certification and compliance, including ensuring that pharmacies are certified in accordance with the requirements in the Mifepristone REMS Program, de-certifying pharmacies that do not maintain compliance with the certification requirements, and ensuring that pharmacy certification can be completed by email and fax to an authorized distributor. Annually, the authorized representative's name and contact information will be verified to ensure it corresponds to that of the current designated authorized representative for the certified pharmacy, and if different, a new authorized representative must certify for the pharmacy. All reference to the requirement in the 2021 Mifepristone REMS Program that mifepristone to be dispensed to patients only in clinics, medical offices and hospitals by or under the supervision of a certified prescriber, and not from retail pharmacies, was removed.

Reviewer Comment: *We agree with the Sponsors' proposal. Changes are in line with the REMS Modification Notification letters sent December 16, 2021. Refer to section 3.3.2.3 Reviewer Comments on Pharmacy Certification for rationale for removing the statement that mifepristone is not distributed to or dispensed from retail pharmacies. Ensuring that the authorized representative's contact information is up to date is necessary to ensure that there is always a point person who is responsible for implementing the Mifepristone REMS Program in their pharmacy and can address any changes that are needed if pharmacy audits identify a need for improvement.*

3.3.3.3. Sponsor Implementation Requirements

The Sponsors proposed that they will ensure that adequate records are maintained to demonstrate that REMS requirements have been met (including but not limited to records of mifepristone distribution, certification of prescribers and pharmacies, and audits of pharmacies and distributors), and that the records must be readily available for FDA inspections. The distributor audit requirement was updated to audit new distributors within 90 calendar days of becoming authorized and annually thereafter (a one-time audit requirement was previously required). The Sponsors also proposed a pharmacy audit requirement whereby certified pharmacies that order mifepristone are audited within 180 calendar days after the pharmacy places its first order of mifepristone, and annually thereafter for pharmacies that ordered in the previous 12 months.

Reviewer's Comment: *We agree with the Sponsors' proposal.*

The number of pharmacies that will certify in the REMS is uncertain; therefore, to obtain a reliable sample size for the audits, the Sponsors will need to audit all certified pharmacies within 180 calendar days after the pharmacy places its first order and annually thereafter for pharmacies that have ordered mifepristone in the previous 12 months. Audits performed at 180 days should allow time for establishment and implementation of audit protocols and for the Sponsors to perform the audits. With the addition of more stakeholders (i.e., certified pharmacies), it is also necessary to audit distributors annually to ensure the REMS requirements are followed. The requirement to conduct audits annually may be revisited if assessment data shows that the REMS is meeting its goal.

3.4. REMS Assessment Timetable

The Sponsors proposed that assessments must be submitted one year from the approval of the modified REMS and annually thereafter, instead of every three years as per the previous requirement.

Reviewer's Comment: *We agree with the Sponsors' proposal. With the addition of new pharmacy stakeholders and removal of the in-person dispensing requirement, more frequent assessment after this REMS modification is needed to ensure REMS processes are being followed and that the REMS is meeting its goal. The requirement can be revisited at a later date if assessment data shows that the modified REMS is meeting its goal. The NDA applicant is required to submit assessment reports as outlined in the timetable for submission of assessments. These reports address requirements for the Mifepristone REMS Program. The Sponsors have indicated that some data will be submitted as separate reports when Sponsor-specific information is needed to address the assessment metrics.*

4. Supporting Document

The Sponsors' REMS Supporting Document was substantially updated to include information regarding the proposed modification under review. Background and rationale from the 12/16/21 REMS Modification Notification letters was included. An updated description of the REMS goal and the ETASU was also included to align with the changes in the REMS Document and provide further clarification. Further explanation of prescriber requirements and rationale for various pharmacy requirements was also included.

Regarding implementation of the modified REMS, the Sponsors additionally proposed that pharmacies that received and shipped mifepristone during the Agency's exercise of enforcement discretion during the COVID-19 PHE, that wish to continue to dispense mifepristone, will be required to comply with the pharmacy certification requirements within 120 days of approval of the modified REMS.

The communication strategy to alert current and future prescriber and pharmacy stakeholders was outlined. Distributors, certified prescribers that purchased mifepristone in the last twelve months, and various professional organizations will receive information about REMS changes within 120 days of modification approval. The Sponsors proposed to list pharmacies that agree to be publicly disclosed on their respective product websites but disclosure of this nature is not a requirement of the REMS. The Sponsors indicated that they anticipate certified pharmacies that do not agree to public disclosure will communicate with the certified prescribers they wish to work with.

The REMS Assessment Plan is discussed in the following section.

Reviewer's Comment: *We agree with the Sponsors' proposal. The Supporting Document addresses all REMS requirements and provides sufficient clarification of implementation and maintenance of the REMS. The implementation requirements for pharmacies currently dispensing mifepristone under FDA's exercise of enforcement discretion during the COVID-19 PHE provide for continued use of these pharmacies without breaks in service. The communication strategy is also adequate given the efforts to reach both established certified prescribers and potentially new prescribers through professional organizations.*

The Sponsors' plan to communicate which pharmacies are certified to certified prescribers is adequate. For the reasons listed in section 3.3.2.3, confidentiality is a concern for REMS stakeholders. Disclosure of pharmacy certification status should be a choice made by individual certified pharmacies. The Sponsors have indicated that there will be some certified pharmacies that have agreed to publicly disclose their status, making this information available to certified prescribers who wish to use a pharmacy to dispense mifepristone.

5. REMS Assessment Plan

The REMS Assessment Plan is summarized in the REMS Supporting Document and will be included in the REMS Modification Approval letter.

The REMS Assessment Plan was revised to align with the modified REMS goal and objectives.

The goal of the Mifepristone REMS Program is to mitigate the risk of serious complications associated with mifepristone by:

- a. Requiring healthcare providers who prescribe mifepristone to be certified in the Mifepristone REMS Program.
 - This objective will be assessed using REMS Certification Statistics and REMS Compliance metrics.
- b. Ensuring that mifepristone is only dispensed by or under the supervision of certified prescribers, or by certified pharmacies on prescriptions issued by certified prescribers.
 - This objective will be assessed using REMS Certification Statistics and REMS Compliance metrics.
- c. Informing patients about the risk of serious complications associated with mifepristone.
 - This objective will be indirectly assessed using REMS Certification Statistics to avoid compromising patient and prescriber confidentiality. As part of the certification process, healthcare providers agree to:
 - Ensure that the *Patient Agreement Form* is reviewed with the patient and the risks of the mifepristone treatment regimen are fully explained
 - Ensure that the *Patient Agreement Form* is signed by the healthcare provider and the patient
 - Ensure that the patient is provided with a copy of the *Patient Agreement Form* and the Medication Guide
 - Ensure that the signed *Patient Agreement Form* is placed in the patient's medical record

The following revisions were made from the Mifepristone REMS Assessment Plan in the April 11, 2019, Supplement Approval letter:

The Assessment Plan Categories of 1) Program Implementation and Operations and 2) Overall Assessment of REMS Effectiveness were added.

REMS Certification Statistics metrics were added to capture certification numbers for program stakeholders to assess the first objective of requiring healthcare providers who prescribe mifepristone to be certified and the second objective of ensuring that mifepristone is only dispensed by or under the supervision of certified prescribers, or by certified pharmacies on prescriptions issued by certified prescribers. The total number of certified prescribers who certified with the wholesaler/distributor and the total number of certified prescribers who submitted a *Prescriber Agreement Form* to certified pharmacies were added to capture the additional method of prescriber certification. The number of newly certified prescribers and the number of active certified prescribers (i.e., those who ordered mifepristone or submitted a prescription during the reporting period) were added. Metrics were also added to capture the total number of certified, newly certified, and active certified pharmacies as well as the total number of authorized, newly authorized, and active authorized wholesaler/distributors.

Drug Utilization Data metrics were added to obtain information on shipment and dispensing of mifepristone. Metrics were added to capture the total number of tablets shipped by the wholesaler/distributor and the number of prescriptions dispensed.

REMS Compliance Data metrics were added to assess the first objective of requiring healthcare providers who prescribe mifepristone to be certified and the second objective of ensuring that mifepristone is only dispensed by or under the supervision of certified prescribers, or by certified pharmacies on prescriptions issued by certified prescribers. These metrics capture program deviations and evaluate overall if the REMS is operating as intended. Metrics include certified pharmacies and wholesaler/distributor audit results and a summary of instances of non-compliance and actions taken to address non-compliance. Prescriber compliance metrics were added to assess if prescribers are decertified along with reasons why. Pharmacy compliance metrics were added to assess if prescriptions were dispensed that were written by non-certified prescribers or if mifepristone tablets were dispensed by non-certified pharmacies as well as the number of pharmacies that were decertified along with reasons why. Wholesaler/distributor metrics were added to assess if shipments were sent to non-certified prescribers and non-certified pharmacies and corrective actions taken. The audit plan and non-compliance plans will be submitted for FDA review within 60 days after the REMS modification approval.

The Sponsors were asked to develop an assessment of prescription delivery timelines to determine what percentage of prescriptions were delivered on time (within four calendar days) and what percentage were delivered late (more than four calendar days) along with the length of the delay and reasons for the delay (e.g., mifepristone is out of stock shipment issues, other). The protocol for this assessment will be submitted for FDA review within 60 days after the REMS modification approval.

The revised REMS Assessment Plan is in the Appendix.

Reviewer's Comment: *We agree with the Sponsors' proposed REMS Assessment Plan.*

6. Discussion

The Sponsors submitted changes to the REMS to remove the requirement that mifepristone be dispensed only in certain healthcare settings (i.e., the "in-person dispensing requirement") and to add that certified pharmacies can dispense the drug in order to minimize the burden on the healthcare delivery system of complying with the REMS and to ensure that the benefits of the drug outweigh the risks. The REMS goal was updated to this effect. Changes were required for prescriber requirements and Sponsors to support the change in ETASU, and new pharmacy requirements were introduced.

The qualifications to become a certified prescriber have not changed as a result of the modification to the Mifepristone REMS Program; however, clarification has been provided for certain prescriber requirements and new prescriber requirements have been added to support pharmacy dispensing. Although certain responsibilities may be conducted by staff under the supervision of a certified prescriber, the certified prescriber remains responsible for ensuring compliance with the requirements of the Mifepristone REMS Program. In order to clarify this, revisions were made throughout the prescriber requirements and REMS materials to reflect that the certified prescriber is responsible for ensuring that the prescriber requirements are met. Additionally, the review team finds it acceptable that certified prescribers who wish to use a certified pharmacy to dispense mifepristone submit their *Prescriber Agreement Form* to the dispensing certified pharmacy (b) (4).

. The burden to prescriber and

pharmacy stakeholders of having certified prescribers submit the form directly to the certified pharmacy that will be dispensing the mifepristone is not unreasonable and has been minimized to the extent possible; it does not impact the safe use of the product. Prescriber requirements necessitated by the addition of some pharmacy requirements were added as well and include prescriber responsibilities in deciding whether or not mifepristone should be dispensed if the patient will receive the drug from the certified pharmacy more than four days after the pharmacy receives the prescription, and prescriber adverse event reporting requirements if a prescriber becomes aware of a patient death and the mifepristone was dispensed from a certified pharmacy. The addition of the latter requirements will ensure consistent adverse event data is relayed to the relevant Mifepristone Sponsor.

Changes were made to the *Patient Agreement Form*. Changes to the form were added to improve clarity of the safety messages. After further consideration, the patient's agreement to take the Medication Guide with them if they visit an emergency room or HCP who did not give them mifepristone so the emergency room or HCP will understand that the patient is having a medical abortion has been removed from the *Patient Agreement Form*. The Medication Guide is not typically carried by patients and this information can be obtained at the point of care. Changes align with updates to labeling submitted with this modification.^{13, 14}

The Agency and Sponsors agreed during this modification to focus on certification of pharmacies that can receive *Prescriber Agreement Forms* via email or fax to complete the prescriber certification process. The proposed pharmacy certification requirements also support timely dispensing of mifepristone. If the mifepristone is shipped to the patient, the REMS requires that it must be delivered within four calendar days from the receipt of the prescription by the pharmacy; if the patient will receive the mifepristone more than four calendar days from pharmacy receipt of prescription, the REMS requires the pharmacist to confirm with the certified prescriber that it is still appropriate to dispense the drug to the patient. This allows prescribers to make treatment decisions based on individual patient situations. A requirement to maintain confidentiality was also added to avoid unduly burdening patient access since patients and prescribers may not utilize pharmacy dispensing if they believe their personal information is at risk. Ultimately, the addition of pharmacy distribution with the proposed requirements will offer another option for dispensing mifepristone, alleviating burden associated with the REMS.

(b) (4)



The Agency reviewed the REMS in 2021, and per the review team's conclusions, a REMS modification was necessary to remove the in-person dispensing requirement and add a requirement that pharmacies that dispense the drug be specially certified; the review team concluded that these changes could occur without compromising patient safety. There have been no new safety concerns identified relevant to the REMS ETASUs that the applicants proposed modifying in their June 22, 2022 submissions since the REMS Modification Notification letters dated 12/16/2021. It is still the position of the review team that the proposed modification is acceptable.

Because the modification proposed include changes to the ETASU of the Mifepristone REMS Program, the assessment plan and timetable of assessments were changed. The assessment plan will capture information on pharmacy dispensing and provide valuable insight as to whether the program is operating as intended Annual assessments are consistent with other approved REMS modifications for major modifications necessitating extensive assessment plan changes.

As part of the REMS Assessment Plan, the REMS goal and objectives are assessed using Program Implementation and Operations Metrics, including REMS Certification Statistics and REMS Compliance Data. The metrics will provide information on the number of certified prescribers, certified pharmacies, and authorized wholesalers/distributors as well as if mifepristone is dispensed by non-certified prescribers or pharmacies. The Sponsors will use the indirect measure of healthcare provider certification to address the objective of informing patients of the risk of serious complications of mifepristone, due to concerns with prescriber and patient confidentiality. Although we typically assess whether patients are informed of the risks identified in a REMS through patient surveys and/or focus groups, we agree that the Sponsors' continued use of the indirect measure of healthcare provider certification adequately addresses the Mifepristone REMS Program objective of informing patients. In addition, because of these prescriber and patient confidentiality concerns, we believe it is unlikely that the Agency would be able to use the typical methods of assessment of patient knowledge and understanding of the risks and safe use of mifepristone.

7. Conclusions and Recommendations

The review team finds the proposed REMS modification for the Mifepristone REMS Program, as submitted on June 22, 2022, and amended on October 19, 2022 (Danco) and October 20, 2022 (GBP), November 30, 2022 (both), December 9 (both), and December 16 (both) acceptable. The REMS materials were amended to be consistent with the revised REMS document. The review team recommends approval of the Mifepristone REMS Program, received on June 22, 2022, and last amended on December 16, 2022, and appended to this review.

8. References

1. (b) (6) Clinical Review of SE-2 Efficacy Supplement for mifepristone, NDA 020687. March 29, 2016. DARRTS Reference ID: 3909590.
2. (b) (6) Summary Review for Regulatory Action for mifepristone, NDA 020687. March 29, 2016. DARRTS Reference ID: 3909594.
3. (b) (6) REMS Review for mifepristone, NDA 020687. March 29, 2016. DARRTS Reference ID: 3909588.
4. (b) (6) REMS Review for mifepristone, NDA 020687. March 29, 2016. DARRTS Reference ID: 3909587.
5. Approval Letter for SE-2 Efficacy Supplement for mifepristone, NDA 020687. March 29, 2016. DARRTS Reference ID: 3909592.
6. (b) (6) REMS Review for mifepristone, NDA 020687. February 22, 2018. DARRTS Reference ID: 4224674.
7. Approval Letter for SE-20 REMS Supplement for mifepristone, NDA 020687. March 29, 2016. DARRTS Reference ID: 4418041.
8. *Am. Coll. of Obstetricians & Gynecologists v. FDA*, 472 F. Supp. 3d 183, 233 (D. Md. July 13, 2020), order clarified, 2020 WL 8167535 (D. Md. Aug. 19, 2020) (preliminarily enjoining FDA from enforcing the in-person dispensing requirement and any other in-person requirements of the

Mifepristone SSS REMS); *FDA v. Am. Coll. of Obstetricians & Gynecologists*, 141 S. Ct. 578 (Jan. 12, 2021) (staying the preliminary injunction imposed by the District Court).

9. (b) (6) REMS Modification Rationale Review for mifepristone, NDA 020687. December 16, 2021. DARRTS Reference ID: 4905882.

10. General Advice Letter for the single, shared system Risk Evaluation and Mitigation Strategy (REMS) for mifepristone, NDA 020687, April 15, 2022. DARRTS ID 4969358.

11. Format and Content of a REMS Document Guidance for Industry <https://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM184128.pdf>. Accessed on December 18, 2022.

12. Grossman D, Raifman S, Morris N, et.al. Mail-order pharmacy dispensing of mifepristone for medication abortion after in-person clinical assessment. *Contraception* 2022; 107:36-41. <https://doi.org/10.1016/j.contraception.2021.09.008>. This article was included in the literature review for the December 16, 2021 REMS Modification Rationale Review, while the article was still in press.

9. Appendices

REMS Document

Prescriber Agreement Form for Danco Laboratories, LLC

Prescriber Agreement Form for GenBioPro, Inc.

Patient Agreement Form

Pharmacy Agreement Form for Danco Laboratories, LLC

Pharmacy Agreement Form for GenBioPro, Inc.

Mifepristone REMS Assessment Plan

Initial Shared System REMS approval: 04/2019

Most Recent Modification: 01/2023

Mifepristone Tablets, 200 mg

Progestin Antagonist

RISK EVALUATION AND MITIGATION STRATEGY (REMS) SINGLE SHARED SYSTEM FOR MIFEPRISTONE 200 MG

I. GOAL

The goal of the REMS for mifepristone is to mitigate the risk of serious complications associated with mifepristone by:

- a) Requiring healthcare providers who prescribe mifepristone to be certified in the Mifepristone REMS Program.
- b) Ensuring that mifepristone is only dispensed by or under the supervision of certified prescribers, or by certified pharmacies on prescriptions issued by certified prescribers.
- c) Informing patients about the risk of serious complications associated with mifepristone.

II. REMS ELEMENTS

A. Elements to Assure Safe Use

1. Healthcare providers who prescribe mifepristone must be specially certified.
 - a. To become specially certified to prescribe mifepristone, healthcare providers must:
 - i. Review the Prescribing Information for mifepristone.
 - ii. Complete a *Prescriber Agreement Form*. By signing¹ a *Prescriber Agreement Form*, prescribers agree that:
 - 1) They have the following qualifications:
 - a) Ability to assess the duration of pregnancy accurately
 - b) Ability to diagnose ectopic pregnancies
 - c) Ability to provide surgical intervention in cases of incomplete abortion or severe bleeding, or to have made plans to provide such care through others, and ability to assure patient access to medical facilities equipped to provide blood transfusions and resuscitation, if necessary
 - 2) They will follow the guidelines for use of mifepristone (see b.i-vii below).
 - b. As a condition of certification, prescribers must follow the guidelines for use of mifepristone described below:
 - i. Ensure that the *Patient Agreement Form* is reviewed with the patient and the risks of the mifepristone treatment regimen are fully explained. Ensure any questions the patient may have prior to receiving mifepristone are answered.
 - ii. Ensure that the healthcare provider and patient sign the *Patient Agreement Form*.

¹ In this REMS, the terms “sign” and “signature” include electronic signatures.

- iii. Ensure that the patient is provided with a copy of the *Patient Agreement Form* and Medication Guide.
- iv. Ensure that the signed *Patient Agreement Form* is placed in the patient's medical record.
- v. Ensure that any deaths are reported to the Mifepristone Sponsor that provided the mifepristone, identifying the patient by a non-identifiable reference and including the NDC and lot number from the package of mifepristone that was dispensed to the patient.
- vi. If mifepristone will be dispensed by a certified pharmacy:
 - 1) Provide the certified pharmacy a signed *Prescriber Agreement Form*.
 - 2) Assess appropriateness of dispensing mifepristone when contacted by a certified pharmacy about patients who will receive mifepristone more than 4 calendar days after the prescription was received by the certified pharmacy.
 - 3) Obtain the NDC and lot number of the package of mifepristone the patient received in the event the prescriber becomes aware of the death of the patient.
- vii. The certified prescriber who dispenses mifepristone or who supervises the dispensing of mifepristone must:
 - 1) Provide an authorized distributor with a signed *Prescriber Agreement Form*.
 - 2) Ensure that the NDC and lot number from each package of mifepristone dispensed are recorded in the patient's record.
 - 3) Ensure that healthcare providers under their supervision follow guidelines i.-v.
- c. Mifepristone Sponsors must:
 - i. Ensure that healthcare providers who prescribe their mifepristone are specially certified in accordance with the requirements described above and de-certify healthcare providers who do not maintain compliance with certification requirements.
 - ii. Ensure prescribers previously certified in the Mifepristone REMS Program complete the new *Prescriber Agreement Form*:
 - 1) Within 120 days after approval of this modification, for those previously certified prescribers submitting prescriptions to certified pharmacies.
 - 2) Within one year after approval of this modification, if previously certified and ordering from an authorized distributor.
 - iii. Ensure that healthcare providers can complete the certification process by email or fax to an authorized distributor and/or certified pharmacy.
 - iv. Provide the Prescribing Information and their *Prescriber Agreement Form* to healthcare providers who inquire about how to become certified.
 - v. Ensure annually with each certified prescriber that their locations for receiving mifepristone are up to date.

The following materials are part of the Mifepristone REMS Program:

- *Prescriber Agreement Form for Danco Laboratories, LLC*
- *Prescriber Agreement Form for GenBioPro, Inc.*
- *Patient Agreement Form*

2. Pharmacies that dispense mifepristone must be specially certified
 - a. To become specially certified to dispense mifepristone, pharmacies must:
 - i. Be able to receive *Prescriber Agreement Forms* by email and fax.
 - ii. Be able to ship mifepristone using a shipping service that provides tracking information.
 - iii. Designate an authorized representative to carry out the certification process on behalf of the pharmacy.
 - iv. Ensure the authorized representative oversees implementation and compliance with the Mifepristone REMS Program by doing the following:
 - 1) Review the Prescribing Information for mifepristone.
 - 2) Complete a *Pharmacy Agreement Form*. By signing a *Pharmacy Agreement Form*, the authorized representative agrees that the pharmacy will put processes and procedures in place to ensure the following requirements are completed:
 - a) Verify that the prescriber is certified by confirming their completed *Prescriber Agreement Form* was received with the prescription or is on file with the pharmacy.
 - b) Dispense mifepristone such that it is delivered to the patient within 4 calendar days of the date the pharmacy receives the prescription, except as provided in c) below.
 - c) Confirm with the prescriber the appropriateness of dispensing mifepristone for patients who will receive the drug more than 4 calendar days after the date the pharmacy receives the prescription and document the prescriber's decision.
 - d) Record in the patient's record the NDC and lot number from each package of mifepristone dispensed.
 - e) Track and verify receipt of each shipment of mifepristone.
 - f) Dispense mifepristone in its package as supplied by the Mifepristone Sponsor.
 - g) Report any patient deaths to the prescriber, including the NDC and lot number from the package of mifepristone dispensed to the patient, and remind the prescriber of their obligation to report the deaths to the Mifepristone Sponsor that provided the mifepristone. Notify the Mifepristone Sponsor that provided the dispensed mifepristone that the pharmacy submitted a report of death to the prescriber, including the name and contact information for the prescriber and the NDC and lot number of the dispensed product.
 - h) Not distribute, transfer, loan or sell mifepristone except to certified prescribers or other locations of the pharmacy.
 - i) Maintain records of *Prescriber Agreement Forms*.
 - j) Maintain records of dispensing and shipping.
 - k) Maintain records of all processes and procedures including compliance with those processes and procedures.
 - l) Maintain the identity of the patient and prescriber as confidential, including limiting access to patient and prescriber identity only to those personnel necessary to dispense mifepristone in accordance with the Mifepristone REMS Program requirements, or as necessary for payment and/or insurance purposes.
 - m) Train all relevant staff on the Mifepristone REMS Program requirements.

- n) Comply with audits carried out by the Mifepristone Sponsors or a third party acting on behalf of the Mifepristone Sponsors to ensure that all processes and procedures are in place and are being followed.
- b. Mifepristone Sponsors must:
 - i. Ensure that pharmacies are specially certified in accordance with the requirements described above and de-certify pharmacies that do not maintain compliance with certification requirements.
 - ii. Ensure that pharmacies can complete the certification process by email and fax to an authorized distributor.
 - i. Verify annually that the name and contact information for the pharmacy's authorized representative corresponds to that of the current designated authorized representative for the certified pharmacy, and if different, require the pharmacy to recertify with the new authorized representative.

The following materials are part of the Mifepristone REMS Program:

- *Pharmacy Agreement Form for Danco Laboratories, LLC*
 - *Pharmacy Agreement Form for GenBioPro, Inc.*
- 3. Mifepristone must be dispensed to patients with evidence or other documentation of safe use conditions as ensured by the certified prescriber in signing the *Prescriber Agreement Form*.
 - a. The patient must sign a *Patient Agreement Form* indicating that the patient has:
 - i. Received, read and been provided a copy of the *Patient Agreement Form*.
 - ii. Received counseling from the healthcare provider regarding the risk of serious complications associated with mifepristone.

B. Implementation System

- 1. Mifepristone Sponsors must ensure that their mifepristone is only distributed to certified prescribers and certified pharmacies by:
 - a. Ensuring that distributors who distribute their mifepristone comply with the program requirements for distributors.
 - i. The distributors must put processes and procedures in place to:
 - 1) Complete the certification process upon receipt of a *Prescriber Agreement Form* or *Pharmacy Agreement Form*.
 - 2) Notify healthcare providers and pharmacies when they have been certified by the Mifepristone REMS Program.
 - 3) Ship mifepristone only to certified pharmacies or locations identified by certified prescribers.
 - 4) Not ship mifepristone to pharmacies or prescribers who become de-certified from the Mifepristone REMS Program.
 - 5) Provide the Prescribing Information and their Prescriber Agreement Form to healthcare providers who (1) attempt to order mifepristone and are not yet certified, or (2) inquire about how to become certified.
 - ii. Put processes and procedures in place to maintain a distribution system that is secure,

confidential and follows all processes and procedures, including those for storage, handling, shipping, tracking package serial numbers, NDC and lot numbers, proof of delivery and controlled returns of mifepristone.

- iii. Train all relevant staff on the Mifepristone REMS Program requirements.
 - iv. Comply with audits by Mifepristone Sponsors or a third party acting on behalf of Mifepristone Sponsors to ensure that all processes and procedures are in place and are being followed for the Mifepristone REMS Program. In addition, distributors must maintain appropriate documentation and make it available for audits.
- b. Ensuring that distributors maintain secure and confidential distribution records of all shipments of mifepristone.
2. Mifepristone Sponsors must monitor their distribution data to ensure compliance with the Mifepristone REMS Program.
 3. Mifepristone Sponsors must ensure that adequate records are maintained to demonstrate that the Mifepristone REMS Program requirements have been met, including, but not limited to records of mifepristone distribution; certification of prescribers and pharmacies; and audits of pharmacies and distributors. These records must be readily available for FDA inspections.
 4. Mifepristone Sponsors must audit their new distributors within 90 calendar days and annually thereafter after the distributor is authorized to ensure that all processes and procedures are in place and functioning to support the requirements of the Mifepristone REMS Program. Mifepristone Sponsors will take steps to address their distributor compliance if noncompliance is identified.
 5. Mifepristone Sponsors must audit their certified pharmacies within 180 calendar days after the pharmacy places its first order of mifepristone, and annually thereafter audit certified pharmacies that have ordered mifepristone in the previous 12 months, to ensure that all processes and procedures are in place and functioning to support the requirements of the Mifepristone REMS Program. Mifepristone Sponsors will take steps to address their pharmacy compliance if noncompliance is identified.
 6. Mifepristone Sponsors must take reasonable steps to improve implementation of and compliance with the requirements of the Mifepristone REMS Program based on monitoring and assessment of the Mifepristone REMS Program.
 7. Mifepristone Sponsors must report to FDA any death associated with mifepristone whether or not considered drug-related, as soon as possible but no later than 15 calendar days from the initial receipt of the information by the Mifepristone Sponsor. This requirement does not affect the sponsors' other reporting and follow-up requirements under FDA regulations.

C. Timetable for Submission of Assessments

The NDA Sponsor must submit REMS assessments to FDA one year from the date of the approval of the modified REMS (1/3/2023) and annually thereafter. To facilitate inclusion of as much information as possible while allowing reasonable time to prepare the submission, the reporting interval covered by each assessment should conclude no earlier than 90 calendar days before the submission date for that assessment. The NDA Sponsor must submit each assessment so that it will be received by the FDA on or before the due date.

MIFEPREX® (Mifepristone) Tablets, 200 mg

PRESCRIBER AGREEMENT FORM

Mifeprex* (Mifepristone) Tablets, 200 mg, is indicated, in a regimen with misoprostol, for the medical termination of intrauterine pregnancy through 70 days gestation. Please see Prescribing Information and Medication Guide for complete safety information.

To **become a certified prescriber**, you must:

- **If you submit Mifeprex prescriptions for dispensing from certified pharmacies:**
 - o Submit this form to each certified pharmacy to which you intend to submit Mifeprex prescriptions. The form must be received by the certified pharmacy before any prescriptions are dispensed by that pharmacy.
- **If you order Mifeprex for dispensing by you or healthcare providers under your supervision:**
 - o Submit this form to the distributor. This form must be received by the distributor before the first order will be shipped to the healthcare setting.
 - o Healthcare settings, such as medical offices, clinics, and hospitals, where Mifeprex will be dispensed by or under the supervision of a certified prescriber in the Mifepristone REMS Program do not require pharmacy certification.

Prescriber Agreement: By signing this form, you agree that you meet the qualifications below and will follow the guidelines for use. You are responsible for overseeing implementation and compliance with the Mifepristone REMS Program. You also understand that if the guidelines below are not followed, the distributor may stop shipping mifepristone to the locations that you identify and certified pharmacies may stop accepting your mifepristone prescriptions.

Mifepristone must be provided by or under the supervision of a certified prescriber who meets the following qualifications:

- Ability to assess the duration of pregnancy accurately.
- Ability to diagnose ectopic pregnancies.
- Ability to provide surgical intervention in cases of incomplete abortion or severe bleeding, or have made plans to provide such care through others, and be able to assure patient access to medical facilities equipped to provide blood transfusions and resuscitation, if necessary.
- Has read and understood the Prescribing Information for mifepristone. The Prescribing Information is available by calling 1-877-4 EARLY OPTION (1-877-432-7596 toll-free), or by visiting www.earlyoptionpill.com.

In addition to meeting these qualifications, you also agree to follow these guidelines for use:

- Ensure that the *Patient Agreement Form* is reviewed with the patient and the risks of the mifepristone treatment regimen are fully explained. Ensure any questions the patient may have prior to receiving mifepristone are answered.
- Ensure the healthcare provider and patient sign the *Patient Agreement Form*.
- Ensure that the patient is provided with a copy of the *Patient Agreement Form* and Medication Guide.
- Ensure that the signed *Patient Agreement Form* is placed in the patient's medical record.
- Ensure that any deaths of patients who received Mifeprex are reported to Danco Laboratories, LLC, identifying the patient by a non-identifiable reference and including the NDC and lot number from the package of Mifeprex that was dispensed to the patient.



*MIFEPREX is a registered trademark of Danco Laboratories, LLC
P.O. Box 4816-New York, NY 10185

1-877-4-EARLY-OPTION (1-877-432-7596) www.earlyoptionpill.com

Ensure that healthcare providers under your supervision follow the guidelines listed above.

- If Mifeprex will be dispensed through a certified pharmacy:
 - o Assess appropriateness of dispensing Mifeprex when contacted by a certified pharmacy about patients who will receive Mifeprex more than 4 calendar days after the prescription was received by the certified pharmacy.
 - o Obtain the NDC and lot number of the package of Mifeprex the patient received in the event the prescriber becomes aware of the death of a patient.
- If Mifeprex will be dispensed by you or by healthcare providers under your supervision:
 - o Ensure the NDC and lot number from each package of Mifeprex are recorded in the patient's record.

I understand that a certified pharmacy may dispense mifepristone made by a different manufacturer than that stated on this Prescriber Agreement Form.

Print Name: _____ Title: _____

Signature: _____ Date: _____

Medical License # _____ State _____

NPI # _____

Practice Setting Address: _____

Return completed form to Mifeprex@dancodistributor.com or fax to 1-866-227-3343.

Approved 01/2023 [Doc control ID]



*MIFEPREX is a registered trademark of Danco Laboratories, LLC
P.O. Box 4816-New York, NY 10185
1-877-4-EARLY-OPTION (1-877-432-7596) www.earlyoptionpill.com

PRESCRIBER AGREEMENT FORM

Mifepristone Tablets, 200 mg

Mifepristone Tablets, 200 mg, is indicated, in a regimen with misoprostol, for the medical termination of intrauterine pregnancy through 70 days gestation. Please see Prescribing Information and Medication Guide for complete safety information.

To **become a certified prescriber**, you must:

- **If you submit mifepristone prescriptions for dispensing from certified pharmacies:**
 - o Submit this form to each certified pharmacy to which you intend to submit mifepristone prescriptions. The form must be received by the certified pharmacy before any prescriptions are dispensed by that pharmacy.
- **If you order mifepristone for dispensing by you or healthcare providers under your supervision:**
 - o Submit this form to the distributor. This form must be received by the distributor before the first order will be shipped to the healthcare setting.
 - o Healthcare settings, such as medical offices, clinics, and hospitals, where mifepristone will be dispensed by or under the supervision of a certified prescriber in the Mifepristone REMS Program do not require pharmacy certification.

Prescriber Agreement: By signing this form, you agree that you meet the qualifications below and will follow the guidelines for use. You are responsible for overseeing implementation and compliance with the Mifepristone REMS Program. You also understand that if the guidelines below are not followed, the distributor may stop shipping mifepristone to the locations that you identify and certified pharmacies may stop accepting your mifepristone prescriptions.

Mifepristone must be provided by or under the supervision of a certified prescriber who meets the following qualifications:

- Ability to assess the duration of pregnancy accurately.
- Ability to diagnose ectopic pregnancies.
- Ability to provide surgical intervention in cases of incomplete abortion or severe bleeding, or have made plans to provide such care through others, and be able to assure patient access to medical facilities equipped to provide blood transfusions and resuscitation, if necessary.
- Has read and understood the Prescribing Information for mifepristone. The Prescribing Information is available by calling 1-855-MIFE-INFO (1-855—643-3463 toll-free), or by visiting www.MifeInfo.com.

In addition to meeting these qualifications, you also agree to follow these guidelines for use:

- Ensure that the *Patient Agreement Form* is reviewed with the patient and the risks of the mifepristone treatment regimen are fully explained. Ensure any questions the patient may have prior to receiving mifepristone are answered.
- Ensure the healthcare provider and patient sign the *Patient Agreement Form*.
- Ensure that the patient is provided with a copy of the *Patient Agreement Form* and Medication Guide.
- Ensure that the signed *Patient Agreement Form* is placed in the patient's medical record.
- Ensure that any deaths of patients who received mifepristone are reported to GenBioPro, Inc. that provided the mifepristone, identifying the patient by a non-identifiable reference and including the NDC and lot number from the package of mifepristone that was dispensed to the patient.

Ensure that healthcare providers under your supervision follow the guidelines listed above.



GenBioPro Inc. - PO Box 32011 - Las Vegas, NV 89103
1-855-MIFE-INFO (1-855-643-3463) - www.MifeInfo.com

- If mifepristone will be dispensed through a certified pharmacy:
 - o Assess appropriateness of dispensing mifepristone when contacted by a certified pharmacy about patients who will receive mifepristone more than 4 calendar days after the prescription was received by the certified pharmacy.
 - o Obtain the NDC and lot number of the package of mifepristone the patient received in the event the prescriber becomes aware of the death of a patient.
- If mifepristone will be dispensed by you or by healthcare providers under your supervision:
 - o Ensure the NDC and lot number from each package of mifepristone are recorded in the patient's record.

I understand that a certified pharmacy may dispense mifepristone made by a different manufacturer than that stated on this Prescriber Agreement Form.

Print Name: _____ Title: _____

Signature: _____ Date: _____

Medical License # _____ State _____

NPI # _____

Practice Setting Address: _____

Return completed form to RxAgreements@GenBioPro.com or fax to 1-877-239-8036

Approved 01/2023 [Doc control ID]

PATIENT AGREEMENT FORM

Mifepristone Tablets, 200 mg

Healthcare Providers: *Counsel the patient on the risks of mifepristone. Both you and the patient must provide a written or electronic signature on this form.*

Patient Agreement:

1. I have decided to take mifepristone and misoprostol to end my pregnancy and will follow my healthcare provider's advice about when to take each drug and what to do in an emergency.
2. I understand:
 - a. I will take mifepristone on Day 1.
 - b. I will take the misoprostol tablets 24 to 48 hours after I take mifepristone.
3. My healthcare provider has talked with me about the risks, including:
 - heavy bleeding
 - infection
4. I will contact the clinic/office/provider right away if in the days after treatment I have:
 - a fever of 100.4°F or higher that lasts for more than four hours
 - heavy bleeding (soaking through two thick full-size sanitary pads per hour for two hours in a row)
 - severe stomach area (abdominal) pain or discomfort, or I am “feeling sick,” including weakness, nausea, vomiting, or diarrhea, more than 24 hours after taking misoprostol
— these symptoms may be a sign of a serious infection or another problem (including an ectopic pregnancy, a pregnancy outside the womb).

My healthcare provider has told me that these symptoms listed above could require emergency care. If I cannot reach the clinic/office/provider right away, my healthcare provider has told me who to call and what to do.
5. I should follow up with my healthcare provider about 7 to 14 days after I take mifepristone to be sure that my pregnancy has ended and that I am well.
6. I know that, in some cases, the treatment will not work. This happens in about 2 to 7 out of 100 women who use this treatment. If my pregnancy continues after treatment with mifepristone and misoprostol, I will talk with my provider about a surgical procedure to end my pregnancy.
7. If I need a surgical procedure because the medicines did not end my pregnancy or to stop heavy bleeding, my healthcare provider has told me whether they will do the procedure or refer me to another healthcare provider who will.
8. I have the MEDICATION GUIDE for mifepristone.
9. My healthcare provider has answered all my questions.

Patient Signature: _____ **Patient Name (print):** _____ **Date:** _____

Provider Signature: _____ **Provider Name (print):** _____ **Date:** _____

Patient Agreement Forms may be provided, completed, signed, and transmitted in paper or electronically.

01/2023

MIFEPREX®(Mifepristone) Tablets, 200mg

PHARMACY AGREEMENT FORM

Pharmacies must designate an authorized representative to carry out the certification process and oversee implementation and compliance with the Mifepristone REMS Program on behalf of the pharmacy.

Healthcare settings, such as medical offices, clinics, and hospitals, where mifepristone will be dispensed by or under the supervision of a certified prescriber in the Mifepristone REMS Program do not require pharmacy certification.

By signing this form, as the Authorized Representative I certify that:

- Each location of my pharmacy that will dispense Mifeprex is able to receive *Prescriber Agreement Forms* by email and fax.
- Each location of my pharmacy that will dispense Mifeprex is able to ship Mifeprex using a shipping service that provides tracking information.
- I have read and understood the Prescribing Information for Mifeprex. The Prescribing Information is available by calling 1-877-4 EARLY OPTION (1-877-432-7596 toll-free) or online at www.earlyoptionpill.com; and
- Each location of my pharmacy that will dispense Mifeprex will put processes and procedures in place to ensure the following requirements are completed. I also understand that if my pharmacy does not complete these requirements, the distributor may stop accepting Mifeprex orders.
 - o Verify that the prescriber is certified in the Mifepristone REMS Program by confirming their completed *Prescriber Agreement Form* was received with the prescription or is on file with your pharmacy.
 - o Dispense Mifeprex such that it is delivered to the patient within 4 calendar days of the date the pharmacy receives the prescription, except as provided in the following bullet.
 - o Confirm with the prescriber the appropriateness of dispensing Mifeprex for patients who will receive the drug more than 4 calendar days after the date the pharmacy receives the prescription and document the prescriber's decision.
 - o Record in the patient's record the NDC and lot number from each package of Mifeprex dispensed.
 - o Track and verify receipt of each shipment of Mifeprex.
 - o Dispense mifepristone in its package as supplied by Danco Laboratories, LLC.
 - o Report any patient deaths to the prescriber, including the NDC and lot number from the package of Mifeprex dispensed to the patient, and remind the prescriber of their obligation to report the deaths to Danco Laboratories, LLC. Notify Danco that your pharmacy submitted a report of death to the prescriber, including the name and contact information for the prescriber and the NDC and lot number of the dispensed product.
 - o Not distribute, transfer, loan or sell mifepristone except to certified prescribers or other locations of the pharmacy.
 - o Maintain records of *Prescriber Agreement Forms*, dispensing and shipping, and all processes and procedures including compliance with those processes and procedures.
 - o Maintain the identity of Mifeprex patients and prescribers as confidential and protected from disclosure except to the extent necessary for dispensing under this REMS or as necessary for payment and/or insurance.
 - o Train all relevant staff on the Mifepristone REMS Program requirements.
 - o Comply with audits carried out by the Mifepristone Sponsors or a third party acting on behalf of the Mifepristone Sponsors to ensure that all processes and procedures are in place and are being followed.

Any new authorized representative must complete and submit the *Pharmacy Agreement Form*.

Authorized Representative Name: _____ Title: _____



*MIFEPREX is a registered trademark of Danco Laboratories, LLC

P.O. Box 4816-New York, NY 10185

1-877-4-EARLY-OPTION (1-877-432-7596) www.earlyoptionpill.com

Signature: _____ Date: _____

Email: _____ Phone: _____ Preferred __ email __ phone

Pharmacy Name: _____

Pharmacy Address: _____

Return completed form to Mifeprex@dancodistributor.com or fax to 1-866-227-3343.



*MIFEPREX is a registered trademark of Danco Laboratories, LLC
P.O. Box 4816-New York, NY 10185
1-877-4-EARLY-OPTION (1-877-432-7596) www.earlyoptionpill.com

PHARMACY AGREEMENT FORM

Mifepristone Tablets, 200 mg

Pharmacies must designate an authorized representative to carry out the certification process and oversee implementation and compliance with the Mifepristone REMS Program on behalf of the pharmacy.

Healthcare settings, such as medical offices, clinics, and hospitals, where mifepristone will be dispensed by or under the supervision of a certified prescriber in the Mifepristone REMS Program do not require pharmacy certification.

By signing this form, as the Authorized Representative I certify that:

- Each location of my pharmacy that will dispense mifepristone is able to receive *Prescriber Agreement Forms* by email and fax.
- Each location of my pharmacy that will dispense mifepristone is able to ship mifepristone using a shipping service that provides tracking information.
- I have read and understood the Prescribing Information for mifepristone. The Prescribing Information is available by calling 1-855-MIFE-INFO (1-855-643-3463 toll-free) or online at www.MifeInfo.com; and
- Each location of my pharmacy that will dispense mifepristone will put processes and procedures in place to ensure the following requirements are completed. I also understand that if my pharmacy does not complete these requirements, the distributor may stop accepting mifepristone orders.
 - o Verify that the prescriber is certified in the Mifepristone REMS Program by confirming their completed *Prescriber Agreement Form* was received with the prescription or is on file with your pharmacy.
 - o Dispense mifepristone such that it is delivered to the patient within 4 calendar days of the date the pharmacy receives the prescription, except as provided in the following bullet.
 - o Confirm with the prescriber the appropriateness of dispensing mifepristone for patients who will receive the drug more than 4 calendar days after the date the pharmacy receives the prescription and document the prescriber's decision.
 - o Record in the patient's record the NDC and lot number from each package of mifepristone dispensed.
 - o Track and verify receipt of each shipment of mifepristone.
 - o Dispense mifepristone in its package as supplied by GenBioPro, Inc.
 - o Report any patient deaths to the prescriber, including the NDC and lot number from the package of mifepristone dispensed to the patient, and remind the prescriber of their obligation to report the deaths to GenBioPro, Inc. Notify GenBioPro that your pharmacy submitted a report of death to the prescriber, including the name and contact information for the prescriber and the NDC and lot number of the dispensed product.
 - o Not distribute, transfer, loan or sell mifepristone except to certified prescribers or other locations of the pharmacy.
 - o Maintain records of *Prescriber Agreement Forms*, dispensing and shipping, all processes and procedures including compliance with those processes and procedures.
 - o Maintain the identity of mifepristone patients and prescribers as confidential and protected from disclosure except to the extent necessary for dispensing under this REMS or as necessary for payment and/or insurance purposes.
 - o Train all relevant staff on the Mifepristone REMS Program requirements.
 - o Comply with audits carried out by the Mifepristone Sponsors or a third party acting on behalf of the Mifepristone Sponsors to ensure that all processes and procedures are in place and are being followed.

Any new authorized representative must complete and submit the *Pharmacy Agreement Form*.

Authorized Representative Name: _____ Title: _____

Signature: _____ Date: _____

Email: _____ Phone: _____ Preferred ___ email ___ phone

Pharmacy Name: _____

Pharmacy Address: _____

Return completed form to RxAgreements@GenBioPro.com or fax to 1-877-239-8036.



GenBioPro Inc. - PO Box 32011 - Las Vegas, NV 89103
1-855-MIFE-INFO (1-855-643-3463) - www.MifeInfo.com

The REMS Assessment Plan must include but is not limited to the following items.

Program Implementation and Operations

1. REMS Certification Statistics

a. Prescribers

- i. Number of certified prescribers who have certified with the Sponsor's distributor(s) and number who have submitted *Prescriber Agreement Forms* to Certified Pharmacies
- ii. Number and percentage of newly certified prescribers
- iii. Number and percentage of active certified prescribers (i.e., who ordered mifepristone or submitted a prescription during the reporting period)

b. Pharmacies

- i. Number of certified pharmacies
- ii. Number and percentage of newly certified pharmacies
- iii. Number and percentage of active certified pharmacies (i.e., that dispensed mifepristone during the reporting period)

c. Wholesalers/Distributors

- i. Number of authorized wholesalers/distributors
- ii. Number and percentage of newly authorized wholesalers/distributors
- iii. Number and percentage of active authorized wholesalers/distributors (i.e. that shipped mifepristone during the reporting period)

2. Utilization Data

- a. Total number of tablets shipped by wholesalers/distributors, stratified by Certified Prescriber or Certified Pharmacy location
- b. Number of prescriptions dispensed from pharmacies

3. REMS Compliance Data

- a. Audits: Summary of audit activities for each stakeholder (i.e., certified pharmacies and wholesalers/distributors) including but not limited to:
 - i. A copy of the final audit plan for each stakeholder type (provide for the current reporting period)
 - ii. The number of audits expected, and the number of audits performed
 - iii. The number and type of deficiencies noted
 - iv. For those with deficiencies noted, report the corrective and preventive actions (CAPAs) required, if any, to address the deficiencies, including the status (e.g., completed, not completed, in progress) (provide for the current reporting period)
 - v. For any stakeholders that did not complete the CAPA within the timeframe specified in the audit plan, describe actions taken (provide for the current reporting period)

- vi. A summary report of all resulting changes to processes and procedures necessary to ensure compliance with the REMS requirements (provide for the current reporting period)
- b. A summary report of non-compliance, associated corrective action plans (CAPAs), and the status of CAPAs including but not limited to:
 - i. A copy of the final non-compliance plans for Pharmacies and Distributors (provide for the current reporting period)
 - ii. For each instance of noncompliance below (iii-v), report the following information (provide for the current reporting period):
 - 1. A unique, anonymized ID for the stakeholder(s) associated with the non-compliance event to enable tracking over time
 - 2. The source of the non-compliance data (e.g., self-reported, audit, other)
 - 3. A root cause analysis of the non-compliance
 - 4. Actions to prevent future occurrences and outcomes of such actions
 - iii. Prescriber compliance
 - 1. Number and percentage of certified prescribers who became decertified as a result of non-compliance
 - Provide a summary of reasons for decertification (provide for the current reporting period)
 - 2. Summary and analysis of any program deviations and corrective actions taken (provide for the current reporting period)
 - iv. Pharmacy compliance
 - 1. Number and percentage of prescriptions dispensed that were written by prescriber(s) who did not submit a Prescriber Agreement to the dispensing Certified Pharmacy
 - 2. Number and percentage of mifepristone tablets dispensed by non-certified pharmacies
 - 3. Number and percentage of pharmacies that became decertified as a result of non-compliance
 - Provide a summary of reasons for decertification (provide for the current reporting period)
 - 4. An assessment of prescription delivery timelines, including percentage delivered more than four days after receipt of the prescription, duration and causes for delay. A proposal for this assessment will be submitted within 60 days of the approval of the REMS Modification.
 - 5. Summary and analysis of any program deviations and corrective actions taken (provide for the current reporting period)
 - v. Wholesaler/distributor compliance
 - 1. Number of healthcare providers who successfully ordered mifepristone who were not certified
 - 2. Number of non-certified pharmacies that successfully ordered mifepristone
 - 3. Number of shipments sent to non-certified prescriber receiving locations
 - 4. Number of shipments sent to non-certified pharmacy receiving locations

5. Summary and analysis of any program deviations and corrective actions taken (provide for the current reporting period)

The requirements for assessments of an approved REMS under section 505-1(g)(3) include with respect to each goal included in the strategy, an assessment of the extent to which the approved strategy, including each element of the strategy, is meeting the goal or whether one or more such goals or such elements should be modified.

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

(b) (6)

01/03/2023 05:18:27 PM

(b) (6)

01/03/2023 05:19:15 PM

(b) (6)

01/03/2023 05:24:28 PM

(b) (6)

01/03/2023 05:27:04 PM

(b) (6)

01/03/2023 05:27:58 PM

(b) (6)

01/03/2023 05:29:45 PM

(b) (6)

01/03/2023 05:33:47 PM

Center for Drug Evaluation and Research (CDER)

(b) (6) (b) (6)
(b) (6) (b) (6)
(b) (6) (b) (6)
(b) (6) (b) (6)
(b) (6) (b) (6)
(b) (6) (b) (6)
(b) (6) (b) (6)

Application Type

NDA and ANDA

Application Number

020687 and 91178

Reviewer Names

(b) (6) (b) (6)
(b) (6)
(b) (6) (b) (6) (b) (6)

(b) (6)

(b) (6) (b) (6)

(b) (6) (b) (6)

(b) (6) (b) (6)

(b) (6)

(b) (6)

(b) (6) (b) (6)

(b) (6)

(b) (6)

(b) (6)

(b) (6)

(b) (6)

Review Completion Date

December 16, 2021

Subject	REMS Modification Rationale Review
Established Name	Mifepristone REMS
Name of Applicants	Danco Laboratories, LLC and GenBioPro, Inc.
Therapeutic Class	Progestin antagonist
Formulation	Oral tablets

Table of Contents

EXECUTIVE SUMMARY	4
1. Introduction	5
2. Background	5
2.1. PRODUCT AND REMS INFORMATION	5
2.2. REGULATORY HISTORY AND EVENTS RELEVANT TO THIS REMS MODIFICATION RATIONALE REVIEW.....	7
3. Rationale for Proposed REMS Modification	9
3.1. CURRENT REQUIREMENTS FOR THE APPROVED REMS	10
3.2. EVALUATION OF THE EVIDENCE	10
3.2.1. Evaluation of the requirement for healthcare providers who prescribe the drug to be specially certified (ETASU A)	12
3.2.2. Evaluation of the requirement for the drug to be dispensed with evidence or other documentation of safe-use conditions (ETASU D)	14
3.2.3. Evaluation of the requirement for drug to be dispensed only in certain healthcare settings (ETASU C)	19
4. Discussion.....	36
5. Conclusions and Recommendations	41
6. References	42
7. Appendix.....	45

EXECUTIVE SUMMARY

This review provides the (b) (6) (b) (6) and (b) (6) (b) (6) rationale and conclusions regarding modifications to the single, shared system Risk Evaluation and Mitigation Strategy (REMS) for mifepristone 200 mg (Mifepristone REMS Program) for new drug application (NDA) 20687 and abbreviated new drug application (ANDA) 91178.

ANDA 91178 was approved with the approval of the Mifepristone REMS Program on April 11, 2019 to mitigate the risk of serious complications associated with mifepristone 200 mg. The most recent REMS modification was approved on May 14, 2021. The REMS consists of elements to assure safe use (ETASU) under ETASU A, C and D, an implementation system, and a timetable for submission of assessments. To determine whether a modification to the REMS was warranted, FDA undertook a comprehensive review of the published literature; safety information collected during the COVID-19 public health emergency (PHE); the one-year REMS assessment report of the Mifepristone REMS Program; adverse event data; and information provided by advocacy groups, individuals and the Applicants. Our review also included an examination of literature references provided by plaintiffs in the *Chelius v. Becerra* litigation discussed below.

The modifications to the REMS will consist of:

- Removing the requirement under ETASU C that mifepristone be dispensed only in certain healthcare settings, specifically clinics, medical offices, and hospitals (referred to here as the “in-person dispensing requirement” for brevity)
- Adding a requirement under ETASU B that pharmacies that dispense the drug be specially certified

A REMS Modification Notification letter will be sent to both Applicants in the Single Shared System.

1. Introduction

In connection with the *Chelius v. Becerra* litigation, FDA agreed to undertake a full review of the Mifepristone REMS Program, in accordance with the REMS assessment provisions of the Federal Food, Drug, and Cosmetic Act (FD&C Act).^a This review provides the analysis of the (b) (6) (b) (6) and the (b) (6) (b) (6) regarding whether any changes are warranted to the single, shared system Risk Evaluation and Mitigation Strategy (REMS) for mifepristone (hereafter referred to as the Mifepristone REMS Program) for new drug application (NDA) 20687 and abbreviated new drug application (ANDA) 91178. The Mifeprex REMS was initially approved in 2011; the single, shared system REMS for mifepristone 200 mg, known as the Mifepristone REMS Program, was approved in 2019.

The last time the existing REMS elements to assure safe use (under ETASU A, C and D) were reviewed was in the context of our review of supplement S-020 to NDA 20687; these ETASU were updated following review and approval of supplement S-020 on March 29, 2016. The key changes approved in 2016 are summarized below.

Changes to labeling included:

- Changing the dosing of Mifeprex to 200 mg orally x 1
- Extension of maximum gestational age through 70 days
- Inclusion of misoprostol in the indication statement
- Replacing the term “physician” with “licensed healthcare provider”
- Removal of the phrase “Under Federal Law”

The Mifeprex REMS and REMS materials were updated to reflect the changes above, and additional changes were made including:

- Removing the Medication Guide as part of the REMS but retaining it as part of labeling.

2. Background

2.1. PRODUCT AND REMS INFORMATION

^a Section 505-1(g)(2) of the FD&C Act (21 U.S.C. § 355-1(g)(2)).

Mifepristone is a progestin antagonist indicated, in a regimen with misoprostol, for the medical termination of intrauterine pregnancy (IUP) through 70 days gestation. Mifepristone is available as 200 mg tablets for oral use.

Mifeprex (mifepristone) was approved on September 28, 2000 with a restricted distribution program under 21 CFR 314.520 (subpart H)^b to ensure that the benefits of the drug outweighed the risk of serious complications associated with mifepristone when used for medical abortion. Mifeprex was deemed to have a REMS under section 505-1 of the Federal Food, Drug, and Cosmetic Act with the passage of the Food and Drug Administration Amendments Act (FDAAA) of 2007, and the Mifeprex REMS was approved on June 8, 2011. On March 29, 2016, as noted above, a supplemental application and REMS modification was approved for Mifeprex. On April 11, 2019, ANDA 091178 was approved, and the Mifepristone REMS Program was approved. The Mifepristone REMS Program is a single, shared system REMS that includes NDA 020687 and ANDA 91178.

The goal of the REMS for mifepristone is to mitigate the risk of serious complications associated with mifepristone by:

- a. Requiring healthcare providers who prescribe mifepristone to be certified in the Mifepristone REMS Program (under ETASU A).
- b. Ensuring that mifepristone is only dispensed in certain healthcare settings, by or under the supervision of a certified prescriber (under ETASU C).
- c. Informing patients about the risk of serious complications associated with mifepristone (under ETASU D).

Under ETASU A, to become specially certified to prescribe mifepristone, a healthcare provider must review the prescribing information, complete and sign the *Prescriber Agreement Form*, and follow the guidelines for use of mifepristone. Under ETASU C, mifepristone must be dispensed to patients only in certain healthcare settings, specifically clinics, medical offices, and hospitals, by or under the supervision of a certified prescriber. Under ETASU D, mifepristone must be dispensed to patients with evidence or other documentation of safe use conditions (i.e., the patient must sign a *Patient Agreement Form*). The Mifepristone REMS Program also includes an implementation system, and a timetable for assessments (one year from the date of the initial approval of the REMS on April 11, 2019, and every three years thereafter).

^b NDA approval letter Mifeprex (NDA 020687) dated September 28, 2000.

2.2. REGULATORY HISTORY AND EVENTS RELEVANT TO THIS REMS MODIFICATION RATIONALE REVIEW

The following is a summary of significant regulatory history since approval of the REMS modification on March 29, 2016:

- 03/29/2016: FDA approved an efficacy supplement (S-020) that, among other things, provided a new dosing regimen (200 mg mifepristone, followed in 24 to 48 hours by 800 mcg buccal misoprostol), increased the gestational age (GA) to which mifepristone may be used (through 70 days gestation), and modified the REMS.
- 03/29/2019: A Citizen Petition was received requesting that FDA revise the product labeling to reflect pre-2016 provisions (including limiting GA to 49 days and requiring patients to make 3 office visits) and that FDA maintain the REMS.
- 04/11/2019: ANDA 91178 was approved along with the Single Shared System REMS for Mifepristone 200 mg (Mifepristone REMS Program) for NDA 20687 and ANDA 91178.
- 01/31/2020: the COVID-19 public health emergency (PHE) was declared by the Secretary of Health and Human Services (HHS) as having existed since January 27, 2020.^c
- 7/13/2020: The United States (US) District Court of Maryland granted a preliminary injunction in the *ACOG v. FDA* litigation to temporarily bar enforcement of the Mifepristone REMS Program in-person dispensing requirement during the COVID-19 PHE.
- 1/12/2021: US Supreme Court granted a stay of that injunction.
- 04/12/2021: FDA issued a General Advice Letter to both the NDA and ANDA Applicants, stating that provided that all other requirements of the Mifepristone REMS Program are met, and given that in-person dispensing of mifepristone for medical termination of early pregnancy may present additional COVID-related risks to patients and healthcare

^c See Secretary of Health and Human Services, Determination that a Public Health Emergency Exists (originally issued January 31, 2020, and subsequently renewed), available at <https://www.phe.gov/emergency/news/healthactions/phe/Pages/default.aspx>

personnel because it may involve a clinical visit solely for this purpose, FDA intends to exercise enforcement discretion during the COVID-19 PHE with respect to the in-person dispensing requirement in the Mifepristone REMS Program, including any in-person requirements that may be related to the *Patient Agreement Form*. FDA further stated that to the extent all of the other requirements of the Mifepristone REMS Program are met, FDA intends to exercise enforcement discretion during the COVID-19 PHE with respect to the dispensing of mifepristone through the mail, either by or under the supervision of a certified prescriber, or through a mail-order pharmacy when such dispensing is done under the supervision of a certified prescriber.

- 05/07/2021: FDA stated that it would be reviewing the elements of the Mifepristone REMS Program in accordance with the REMS assessment provisions of section 505-1 of the FD&C Act.
- 05/14/2021: A modification was approved for the Mifepristone REMS Program. This modification was to revise the *Patient Agreement Form* to include gender-neutral language.
- 06/30/2021: An Information Request (IR) was sent to the Applicants for additional information on shipments and any program deviations, adverse events, or noncompliance with the REMS that occurred during the period from April 1, 2021 through September 30, 2021.
- 7/15/2021: An IR was sent to the Applicants to provide the total number of shipments during the period from April 1, 2021 to September 30, 2021 and details on whether any of those shipments were involved in any program deviation or non-compliance.
- 8/5/2021: An IR was sent to the Applicants for additional clinical and other information (e.g., adverse events and units of mifepristone shipped) for the period of March 29, 2016 through June 30, 2021, to be provided by August 31, 2021. This IR also requested information covering the period of July 1, 2021 through September 30, 2021 and an

aggregate summary (for the period of March 29, 2016 through September 30, 2021), to be provided by October 12, 2021.^d

- 8/26/2021: The ANDA Applicant submitted a response to the IR issued on 8/5/2021.
- 08/27/2021: The NDA Applicant submitted a response to the IR issued on 8/5/2021.
- 10/08/2021: The NDA Applicant submitted a response to the June 30 and July 15, 2021 IRs as well as an aggregate summary for the period March 29, 2016 through September 30, 2021 in response to the August 5, 2021 IR. The NDA Applicant also included a follow-up to their initial response provided on August 27, 2021 to the August 5, 2021 IR.
- 10/12/2021: The ANDA Applicant submitted a response to the June 30 and July 15, 2021 IRs as well as an aggregate summary for the period March 29, 2016 through September 30, 2021 in response to the August 5, 2021 IR.
- 10/16/2021: The ANDA Applicant revised their Oct 12, 2012 response to provide a correction to the number of mifepristone tablets.
- [REDACTED] (b) (4)
[REDACTED]
- 11/02/2021: A [REDACTED] (b) (6) [REDACTED] (b) (6) meeting was convened to obtain CDER concurrence on the removal of the in-person dispensing requirement and the addition of a certification requirement for pharmacies. The [REDACTED] (b) (6) [REDACTED] (b) (6) and senior CDER leadership concurred with removing the in-person dispensing and adding pharmacy certification.

3. Rationale for Proposed REMS Modification

^d Multiple Information Requests were issued to obtain additional information on drug shipments, any program deviations or noncompliance, and use of alternative methods for drug distribution during the COVID-19 PHE. These IRs are referenced as appropriate in this document and the one-year REMS Assessment Review of the Mifepristone REMS Program, December 16, 2021.

3.1. CURRENT REQUIREMENTS FOR THE APPROVED REMS

The Mifepristone REMS Program includes elements to assure safe use (ETASU), an implementation system, and a timetable for submission of assessments. Elements to assure safe use in the current REMS include a prescriber certification requirement (ETASU A), a requirement that mifepristone be dispensed only in certain healthcare settings by or under the supervision of a certified prescriber (ETASU C), and a requirement that mifepristone be dispensed only with documentation of safe use conditions (ETASU D). Documentation of safe use conditions under ETASU D consists of a *Patient Agreement Form* between the prescriber and the patient indicating that the patient has received counseling from the prescriber regarding the risk of serious complications associated with mifepristone 200 mg for medical termination of early pregnancy.

3.2. EVALUATION OF THE EVIDENCE

We reviewed multiple different sources of information, including published literature, safety information submitted to the Agency during the COVID-19 PHE, FDA Adverse Event Reporting System (FAERS) reports, the first REMS assessment report for the Mifepristone REMS Program, and information provided by advocacy groups, individuals, and the Applicants. Our review also included an examination of literature references provided by plaintiffs in the *Chelius v. Becerra* litigation. Below is an overview of how information relevant to the current Mifepristone REMS Program was retrieved, analyzed, and applied to each of the individual ETASUs to determine if further changes should be considered.

Methods for the literature search

(b) (6) conducted a literature search in PubMed and Embase to retrieve publications relevant to this review. The time period used for this literature search was between March 29, 2016 (when the Mifeprex labeling and REMS were last substantially revised) through July 26, 2021. The search terms used were “medical abortion” and “mifepristone” and “pregnancy termination and mifepristone.”

The search retrieved 306 publications from PubMed and 613 from Embase, respectively; the search yielded 646 unique publications after eliminating duplications between the two databases. The result of our literature search was also supplemented by an examination of literature references provided by advocacy groups, individuals, plaintiffs in the *Chelius* litigation, and the Applicants, as well as letters from healthcare providers and researchers.

References included in these letters were considered for inclusion in this review using identical selection criteria to the (b) (6) literature search (outlined below).

For this review of the REMS, (b) (6) focused on publications containing safety data related to outcomes of medical abortion (objective safety data) obtained from our literature search and from the references provided to us relevant to the REMS ETASUs. We excluded systematic reviews and meta-analyses because these publications did not include original safety data related to the outcomes of medical abortion. The following are examples of materials that were excluded from our literature search:

- Information from survey studies or qualitative studies that evaluated perspectives on and/or satisfaction with medical abortion procedures from patients, pharmacists, clinic staff, or providers, even if the study assessed REMS ETASUs. These surveys or qualitative studies did not include objective safety data related to outcomes of medical abortion.
- Opinions, commentaries, or policy/advocacy statements. These publications did not include objective safety data related to outcomes of medical abortion.
- Safety data related to mifepristone use for second trimester medical abortion. These publications reported data not applicable to the approved indication for medical abortion up to 70 days gestation.
- Safety data related to mifepristone use for spontaneous first trimester abortion (i.e., miscarriages). These publications reported data not applicable to the approved indication for medical abortion up to 70 days gestation.
- Safety data that pertained only to surgical abortion or did not separate out medical abortion from surgical abortion.
- Other safety information unrelated to the REMS elements (e.g., articles limited to case reports or those discussing unrelated gynecologic or medical issues)
- Publications for which it was not possible to conduct a full review of the methods or results, i.e., the references were limited to an abstract of the study methods and results.
- Publications that provided only general statistics on abortion care in the United States.

- Information pertinent to molecular or other basic science aspects of mifepristone.
- Data on the logistics of accessing abortion care in general, such as time to appointment or the distance traveled to obtain care.
- Publications that provided data not related specifically to abortion care or the REMS (e.g., references focused on federal poverty guidelines, poverty data, or the financial impact of the COVID-19 pandemic).

One exception to the above literature search criteria was the inclusion in Section 3.2.2 of this review, which discusses the *Patient Agreement Form*, of publications that discussed changes in provider volume. The data discussed in relation to provider volume was obtained from surveys. This data was included because changes in provider volume could only be obtained from well-conducted survey studies.

Regarding medical/scientific references submitted with letters from the plaintiffs in the *Chelius* litigation, we applied the same criteria as for the literature search, as described above.

Letters from the plaintiffs in the *Chelius* litigation included several references that preceded our 2016 review of the REMS. Two of those pre-2016 studies were not captured in our 2016 literature search. These two studies were assessed as part of our current review; their results are consistent with the existing safety profile of the approved medical abortion regimen, and therefore, support our current conclusions regarding the REMS. See Appendix A.

3.2.1. Evaluation of the requirement for healthcare providers who prescribe the drug to be specially certified (ETASU A)

In order to become specially certified, prescribers must: 1) review the prescribing information for mifepristone and 2) complete the *Prescriber Agreement Form*. In signing the *Prescriber Agreement Form*, prescribers agree they meet the qualifications listed below:

- Ability to assess the duration of pregnancy accurately
- Ability to diagnose ectopic pregnancies
- Ability to provide surgical intervention in cases of incomplete abortion or severe bleeding, or to have made plans to provide such care through others, and ability to

ensure patient access to medical facilities equipped to provide blood transfusions and resuscitation, if necessary.

- Has read and understood the Prescribing Information of mifepristone (which the provider can access by phone or online).

In addition to meeting these qualifications, as a condition of certification the healthcare provider also agrees to follow the guidelines for use below:

- Review the *Patient Agreement Form* with the patient and fully explain the risks of the mifepristone treatment regimen. Answer any questions the patient may have prior to receiving mifepristone.
- Sign and obtain the patient's signature on the *Patient Agreement Form*.
- Provide the patient with a copy of the *Patient Agreement Form* and the Medication Guide.
- Place the signed *Patient Agreement Form* in the patient's medical record.
- Record the serial number from each package of mifepristone in each patient's record.
- Report deaths to the Applicant, identifying the patient by a non-identifiable patient reference and the serial number from each package of mifepristone.

The literature review was the primary source of information that contributed to our reassessment of ETASU A.

We continue to be concerned that absent these provider qualifications, serious and potentially fatal complications associated with medical abortion, including missed ectopic pregnancy and heavy bleeding from incomplete abortion, would not be detected or appropriately managed. Our review of the literature did not identify any studies comparing providers who met these qualifications with providers who did not. In the absence of such studies, there is no evidence to contradict our previous finding that prescribers' ability to accurately date pregnancies, diagnose ectopic pregnancies, and provide surgical intervention or arrange for such care through others if needed, is necessary to mitigate the serious risks associated with the use of mifepristone in a regimen with misoprostol. Therefore, our review continues to support the conclusion that a healthcare provider who prescribes mifepristone should meet the above qualifications. We conclude it is reasonable to maintain the requirement for a one-time prescriber certification where prescribers attest to having the ability to diagnose an intrauterine

pregnancy, to diagnose an ectopic pregnancy,^e and to either manage serious complications themselves or arrange for other providers to provide the needed care in a timely manner.

In addition, in signing the *Prescriber Agreement Form* and placing it in the patient's medical record, the prescribers acknowledge the requirement to report patient deaths associated with mifepristone to the manufacturer. Such a requirement ensures that the manufacturer receives all reports of patient deaths and, in turn, fulfills its regulatory obligations to report those deaths to the FDA.

As discussed in Section 3.2.2 below, there is a potential for doubling of the number of prescribers of mifepristone if the in-person dispensing requirement in ETASU C is removed from the Mifepristone REMS Program. Given the potential addition of new prescribers, in addition to the considerations described above, we conclude that we should maintain the requirement for prescriber certification, to ensure that providers meet the necessary qualifications and adhere to the guidelines for use. Our literature review supports that these requirements are still necessary, and the potential increase in new prescribers under the REMS is a further reason to maintain prescriber certification. Healthcare provider certification continues to be a necessary component of the REMS to ensure the benefits of mifepristone for medical abortion outweigh the risks. The burden of prescriber certification has been minimized to the extent possible by requiring prescribers to certify only one time for each applicant.

3.2.2. Evaluation of the requirement for the drug to be dispensed with evidence or other documentation of safe-use conditions (ETASU D)

In order to receive mifepristone for medical termination of pregnancy through 70 days gestation, the patient must sign a *Patient Agreement Form* indicating that the patient has received, read, and been provided a copy of the *Patient Agreement Form* and received counseling from the prescriber regarding the risk of serious complications associated with mifepristone for this indication. The *Patient Agreement Form* ensures that patients are informed of the risks of serious complications associated with mifepristone for this indication.

^e American College of Obstetricians and Gynecologists (ACOG) Practice Bulletin Number 191, February 2018. Tubal Ectopic Pregnancy. <https://www.acog.org/clinical/clinical-guidance/practice-bulletin/articles/2018/03/tubal-ectopic-pregnancy>. Mifepristone is not effective for terminating ectopic pregnancy. Some of the expected symptoms experienced with a medical abortion (abdominal pain, uterine bleeding) may be similar to those of a ruptured ectopic pregnancy. A missed ectopic pregnancy that ruptures is a medical emergency that requires immediate surgical intervention.

In a number of approved REMS, *Patient Agreement Forms* or *Patient Enrollment Forms* ensure that patients are counseled about the risks of the product and/or informed of appropriate safe use conditions.^f

As a condition of certification under the Mifepristone REMS Program, healthcare providers must follow the guidelines for use of mifepristone, including reviewing the *Patient Agreement Form* with the patient, fully explaining the risks of the treatment regimen, and answering any questions the patient may have before receiving the medication. With this form, the patient acknowledges that they have received and read the form, and that they have received the counseling regarding when to take mifepristone, the risk of serious complications associated with mifepristone and what to do if they experience adverse events (e.g., fever, heavy bleeding). Both the healthcare provider and patient must sign the document and the patient must receive a copy of the signed form. In addition to the counseling described in the *Patient Agreement Form*, patients also receive a copy of the Medication Guide for mifepristone. Ultimately, the *Patient Agreement Form* serves as an important counseling component, and documentation that the safe use conditions of the Mifepristone REMS Program have been satisfied, as the prescriber is required to place the signed *Patient Agreement Form* in the patient's medical record.

Prior to the March 29, 2016 approval of the S-020 efficacy supplement for Mifeprex, FDA undertook a review of all elements of the REMS. At that time, the (b) (6), (b) (6), along with the (b) (6), (b) (6), recommended removal of the *Patient Agreement Form* (ETASU D). This recommendation received concurrence from the (b) (6) on February 23, 2016. The rationale for this recommendation in the 2016 (b) (6) review^g is summarized here as follows:

- The safety profile of Mifeprex is well-characterized over 15 years of experience, with known risks occurring rarely; the safety profile has not changed over the period of surveillance.
- Established clinical practice includes patient counseling and documentation of informed consent and evidence shows that practitioners are providing appropriate patient

^f REMS@FDA, <https://www.accessdata.fda.gov/scripts/cder/remis/index.cfm>, Accessed November 15, 2021.

^g (b) (6) Clinical Review, NDA 020687/S20, dated March 29, 2016. https://darrrts.fda.gov/darrrts/faces/ViewDocument?documentId=090140af803dc7bd&_afRedirect=386175573203745

counseling and education; the *Patient Agreement Form* is duplicative of these established practices.

- Medical abortion with Mifeprex is provided by a small group of organizations and their associated providers. Their documents and guidelines are duplicated in the *Patient Agreement Form*.
- ETASUs A and C remain in place: The *Prescriber Agreement Form* and the requirement that Mifeprex be dispensed to patients only in certain healthcare settings, specifically, clinics, medical offices, and hospitals under the supervision of a certified prescriber, remain in place.

In light of a memorandum from the Director of the Center for Drug Evaluation and Research, an addendum to the (b) (6) March 29, 2016 review and a memorandum from the signatory authority in (b) (6) indicated that the *Patient Agreement Form* would be retained in the REMS.^{h,i}

The current review of literature from March 29, 2016 to July 26, 2021, is relevant to our assessment of the necessity of the *Patient Agreement Form* as part of the REMS. While our literature search yielded no publications which directly addressed this element of the REMS, we identified the following literature that focused on the informed consent process. These studies were reviewed for their potential relevance on this topic, though the articles do not directly assess the need for the *Patient Agreement Form* as a condition necessary to assure safe use of Mifepristone under ETASU D.

- Two studies^{1,2} (both authored by Dr. Grossman in 2021) used the *Patient Agreement Form* and additional clinic-specific written informed consent forms as part of the study methodology. One study evaluated medical abortion with pharmacist dispensing of mifepristone and another evaluated mail-order pharmacy dispensing. Safety and efficacy outcomes were not assessed regarding the element of consent in isolation or the *Patient Agreement Form*.
- Several studies included use of electronic or verbal consent. Two studies were conducted using signed electronic consent (Chong³, Kerestes⁴). Aiken⁵ reported that patients had the option of providing consent verbally and the discussion had to be recorded in the notes. Rocca⁶ described obtaining verbal informed consent from patients seeking medical abortion provided in pharmacies or government-certified

^h (b) (6) Review of proposed REMS modifications to Mifeprex. March 29, 2106.

ⁱ (b) (6) Summary of Regulatory Action for Mifeprex. March 29, 2016.

public health facilities by auxiliary nurse midwives (ANMs) in Nepal. Outcomes were not assessed regarding the single element of consent and its role in the efficacy of medical abortion.

- A retrospective chart review (Wiebe⁷) was conducted in Canada. This study included telemedicine abortions between January 31, 2017 and January 31, 2019 and a similar group of controls seen in the clinic during the same time frame, matched by date of initial appointment. As part of the telemedicine process, patients read a consent form (not specified whether they could view an electronic version) and gave verbal consent “witnessed by the counselor”. Again, outcomes were not assessed regarding the single element of consent and its role in the efficacy of medical abortion.

After review, we conclude that there are no outcome data from these studies that address the need for the *Patient Agreement Form* as a condition necessary to assure safe use of mifepristone. Nor do any of these studies provide evidence of whether the patient’s informed consent has been adequately documented under the process set out in the study protocol. Therefore, these studies do not provide evidence that would support removing ETASU D.

Although (b) (6) agrees that informed consent in medicine is an established practice, the National Abortion Federation’s 2020 Clinical Policy Guidelines for Abortion Care⁸ continue to include a detailed section on patient education, counseling, and informed consent. The guidelines state that these steps are essential parts of the abortion process; that they should be conducted by appropriate personnel, with accurate information, including about alternatives and potential risks and benefits; and that the patients must have an opportunity to have any questions answered to their satisfaction prior to any intervention. Under these guidelines, documentation must show that the patient affirms that they understand all the information provided and that the decision to undergo an abortion is voluntary. The guidelines specifically list the risks that must be addressed at a minimum, including those pertinent to medical abortion: hemorrhage, infection, continuing pregnancy, and death. Additionally, Practice Bulletins from ACOG⁹ and the Society of Family Planning also support detailed patient counseling.

In addition, trends in US clinical practice are developing which could negatively impact adequate patient counseling about the risks of medical abortion. One survey by Jones 2017¹⁰ of abortion providers in the United States and Canada prior to the COVID-19 pandemic did reveal strong adherence to evidence-based guidelines. However, this same survey noted continued increasing uptake of medical abortion by US providers. Grossman¹¹ conducted a US survey in

2019 which suggested that the number of obstetrician/gynecologists providing medical abortion care may be increasing and that uptake might increase if mifepristone were dispensed by pharmacies instead of being dispensed in-person. A subsequent survey of US obstetricians/gynecologists by Daniel in 2021¹² evaluated a subsample (n = 868) from a prior national survey of providers and found that 164 (19%) reported providing medical abortion in the previous year. Of those obstetrician/gynecologists not providing medical abortion, 171 (24%) said they would offer the method to their patients if the in-person dispensing requirement for mifepristone were removed. This indicates a potential doubling of providers (+ 104%, 95% confidence interval (CI): 97% –112%). There were geographical variations, with the largest potential increases being in the Midwest (+ 189%, 95% CI: 172% –207%) and the South (+ 118%, 95% CI: 103% –134%).

Based on the articles discussed above, removal of the in-person dispensing requirement from the Mifepristone REMS Program (as discussed below in section 3.2.3) could significantly increase the number of providers to a larger group of practitioners. The *Patient Agreement Form* is an important part of standardizing the medication information on the use of mifepristone that prescribers communicate to their patients, and also provides the information in a brief and understandable format for patients. The requirement to counsel the patient, to provide the patient with the *Patient Agreement Form*, and to have the healthcare provider and patient sign the *Patient Agreement Form*, ensures that each provider, including new providers, informs each patient of the appropriate use of mifepristone, risks associated with treatment, and what to do if the patient experiences symptoms that may require emergency care. The single-page *Patient Agreement Form* is in line with other elements of this REMS, in that it supports the requirement that certified prescribers be able to accurately assess a patient, counsel a patient appropriately and recognize and manage potential complications. The form is placed in the patient's medical record to document the patient's acknowledgment of receiving the information from the prescriber and a copy is provided to the patient. We determined, consistent with section 505-1(f)(2) of the FD&C Act, that this does not impose an unreasonable burden on providers or patients, and that the *Patient Agreement Form* remains necessary to assure the safe use of Mifepristone.

After considering potential burden on healthcare providers and patients and considering the available data discussed above, including the potential for increased prescribing of mifepristone if in-patient dispensing is removed from the REMS, we conclude that the *Patient Agreement Form* should remain a safe use condition in the REMS.

3.2.3. Evaluation of the requirement for drug to be dispensed only in certain healthcare settings (ETASU C)

Mifepristone applicants must ensure that mifepristone is available to be dispensed to patients only in clinics, medical offices, and hospitals by or under the supervision of a certified prescriber. This creates what we refer to in this document as an in-person dispensing requirement under the REMS; i.e., the patient must be present in person in the clinic, medical office or hospital when the drug is dispensed. The mifepristone REMS document states that mifepristone may not be distributed to or dispensed through retail pharmacies or settings other than these.

The following information contributed to our analysis of this requirement: Mifepristone REMS Program year-one assessment data, postmarketing safety information and literature review.

REMS Assessment Data

Reporting period for the Mifepristone REMS Program - April 11, 2019 through February 29, 2020

We evaluated information included in the one-year (1st)^j REMS assessment reports for the Mifepristone REMS Program, which included healthcare provider certification data, program utilization data, compliance data, audit results and patient exposure data.¹³ The assessment reports were submitted on April 10, 2020 by the NDA Applicant and April 15, 2020 by the ANDA Applicant and cover a reporting period from April 11, 2019 through February 29, 2020. During this reporting period, the NDA Applicant reported (b) (4) newly certified healthcare providers, and the ANDA Applicant reported (b) (4) newly certified healthcare providers in the Mifepristone REMS Program. The NDA Applicant reported a total of (b) (4) certified healthcare providers (includes new and previously certified) ordered mifepristone during the assessment reporting period, and the ANDA Applicant reported a total of (b) (4) certified healthcare providers ordered mifepristone during the assessment reporting period. The NDA Applicant estimated that a total of (b) (4) patients were exposed to mifepristone during the assessment reporting period. The ANDA Applicant reported an estimated total of (b) (4) patients were exposed to mifepristone during the reporting period.

During the reporting period, a small number of non-compliance events were reported. The authorized distributor for the NDA applicant reported to the NDA Applicant that they experienced deviations with scanning of the product serial numbers which were confirmed during the February 2020 audit. The authorized distributor conducted a root cause analysis and developed a corrective and preventive action (CAPA) on February 12, 2020. The CAPA was

^j This REMS assessment report was the first to be submitted following the approval of the single, shared system REMS for mifepristone.

validated and deployed with monitoring of the system through April 10, 2020. The corrective action will prevent similar events from occurring in the future.

January 27, 2020 through September 30, 2021

During the timeframe from January 27, 2020 through September 30, 2021, there were periods when the in-person dispensing requirement was not being enforced.

- On July 13, 2020, the United States District Court for the District of Maryland granted a preliminary injunction in the ACOG case to temporarily bar enforcement of the in-person dispensing requirement during the COVID-19 PHE.
- On January 12, 2021, the United States Supreme Court issued a stay of the injunction.
- On April 12, 2021, the FDA issued a General Advice Letter informing the applicants of the Agency's intent to exercise enforcement discretion during the COVID-19 public health emergency regarding the in-person dispensing requirement in the Mifepristone REMS Program.^{k,l}

To better understand whether there was any impact on safety or noncompliance during the periods when the in-person dispensing requirement was not being enforced, we requested additional information from the Applicants to provide for more comprehensive assessment of the REMS for the time period from January 27, 2020 (the effective date of the COVID-19 PHE) to September 30, 2021. We requested the Applicants provide a summary and analysis of any program deviation or noncompliance events from the REMS requirements and any adverse events that occurred during this time period that had not already been submitted to FDA. As part of an additional request for information for the REMS assessment report, the Applicants were also asked to submit the adverse events to FAERS and to notify FDA that the reports were submitted.

Between January 27, 2020 and September 30, 2021, the NDA Applicant distributed (b) (4) shipments representing (b) (4) tablets. The NDA Applicant reported that there were (b) (4) shipments representing a total of (b) (4) tablets sent to (b) (4) non-certified healthcare providers.^{m,n} (b) (4) of these healthcare providers subsequently became certified while (b) (4) did not. Of the (b) (4) healthcare providers who were not subsequently certified, (b) (4) returned a total of 12 of the 13

^k FDA General Advice Letter for NDA 20687, April 12, 2021.

^l FDA General Advice Letter for ANDA 091178, April 12, 2021.

^m NDA 020687 September 9, 2021 response to the FDA's September 2, 2021 Information Request.

ⁿ NDA 020687 October 8, 2021 response to the FDA's June 30, 2021 Information Request.

Mifeprex tablets to the distributor. (b) (4) non-certified healthcare provider dispensed one tablet to a patient; no adverse events were reported. The NDA Applicant attributed the non-compliance observed to the authorized distributor's transition to a new platform. The NDA Applicant implemented a corrective and preventative action to address this issue, which we found to be acceptable.

The ANDA Applicant distributed (b) (4) shipments representing (b) (4) tablets of mifepristone from January 27, 2020 to September 30, 2021 and reported no instances of shipments to non-certified healthcare providers during this timeframe.

The NDA and the ANDA applicants reported a total of eight cases reporting adverse events between January 27, 2020 and September 30, 2021. These eight cases were also identified in the FAERS database and are described in the section below.

The number of adverse events reported to FDA during the COVID-19 PHE with mifepristone use for medical termination of pregnancy is small, and the data provide no indication that any program deviation or noncompliance with the Mifepristone REMS Program contributed to these reported adverse events. Further analysis of the adverse events is included below in the section on Pharmacovigilance Data.

Pharmacovigilance Data

The (b) (6) (b) (6) conducted a search of the FAERS database and the published medical literature to identify U.S. postmarketing adverse events that reportedly occurred from January 27, 2020 through September 30, 2021 with mifepristone use for medical termination of pregnancy.^{o,p}

The data for this time period were then further divided into date ranges when the in-person dispensing requirement was being enforced per the REMS (January 27, 2020 - July 12, 2020 & January 13, 2021 - April 12, 2021) versus when the in-person dispensing requirement was not being enforced (July 13, 2020 - January 12, 2021 (in-person dispensing requirement was temporarily enjoined) & April 13, 2021 - September 30, 2021 (in-person dispensing requirement was not being enforced because of the COVID-19 PHE)).

^c (b) (6). Pharmacovigilance Memorandum: Mifepristone and All Adverse Events. NDA 020687 and ANDA 091178. (b) (6) # 2007-525. Finalized April 12, 2021.

^p (b) (6). Pharmacovigilance Memorandum: Mifepristone and All Adverse Events. NDA 020687 and ANDA 091178. (b) (6) # 2007-525. Finalized December 16, 2021.

A total of eight cases that met the search criteria were identified in FAERS and no additional case reports were identified in the medical literature. Two of the eight cases reported adverse events that occurred when the in-person dispensing requirement in the REMS was being enforced (i.e., January 27, 2020 - July 12, 2020 & January 13, 2021 - April 12, 2021). These two cases reported the occurrence of uterine/vaginal bleeding (case 1) and uterine/vaginal bleeding and sepsis (case 2). Of note, uterine/vaginal bleeding and sepsis are labeled adverse events. Five of the eight cases reported adverse events that occurred when the in-person dispensing requirement was not being enforced (i.e., July 13, 2020 - January 12, 2021 & April 13, 2021 - September 30, 2021). These five cases reported the occurrence of ongoing pregnancy (case 3), drug intoxication and death approximately 5 months after ingestion of mifepristone (case 4), death [cause of death is currently unknown] (case 5), sepsis and death (case 6), and pulmonary embolism (case 7). Although these adverse events occurred during the period when the in-person dispensing requirement was not being enforced, the narratives provided in the FAERS reports for cases 5, 6, and 7 explicitly stated that mifepristone was dispensed in-person. Of note, ongoing pregnancy, and sepsis, including the possibility of fatal septic shock, are labeled adverse events. The remaining case from July 2021 reported the occurrence of oral pain/soreness (case 8) but did not provide sufficient information to determine the exact date of the adverse event. Based upon the U.S. postmarketing data reviewed, no new safety concerns were identified by (b) (6)

In addition to the FAERS data provided above, (b) (6) routinely monitors adverse events reported to FAERS and published in the medical literature for mifepristone for medical termination of pregnancy. (b) (6) has not identified any new safety concerns with the use of mifepristone for medical termination of pregnancy.

To enable additional review of adverse events, the Applicants were requested^q to provide a summary and analysis of adverse events reported with incomplete medical abortion requiring surgical intervention to complete abortion, blood transfusion following heavy bleeding or hemorrhage, ectopic pregnancies, sepsis, infection without sepsis, hospitalization related to medical abortion, and emergency department (ED)/urgent care encounter related to medical abortion. The Applicant for Mifeprex provided a summary of postmarketing safety information from March 29, 2016, when S-020 was approved, through September 30, 2021, on August 27 and October 8, 2021. During the time period in question, (b) (4) tablets were shipped, and

^q On August 5, 2021, an IR was sent to the Applicants requesting a summary and analysis of adverse events from March 29, 2016 through June 30, 2021 and from July 1, 2021 through September 30, 2021.

48 adverse events were received. The 48 adverse events included 4 deaths (one of which occurred in 2010 but was reported in 2017), 25 incomplete abortions requiring surgical intervention, 17 blood transfusions following heavy vaginal bleeding, 2 ectopic pregnancies, 7 infections (1 sepsis and 6 infection without sepsis), 13 hospitalizations, and 43 ED or urgent care visits related to medical abortion. For the period between January 27, 2020 and September 30, 2021, a time frame that includes the entire period when the COVID-19 public health emergency (PHE) has been in effect, there were three adverse events reported corresponding to the above cases from FAERS identified by (b) (6) case 1 (uterine/vaginal bleeding), case 2 (uterine/vaginal bleeding and sepsis), and case 4 (drug intoxication and death).

The ANDA Applicant provided a summary of postmarketing safety information from April 11, 2019 (date of ANDA approval) through September 30, 2021. On August 26, 2021, the Applicant provided distribution and adverse event information from April 11, 2019 through June 30, 2021. During this time period, a total of (b) (4) tablets were shipped. There were 7 adverse events including 3 deaths (1 from sepsis, 1 from bilateral pulmonary artery thromboemboli, 1 in a patient who complained of not being able to breathe), 1 ongoing pregnancy treated with uterine aspiration, 2 blood transfusions, 1 sepsis (with death), 1 hospitalization, and 3 ED or urgent care visits related to medical abortion. On October 12, 2021 the Applicant provided information from July 1, 2021 to September 30, 2021; there were no additional adverse events. For the period between January 27, 2020 and September 30, 2021, there were four adverse events reported corresponding to the above cases from FAERS identified by (b) (6) case 3 (ongoing pregnancy), case 5 (death unknown cause), case 6 (sepsis and death), and case 7 (pulmonary embolism).^r

The postmarketing data from FAERS were analyzed by (b) (6) to determine if there was a difference in adverse events between periods when the in-person dispensing requirement was being enforced and periods when the in-person dispensing requirement was not being enforced. Based on this review, we conclude that there does not appear to be a difference in adverse events between periods when the in-person dispensing requirement was being enforced and periods when the in-person dispensing requirement was not being enforced. This suggests that mifepristone may be safely used without an in-person dispensing requirement.

^r The eighth FAERS case, oral pain/soreness, was not within the scope of the August 5, 2021 IR and was not considered for this review of postmarketing safety information submitted by the Applicants in response to the IRs.

(b) (6) review of the Applicants' IR responses, which included the same cases identified by (b) (6) from FAERS, did not change our conclusion.^s

Literature Review

Published studies have described alternatives in location and method for dispensing mifepristone by a certified prescriber (or an equivalent healthcare provider in countries other than the US). Some studies have examined replacing in-person dispensing in certain health care settings with dispensing at retail pharmacies (Grossman², Wiebe⁷, Rocca⁶) and dispensing mifepristone from pharmacies by mail (Grossman¹, Upadhyay¹⁴, Hyland¹⁵). Other studies have evaluated two modes of dispensing by prescribers: (1) prescribers mailing the medications to women (Gynuity study [Raymond¹⁶, Chong³, Anger¹⁷], Kerestes⁴, Aiken⁵ (2021)) and (2) prescribers using couriered delivery of medications (Reynolds-Wright¹⁸). Other studies have evaluated dispensing mifepristone by mail by an entity described as "a partner organization" (Aiken¹⁹ (2017), Norton²⁰, Endler²¹). For ease of review, in the sections below that describe these studies, we have separated relevant references by the methodology used to dispense mifepristone.

Retail pharmacy dispensing

Three studies report medical abortion outcomes for retail pharmacy dispensing of mifepristone after clinical evaluation. Grossman² conducted a US-based study in which mifepristone and misoprostol were dispensed from a pharmacy partnered with the clinic where the participant had an evaluation by ultrasound and counseling. Of the 266 participants enrolled, 260 had known abortion outcomes. Complete abortion without additional procedure occurred in 243 participants (93.5% of those with known outcomes). Seventeen participants (6.5% of those with known outcomes) were diagnosed with incomplete abortion and underwent uterine aspiration. The reported proportion of complete abortion is within the range described in the approved mifepristone labeling. However, the finding represents a lower-than-expected efficacy based on the cohort's GA (84% of participants were at ≤ 56 days GA, a cohort for which the labeled success rate is 96.8%). No participants experienced a serious adverse event, were hospitalized, or required transfusion. Three participants had ED visits with treatment (intravenous hydration, pain medication, pelvic infection after uterine aspiration for incomplete abortion). The study's

^s The reporting period of (b) (6) assessment of the adverse events in FAERS is not identical to the time period for summaries of adverse events in the IRs to the Applicants. Therefore, the numbers of cases and adverse events summarized in (b) (6) assessment may differ from the numbers of cases and adverse events summarized by the Applicants in their responses to IRs (note that each case report may include more than one adverse event).

safety and efficacy outcomes are consistent with labeled frequencies. The majority of participants (65%) were very satisfied with the experience. There were some complaints from participants about not receiving all prescribed medications at the initial pharmacy visit, privacy not being adequately maintained, and perceived negative pharmacist attitude.

Overall, we conclude that this study has limited generalizability because it was conducted in two US states and involved partnered pharmacies, some of which were in the same building as the clinic. Additionally, all participating pharmacies in this study were required to have a pharmacist on duty during clinic hours who had been trained in the study protocol and was willing to dispense mifepristone. The study conditions may not be generalizable to US retail pharmacies; there is insufficient information to assess this. Rocca⁶ conducted an observational study evaluating 605 participants at ≤ 63 days GA who obtained medical abortions in Nepal by comparing the provision of medical abortion service by newly trained nurse midwives in pharmacies to medical abortion provided in government-certified clinics. Participants who presented to pharmacy study sites underwent clinical screening including a pelvic exam by trained nurse midwives at the pharmacy (which was equipped with an examination room) and if eligible for medical abortion, were dispensed mifepristone and misoprostol in the pharmacy at the time of their visit. Participants who presented to public health facilities underwent clinical screening including pelvic examination by abortion providers including trained nurse midwives and if eligible for medical abortion were dispensed mifepristone and misoprostol in the clinic at the time of their visit. The authors reported that, with respect to complete abortion (>97%) and complications (no hospitalizations or transfusions), evaluation and dispensing in pharmacy was non-inferior to in-clinic evaluation and dispensing.

Wiebe,⁷ in a retrospective, chart review study conducted in Canada, compared abortion outcomes of 182 women at ≤ 70 days GA who underwent medical abortion with telemedicine consult, and either received medications by courier or picked them up at a local pharmacy, with outcomes of a matched control cohort of 199 women who received the medications at a pharmacy after an in-clinic visit. The groups had similar documented complete medical abortion outcomes (90%, calculated maintaining subjects with unknown outcomes in the denominator; $\geq 95\%$ calculated with known outcomes only). The telemedicine group had one case of hemorrhage (0.5%) and one case of infection requiring antibiotics (0.5%) compared with no cases of hemorrhage or infection requiring antibiotics in the in-clinic cohort. The telemedicine group had more ED visits (3.3% compared to 1.5% in-clinic cohort). Both models of dispensing mifepristone resulted in efficacy and safety outcomes within labeled frequency.

None of the three studies described above allow a determination regarding differences in safety between in-person dispensing by a certified prescriber in a health care setting and dispensing through a retail pharmacy, due to limitations on the generalizability of the studies to the current retail pharmacy environment in the US. The outcome findings from the one US study (Grossman²), in which the pharmacies were partnered with prescribers, may not be generalizable to much of the US as they do not reflect typical prescription medication availability with use of retail pharmacy dispensing. Although retail pharmacy dispensing of mifepristone and misoprostol in Canada has been described in the literature, there are important differences in healthcare systems between Canada and the US that render the findings from studies in Canada (Wiebe⁷) not generalizable to the US. In the Wiebe study, timely provision of medication from the retail pharmacy was accomplished by either courier to the woman or faxed prescription to the woman's pharmacy. It is unknown whether conditions that allow timely access to medications for medical abortion would occur in retail pharmacies throughout the US. Canada's federal government has reaffirmed that abortion is an essential health service^t which may have implications affecting access to medical abortion from retail pharmacies in Canada. The Rocca⁶ study evaluated medical abortion provided in Nepali pharmacies and essentially moved the abortion provider and clinical examination into the pharmacy, a scenario that is not, at this time, applicable to the US retail setting.

Mail order pharmacy

Grossman¹ published an interim analysis of an ongoing prospective cohort study evaluating medical abortion with mifepristone and misoprostol dispensed by mail-order pharmacy after in-person clinical assessment. All participants were evaluated for eligibility during a clinic visit with GA up to 63 days confirmed with either an ultrasound or examination; instead of receiving medication at the clinic visit, participants received medications from a mail-order pharmacy. A total of 240 participants have been enrolled; three participants did not take either medication. A total of 227 (94.6%) provided some outcome information, of whom 224 provided abortion outcome information. Complete abortion without additional procedures occurred in 217 participants (96.9% of those with known outcomes). Two (0.9%) participants experienced serious adverse events (SAE); one received a blood transfusion, and one was hospitalized overnight. Nine (4%) participants attended 10 ED visits. In this interim analysis, the outcomes are consistent with labeled frequencies. With respect to the time interval between a

^t As noted in Mark²³ and Martin²⁴, most provincial and federal health insurance programs in Canada cover medical abortion, and covered services are free at the point of care.

participant's clinic visit and receipt of medications, of the 224 participants with known abortion outcomes, 184 (82.1%) received medication within 3 days. However, 17% received between 4-7 days and one participant waited over 7 days for receipt. Seven of 216 (3.2%) participants who completed the day-3 survey reported compromised confidentiality (e.g., someone found their medication, privacy concerns).

Upadhyay¹⁴ reports findings from a retrospective cohort study of 141 women undergoing medical abortion in the US without a consultation or visit. Eligibility was assessed based on a participant-completed online form collecting pregnancy and medical history. Participants who were considered eligible received medication delivered by a mail-order pharmacy. Three interactions via text, messaging or telephone occurred to confirm medication administration, assessment of expulsion and pregnancy symptoms, and results of a 4-week home pregnancy test. Abortion outcome was determined by either the day 3 assessment or the 4-week pregnancy test. The investigators reported a complete abortion rate without additional procedures of 95% (105 participants out of 110 for whom outcomes were known) and stated that no participants had any major adverse events. The proportion of abortion outcomes assessed at 3 days versus 4 weeks is not reported. Regardless, determining outcomes at 3 days is insufficient to determine outcome rates or safety findings because a 3-day follow-up period is too short. Additionally, a substantial number of participants (31) provided no outcomes information. Among the 141 participants enrolled, 128 had any follow-up contact with the study staff, and 110 provided outcomes information. Excluding outcomes of 22% of the cohort is a limitation of this study. This study used a model with numerous deviations from standard provision of medical abortion in the US, such as no synchronous interaction with the prescriber during informed consent or prior to prescribing medication, no confirmation of self-reported medical, surgical, and menstrual history. Further, follow-up information based on a 3-day period is insufficient to determine outcome rates or safety findings. These deviations, limited follow-up information, and small sample size limit the usefulness of this study.

Hyland¹⁵ describes findings from a cohort study in Australia evaluating medical abortion outcomes utilizing telemedicine and a central mail order pharmacy. All participants obtained screening tests including ultrasound confirmation of GA. A total of 1010 participants completed the screening process and were provided mifepristone and misoprostol. Abortion outcomes were determined for 754 (75%) of the 1010. Outcomes for the remaining 256 participants (25%) were not included because 31 provided no relevant information after shipment, 14 reported not taking misoprostol, and 211 did not have "full follow up" (i.e., known outcome of either complete medical abortion, uterine evacuation, or ongoing pregnancy with plan to continue).

Complete abortions without additional procedures occurred in 727 participants (96% of those with definitively documented outcomes) and is consistent with labeled efficacy. Of the 754 participants included in the analysis 717 (95%) had no face-to-face clinical encounters after medications were mailed while 21 (3%) were admitted to the hospital and 16 (2%) had an outpatient encounter. One participant who was hospitalized and underwent a surgical uterine evacuation received a transfusion. Not included in the findings are 7 hospitalizations occurring in 7 participants who did not have “full follow up”. The authors do not report any other adverse events and conclude use of the telemedicine medical abortion service is safe. The reasons for hospitalization are not discussed by the authors; therefore, it is unknown why the patients were hospitalized. Although the reported number of hospitalizations (3%) is higher than the less than 1% in the FDA-approved mifepristone labeling, conclusions regarding the safety findings in this study cannot be made in the absence of information about the reasons for hospitalization. Other limitations of this study include incomplete information about outcomes with face-to-face encounters, and not reporting outcomes of 25% of the enrolled cohort.

Overall, the three studies evaluating mail order pharmacy dispensing suggest that the efficacy of medical abortion is maintained with mail order pharmacy dispensing. In the Grossman¹ study, the interim analysis, although small, does not raise serious safety concerns. We note that 18% of participants did not receive medications within 3 days; the potential for delay in receiving medication by mail could limit the GA eligible for medical abortion through mail order pharmacy dispensing, because women at GA closer to 70 days might not receive medication in time. A small proportion (3%) of participants raised concerns regarding the issues of confidentiality and privacy. Safety findings from the Hyland¹⁵ study are difficult to interpret. Although only one transfusion is reported, and the authors state the findings demonstrate safety, the higher hospitalization rates, and lack of information on the reasons for hospitalization do not allow any conclusions about safety findings. Lastly, the Upadhyay¹⁴ study had no reported adverse events, but the findings are less useful because of the limited follow-up, and because medical abortions were provided using a model with numerous deviations from standard provision of medical abortion in the US.

Clinic dispensing by mail

A total of five studies evaluated clinic dispensing by mail.^{3,4,5,16, 17} Gynuity Health Projects conducted a prospective cohort study (the “TelAbortion” study) evaluating use of telemedicine for remote visits and mifepristone being dispensed from clinics via overnight or regular tracked mail. Three publications reviewed have reported outcomes for the Gynuity population

exclusively: Raymond¹⁶ from May 2016 to December 2018, Chong³ from May 2016 to September 2020 and Anger¹⁷ from March 2020 to September 2020. Due to the pandemic, the Gynuity study deviated from the protocol requirement of confirmation of GA by examination or ultrasound for many participants treated from March 2020 onward (although none of the three publications reported on the single element of dispensing mifepristone from the healthcare setting by mail). A fourth study, Kerestes,⁴ reports outcomes of medical abortion at the University of Hawai'i from April 2020 to November 2020: seventy-five (of whom 71 were enrolled in the Gynuity study) of the 334 participants in Kerestes were dispensed mifepristone by mail after a telemedicine consult. The section below discusses these four studies from the US as well as a large UK study by Aiken⁵ (2021).


Raymond¹⁶ (2019) reported outcomes from the Gynuity study prior to the pandemic. In the TelAbortion study, participants were not required to have an in-person clinic visit; rather, they obtained screening tests at laboratories and radiology offices and then communicated with the abortion provider by videoconference. If the participant was eligible for treatment, the provider dispensed the medications by mail. Of 433 women screened, 165 (38%) either declined to schedule the videoconference or did not keep the videoconference appointment. Among the 268 participants evaluated via videoconference, medication packages were sent to 248. Abortion outcomes were determined for 190 (77%) of the 248; outcomes for 58 (23%) participants were unknown. Complete abortion without additional procedures occurred in 177 participants (93% of those with known outcomes). The investigators obtained follow-up information from 217 participants after package shipment; there were two hospitalizations (one received a transfusion for severe anemia despite having had a complete abortion), and 16 other participants (7%) had clinical encounters in ED and urgent care centers. The reported outcomes in Raymond¹⁶ (2019) are similar to outcomes described in approved labeling except the combined ED/urgent care center encounters (7%) exceeded the ED visits in approved labeling (2.9-4.6%). The authors note that half of the ED/urgent care visits did not entail any medical treatment and opine that the increased number of visits may have been due to the study participants living farther from the abortion providers.¹⁶ All participants received medications within 8 days.

Chong³ updated the findings from the Gynuity study described in Raymond¹⁶ and reported on 1157 medical abortion outcomes, of which approximately 50% occurred during the period of the COVID-19 PHE. Although a screening ultrasound was required per the protocol, sites determined in 52% (346/669) of abortions that occurred during the period of the COVID-19 PHE that, in order to avoid potential exposure to COVID-19 at a health care facility, those

participants were not required to obtain a screening ultrasound. Use of urine pregnancy test to confirm abortion completion also increased from 67% (144/214) in the 6 months prior to the pandemic to 90% (602/669) in the 6 months during the pandemic. Of the 1390 participants to whom medicine packages (containing both mifepristone and misoprostol) were mailed, 1157 (83.2%) had known abortion outcomes. Complete abortion without a procedure occurred in 1103 participants (95% of the those with a known outcome). Ten women experienced an SAE (5 transfusions (0.4%) and 7 hospitalizations (0.7%)) and 70 (6%) participants had unplanned clinical encounters in ED/urgent care. Surgical interventions were required in 47 participants (4.1% of 1390) to complete abortion. The reported outcomes in this study are similar to outcomes described in approved labeling, except that the combined ED/urgent care center encounters (6%) exceeded the ED visits in approved labeling (2.9-4.6%).

Anger¹⁷ compared outcomes among participants enrolled in the Gynuity study who did versus did not have confirmation of GA/intrauterine location with an examination or ultrasound from 10 jurisdictions across the US. These participants were screened for enrollment from March 25 through September 15, 2020. All participants had a telemedicine consultation and received mifepristone and misoprostol by mail from the healthcare facility. Determination of which participants did not require confirmation of GA by examination or ultrasound to be eligible depended on the study clinician's assessment of eligibility for "no-test medication abortion"^u based on a sample protocol published by Raymond²² (2020). There were two key differences between the two groups. Participants for whom the study clinician determined a pre-abortion ultrasound was required were more likely than the participants who had no ultrasound or examination to live further than 150 miles from the clinic (51.2% vs. 31.7%) and were more likely to have a GA above 63 days (12.0% vs. 1.7%). The study sites shipped 503 medication packages during the analysis period; 344 packages went to the "no test" group while 159 went to the "test" medical abortion cohort (see figure below). However, because the two cohorts were not randomized in this study, they had different baseline characteristics. Consequently, findings based on the comparisons between the two cohorts should be interpreted carefully.

^u "No-test medication abortion" refers to medical abortion provided without a pretreatment ultrasound, pelvic examination, or laboratory tests when, in the judgment of the provider, doing so is medically appropriate (appropriateness based on history and symptoms); "no-test medication abortion" does include post-abortion follow up. A sample protocol is described by Raymond et al.²²



Source: Figure 1 in this publication. MA= medical abortion.

The investigators' analyses excluded 91 (18% of 503; 57 in the no-test group and 34 in the test group) participants because they did not provide a date of the last menstrual period (LMP), did not take mifepristone, or did not have a recorded abortion outcome. Overall, 410 participants (81.5% of 503) provided outcomes data. There were no reported ectopic pregnancies in either group. The number of ED/urgent care visits and the proportion of unplanned clinical encounters that led to medical treatment were not reported. In the no-test group, complete medical abortion was confirmed in 271 participants who took medications (94% among those with known outcome). In the no-test cohort, two participants were "hospitalized and/or blood transfusion," and 36 (12.5%) had an unplanned clinical encounter (participant sought in-person medical care related to abortion and the visit was not planned prior to abortion).

In the test medical abortion group, complete abortion was confirmed in 123 participants (of 125 with known outcomes); the completion rate was 98% among those with known outcomes. In the test medical abortion group, one participant was "hospitalized and/or blood transfusion," and 10 (8.0%) had an unplanned clinical encounter. The authors concluded that, compared to participants who had an ultrasound prior to medical abortion, those without an examination prior to medical abortion were more likely to require procedural interventions and had more unplanned clinical encounters.

Kerestes⁴ was the only publication that linked outcomes of medical abortion with different delivery models. Participants included in the report had GA up to 77 days and received

medications in Hawaii between April 2020 and January 2020. A total of 334 medication packages (to 330 unique participants) were dispensed containing mifepristone and misoprostol; three different delivery models were used concurrently: 110 (32.9%) had traditional in-person visits, 149 (44.6%) had telemedicine consultation with in-person pick-up of medications, and 75 (22.5%) were sent medications by mail (71 of these were enrolled through Gynuity's TelAbortion study). Seven participants of the 330 participants who received 334 medication packages reported that they did not take them and were excluded from analysis of the outcomes. Among participants with follow-up data, the rates of successful medical abortion without surgery were 93.6%, 96.8%, and 97.1% in the in-clinic group, telemedicine + in-person pickup group, and telemedicine + mail group, respectively; these were consistent with outcomes in approved labeling. Blood transfusion was given to two participants (both in the telemedicine + in-person pickup group). Eleven participants went to an ED. Although ED visits occurred the most frequently in the telemedicine + mail group (four participants or 5.8%) and the least in the in-person group (two participants or 2.1%), the study reported no increases in other serious adverse events.

Taken together, the three Gynuity study reports^{3,16,17} and Kerestes⁴ support dispensing mifepristone and misoprostol by mail after a telemedicine visit. Efficacy was maintained in all four studies. All of the studies reported SAEs frequencies comparable to labeled rates, except two of the Gynuity study reports (Raymond¹⁶, Chong³) and Kerestes⁴ report a higher frequency of ED/urgent care visits than the labeled frequency of ED visits. We do not know whether the reporting of combined ED and urgent care visits represents an increased rate of ED visits compared to the labeled rate of ED visits (2.9-4.6%). Other labeled SAEs (e.g., transfusion) occur infrequently (< 1%).

Aiken⁵ (2021) reports outcomes of medical abortion up to 70 days GA in the UK before and during the pandemic in a retrospective cohort study. In the UK, prior to the COVID-19 pandemic, all patients attended an in-clinic visit where they received an ultrasound, were administered mifepristone in the clinic, and given misoprostol in-clinic for use at home (traditional model). During the pandemic, medical abortion consultations were performed remotely by telephone or video. Based on the consultation and questionnaire (including date of last menstrual period; menstrual, contraceptive and medical history; symptoms; risk for ectopic pregnancy), an assessment of eligibility for treatment via telemedicine was made. If eligible, medications were delivered to participants via mail or were made available for collection from the clinic for use at home. If the participant was assessed to be ineligible for treatment via

telemedicine, an in-person assessment with ultrasound was performed and medications were provided from the clinic for home use (hybrid model).

The study compared the two cohorts: 22,158 obtained medical abortion before the pandemic and had in-person visits and dispensing (traditional model) and 29,984 obtained medical abortion during the pandemic with either in-person visit and in-person dispensing, or a telemedicine visit and dispensing by mail or picked up from the clinic (hybrid model). Outcomes were obtained from electronic records and incident databases. Outcomes of all hospitalizations related to abortion, ED visits, infection without sepsis, and hemorrhage without transfusion were not reported. The investigators' analysis for non-inferiority determined the efficacy and safety were comparable between both cohorts. Complete abortion occurred in > 98% in both cohorts. Hemorrhage requiring transfusion was reported in 0.04% and 0.02% of the traditional and hybrid cohorts, respectively; this is lower than the labeled 0.5% transfusion rate. There were no severe infections requiring hospitalization, major surgery or deaths reported.

A secondary analysis of the hybrid cohort was reported. Within the 29,984-person hybrid model cohort, 11,549 (39%) abortions were conducted in-person (in-person assessment with ultrasound was performed and medications provided from the clinic for home use) and 18,435 (61%) abortions were provided by telemedicine visit, without tests or confirmation of GA/intrauterine position by ultrasound, and medications either mailed or picked up from the clinic. Outcomes stratified by type of mifepristone dispensing were not reported. The rate of complete abortion was slightly higher in the telemedicine group (99.2%) than that in the in-person group (98.1%). There were no significant differences in the rates of reported SAEs. Adjustments for clinical and demographic characteristics were made because the two groups differed in baseline characteristics, including a higher proportion of pregnancies with GA over 6 weeks in the in-person group (68.2% compared with 55.1%). The authors conclude a hybrid model for medical abortion that includes no-test medical abortion^u (no ultrasound, no pelvic exam, no pregnancy test) is effective and safe.

We conclude that although the Aiken⁵ (2021) study has a large sample size and includes 85% of all medical abortions performed in England and Wales during the study period, the study has limitations. The authors acknowledge the main limitation of their study was that analysis was based on deidentified information in the NHS database and the investigators were unable to verify the outcomes extracted. Other limitations included that their search only captured

outcomes in electronic records and incident databases that met the authors' defined threshold for SAE reporting, and that the labeled abortion outcomes considered serious, such as hospitalizations related to abortion, infection without sepsis, hemorrhage without transfusion, or ED/urgent care visits, were not all included in the authors' definition of serious adverse event.

Data from the mail order dispensing studies with telemedicine visits from Gynuity (Raymond, Chong and Anger),^{3,16,17} Kerestes⁴, and Aiken⁵ (2021) support that efficacy of medical abortion was maintained. The Aiken⁵ study appears to be of sufficient sample size to determine whether safety outcomes with mail dispensing differ from in-person dispensing; however, the study's design did not capture all serious safety outcomes, thus limiting the certainty of the findings. Study reports of Raymond¹⁶ Chong³, and Kerestes⁴ all suggest there may be an increase in ED/urgent care visits with telemedicine visits and dispensing by mail without increases in other adverse events. Anger's¹⁷ comparative analysis suggests a pre-abortion examination may decrease the occurrence of procedural intervention and decrease the number of unplanned visits for postabortion care. Overall, despite the limitations noted, these studies support that dispensing by mail is safe and effective. Although the literature suggests there may be more frequent ED/urgent care visits related to the use of mifepristone when dispensed by mail from the clinic, there are no apparent increases in other SAEs related to mifepristone use. One reason for the increase in frequent ED/urgent care visits in the Raymond¹⁶ publication, according to its authors, may have been that a substantial proportion of participants lived significant distances from their providers and increased distances have been associated with higher use of ED following treatment. Raymond¹⁶ reported that half of the participants who had an ED/urgent care visit did not require medical treatment.

Clinic dispensing by courier

Reynolds-Wright¹⁸ reported findings from a prospective cohort study of 663 women at less than 12 weeks' GA in Scotland undergoing medical abortion at home with use of telemedicine during the pandemic (from April 1 to July 9, 2020). The majority of medical abortions (78.7%) used telemedicine visits, eliminated pre-abortion ultrasound, and provided mifepristone for pick up at the service or by couriered delivery to woman's home. The number of couriered deliveries was not reported; thus, this study does not provide abortion outcomes separately for couriered delivery of mifepristone and misoprostol. With access to NHS regional hospital databases, the investigators were able to verify pregnancy outcomes and complications. Of the 663 participants, 642 (98.2%) were under 10 weeks GA, 21 (1.8%) were between 10 and 12 weeks

GA, and one participant was never pregnant. A total of 650 participants had complete abortion without requiring surgical intervention (98%), 5 (0.8%) an ongoing pregnancy and 4 (0.6%) an incomplete abortion. The outcomes from this study in Scotland are consistent with labeled mifepristone outcomes. The study shares the same limitations as the Aiken⁵ (2021) study.

Partner organization dispensing by mail

Women on Web (WoW), an internet group, connects patients and providers outside of the US and provides medical abortion globally, dispensing mifepristone through “a partner organization” by mail.^v Medical abortion eligibility is determined using an online questionnaire with asynchronous physician review. If eligible, medications are mailed to the women. WoW provides help and support by email or instant messaging.

Aiken¹⁹ (2017) conducted a population-based study analyzing findings from 1,636 women in the Republic of Ireland and Northern Ireland who were sent medications between 2010 and 2012. Receipt of medications was confirmed for 1,181 women, among whom 1,023 confirmed use of mifepristone and misoprostol; outcome information was available for 1,000 (61% of women sent medications). Of the 1,000 women, the majority (781, 78%) were less than 7 weeks GA and 219 (22%) were at 7-9 weeks. Complete abortion without surgical intervention occurred in 947 (94.7% of 1,000 with known outcome); 7 (0.7%) women received a blood transfusion, 26 (2.6%) received antibiotics (route of administration undetermined) and 87 (8.7%) sought medical care at a hospital or clinic for symptoms related to medical abortion. Hospitalizations related to abortion were not reported. The reported proportion of complete abortion is within the range labeled for medical abortion up to 70 days (92.7-98.1%). However, the finding of 94.7% complete abortion represents a lower-than-expected efficacy based on the cohort’s GA (almost 80% less than 7 weeks, labeled success for medical abortion $\leq$ 49 days is 98.1%). This study has limitations, including outcomes based on self-report without validation of completed abortion by examination or laboratory testing, and no known outcomes for 39% of study cohort. Additionally, the authors noted medical abortion was provided in a legally-restrictive setting, where the law provided a maximum penalty of life imprisonment for the woman undergoing the abortion, which may affect participants’ self-reporting.

^v In March 2019, FDA sent a WL to Aidaccess.org, a group affiliated with WoW. Aidaccess.org received this WL because it was introducing misbranded and unapproved new drugs into the U.S. In the context of this REMS review, studies involving WoW are included solely for purposes of evaluating of data regarding the methods of dispensing mifepristone.

Endler²¹ and Norten²⁰ have reported outcomes from WoW cohorts but do not provide relevant information on mifepristone dispensing by mail, because neither provide meaningful outcomes data for consideration. Endler²¹ compared the outcomes of self-reported heavy bleeding and clinical visits occurring during the “first or second day of abortion” that occurred in women undergoing medical abortion at 9 weeks GA or less, with outcomes from women at more than 9 weeks GA. Outcome data from day 1 or 2 is of limited usefulness. Norten²⁰ describes findings from a survey of women who were sent medical abortion medication through WoW and provided self-reported outcomes. Results were based on surveys returned from only 37% of participants, a return rate that is too low for the study to be considered valid.

WoW uses a model with numerous deviations from the standard provision of medical abortion in the US. For example, this model has no synchronous interaction with the prescriber during informed consent or prior to prescribing medication and no confirmation of self-reported medical, surgical, and menstrual history or confirmed pregnancy testing. Further, although Aiken¹⁹ (2017) is a large cohort study, the outcomes are self-reported with no verification of complete abortion by laboratory or clinical evaluation and 39% of outcomes are unaccounted for. These limitations in the Aiken study result in the data being insufficient to determine the safety of dispensing mifepristone by mail through a partner organization.

4. Discussion

After review of the published literature, safety information collected during the COVID-19 PHE, postmarketing data, information from the first Mifepristone REMS Program assessment report, responses to information requests to the Applicants, and information provided by advocacy groups, individuals and the plaintiffs in the *Chelius v. Becerra* litigation, we conclude that the REMS can be modified to reduce burden without compromising patient safety.

Prescriber Certification

None of the publications we reviewed would support a conclusion that a healthcare provider who prescribes mifepristone does not need to meet the qualifications included in the Mifepristone REMS Program as described above in section 3.2.1. Absent these provider qualifications, serious complications associated with medical abortion, including missed ectopic pregnancy and heavy bleeding from incomplete abortion, would not be detected or appropriately managed.

We conclude that prescriber certification (ETASU A) should be maintained. The current process requires the prescriber to agree to the requirements of the Mifepristone REMS Program and to attest that they meet the qualifications described in section 3.2.1 above. The REMS has been structured to minimize burden to prescribers by requiring only a one-time certification by the prescriber for each Applicant. We have determined that healthcare provider certification continues to be necessary to ensure the benefits outweigh the risks, especially considering that, if the in-person dispensing requirement is removed from the Mifepristone REMS Program, the number of new providers may increase (see discussion in section 3.2.2 above).

Drug to be dispensed with evidence or other documentation of safe use conditions

The requirement to counsel the patient and provide them with the *Patient Agreement Form* ensures that each patient is informed of the appropriate use of mifepristone, the risks associated with treatment, and what to do if they experience symptoms that may require emergency care.

In 2016, we initially recommended eliminating the *Patient Agreement Form* (see section 3.2.2), though the form was ultimately maintained as part of the REMS. As discussed above, our current literature review has indicated that there is no basis to remove the *Patient Agreement Form* from the REMS. In addition, surveys we reviewed suggest that if the in-person dispensing requirement for mifepristone is removed, there could be a potential doubling of medical abortion providers. This potential doubling of medical abortion providers supports the continued need to ensure that patients are consistently provided patient education under the Mifepristone REMS Program regarding the use and risks of mifepristone. The *Patient Agreement Form* is an important part of standardizing the medication information that prescribers communicate to their patients, including new prescribers, and also provides the information in a brief and understandable format to patients. We determined, in accordance with section 505-1(f)(2) of the FD&C Act, that this does not impose an unreasonable burden on providers or patients.^w

Given the likelihood of a potential increase in new prescribers if the in-person dispensing requirement is removed from the Mifepristone REMS Program, we conclude that maintaining the *Patient Agreement Form* remains necessary to assure safe use at this time.

^w The *Patient Agreement Form* can be signed in person or through other means.

Drug to be dispensed only in certain healthcare settings

As discussed above in section 3.2.3, our evaluation of information submitted by the applicants in the one-year (1st) REMS assessment report for the Mifepristone REMS Program and in response to follow-up requests from the Agency indicates that the number of adverse events reported to FDA during the COVID-19 PHE with mifepristone use is small, and the data provide no indication that any program deviation or noncompliance with the Mifepristone REMS Program contributed to these adverse events. We further conclude, based our review of the postmarketing safety data from FAERS during the COVID-19 PHE and information submitted by the applicants for the timeframe of January 27, 2020 through September 30, 2021, that there does not appear to be a difference in adverse events between periods during the COVID-19 PHE when the in-person dispensing requirement was being enforced and periods when the in-person dispensing requirement was not being enforced; nor have we identified any new safety concerns with the use of mifepristone for medical termination of early pregnancy.

Alternatives to in-person dispensing of mifepristone have been investigated in several studies and countries. The literature review identified 15 publications^x that assessed safety outcomes from various medication delivery models (US, UK, Canada, Ireland, Australia, Nepal), including dispensing by retail and mail order pharmacies, prescribers mailing medications or using couriered service to deliver medications, and dispensing by “partner organizations”. The ability to generalize the results of these studies to the US population is hampered by differences in pre-abortion care (e.g., telemedicine versus in-person, testing), and the usefulness of the studies is limited in some instances by small sample sizes and lack of follow-up information on outcomes with regard to both safety and efficacy.

In addition, there are factors which complicate the analysis of the dispensing element alone. Some of these factors are: (1) only a few studies have evaluated alternatives for in-person dispensing of mifepristone in isolation; for example, most studies on mail dispensing of mifepristone also include telemedicine consultation, and (2) because most SAEs with medical abortion are infrequent, though they can be life threatening, further evaluation of changes in dispensing would require studies with larger numbers of participants. We did not find any large clinical studies that were designed to collect safety outcomes in healthcare systems similar to the US.

^x The 15 publications correspond to endnote numbers: 1-7, 14-21.

Based on the literature identified by our review, dispensing mifepristone by mail from the clinic or from a mail order pharmacy does not appear to jeopardize the efficacy of medical abortion. The studies we reviewed are not adequate on their own to establish the safety of the model of dispensing mifepristone by mail, although the safety and efficacy outcomes reported in these studies remain within the ranges described in mifepristone labeling except for increased numbers of ED/urgent care visits and hospitalizations.

Four publications (Raymond¹⁶, Chong³, Anger¹⁷ and Kerestes⁴), describe a relevant US cohort where dispensing mifepristone from the clinic by mail was paired with telemedicine visits. These studies showed that efficacy was maintained and there was no increased frequency of SAEs except for higher ED/urgent care visits. The increased ED/urgent care visits were not associated with increases of other SAEs, and in the view of one study's authors (Raymond¹⁶), may be associated with participants being located significant distances from their providers. The Aiken⁵ (2021) study of a large UK cohort where the clinics mailed mifepristone report small (lower than labeled) occurrences of transfusion and no significant infections requiring hospitalization. In Grossman¹ and Hyland¹⁵, where the pharmacies mailed mifepristone after prescribers confirmed GA, efficacy is maintained. Grossman's¹ interim analysis found no increases in SAEs. Hyland¹⁵ reported higher numbers of hospitalizations but did not report increases of other SAEs. Overall, while the studies assessing mifepristone dispensing by mail suggest more frequent encounters with healthcare providers, they generally support a conclusion that dispensing by mail is safe. Despite the limitations of the studies we reviewed, we conclude that overall, the outcomes of these studies are not inconsistent with our conclusion that, based on the 1st year REMS assessment report and postmarketing safety data, mifepristone will remain safe, and efficacy will be maintained if the in-person dispensing requirement is removed from the Mifepristone REMS Program.

Based on the REMS assessment data, FAERS data from the time period when the in-person dispensing requirement was not being enforced, our review of the literature, and information provided by advocacy groups, individuals, the Applicants, and the plaintiffs in the *Chelius v. Becerra* litigation, we conclude that mifepristone will remain safe and effective for medical abortion if the in-person dispensing requirement is removed, provided all the other requirements of the REMS are met, and pharmacy certification is added as described below.

Removing the in-person dispensing requirement will render the REMS less burdensome to healthcare providers and patients and provided all other requirements of the REMS are met, including the additional requirement for pharmacy certification, the REMS will continue to

ensure that the benefits of mifepristone for medical abortion outweigh the risks. Therefore, to reduce the burden imposed by the REMS, the Mifepristone REMS Program should be modified to remove the in-person dispensing requirement, which would allow, for example, dispensing of mifepristone by mail via certified prescribers or pharmacies, in addition to in-person dispensing in clinics, medical offices and hospitals as currently outlined in ETASU C.

New requirement to be added for pharmacy certification

The current distribution model requires the certified prescriber to dispense mifepristone directly to the patient in a clinic, medical office, or hospital. During the periods when the in-person dispensing requirement was not being enforced, both applicants used mail order pharmacies to receive and hold mifepristone on behalf of the certified healthcare providers who had purchased the product.^{j,y,z} Pursuant to a prescription for mifepristone, the mail order pharmacy would ship the product to a named patient.

The Mifepristone REMS Program continues to require that mifepristone be prescribed only by certified prescribers. With the removal of the in-person dispensing requirement, however, the drug is no longer required to be dispensed only in a clinic, medical office or hospital. Under the REMS as modified, mifepristone can be dispensed through a pharmacy, provided the product is prescribed by a certified prescriber and all other requirements of the REMS are met. Given this modification to the dispensing requirements in the REMS, it is necessary to add a requirement for certification of pharmacies under ETASU B. Adding the pharmacy certification requirement incorporates pharmacies into the REMS, ensures that pharmacies are aware of and agree to follow applicable REMS requirements, and ensures that mifepristone is only dispensed pursuant to prescriptions that are written by certified prescribers. Without pharmacy certification, a pharmacy might dispense product that was not prescribed by a certified prescriber. Adding pharmacy certification ensures that ETASU A is met prior to dispensing the product to a patient; certified prescribers, in turn, have agreed to meet all the conditions of the REMS, including ensuring that the *Patient Agreement Form* (ETASU D) is completed. In addition, wholesalers and distributors can only ship to certified pharmacies. Based on our review of the safety data and our consideration of the distribution model implemented by the Applicants during the periods

y ANDA 091178: September 23, 2021 response to the September 15, 2021 information request; October 11 and 16, 2021 responses to the June 30, 2021 and July 15, 2021 information requests; October 26, 2021 response to the October 22, 2021 information request; October 29, 2021 response to the October 27 information request.

z NDA 020687: September 20, 2021 response to the September 15, 2021 information request; October 26, 2021 response to the October 22 information request.

when the in-person dispensing requirement was not being enforced, as well as REMS assessment data and published literature, we conclude that provided all other requirements of the REMS are met, the REMS program, with the removal of the in-person dispensing requirement and the addition of a requirement for pharmacy certification, will continue to ensure the benefits of mifepristone for medical abortion outweigh the risks while minimizing the burden imposed by the REMS on healthcare providers and patients. As modified, the REMS would allow, for example, dispensing by mail order or specialty pharmacies, similar to the distribution model used by applicants during the periods when the in-person dispensing requirement was not being enforced.^{aa}

The above recommendations were discussed with the (b) (6) (b) (6) and senior leadership from CDER on November 2, 2021. The (b) (6) (b) (6) along with senior CDER leadership, concurred with removing the in-person dispensing requirement provided that all of the remaining REMS requirements are met, including but not limited to prescriber certification where prescribers need to attest to having certain qualifications, and maintaining the *Patient Agreement Form*. The (b) (6) (b) (6) and senior leadership from CDER were also in favor of adding pharmacy certification to assure the safe use of mifepristone.

5. Conclusions and Recommendations

Based on the results of REMS assessments; our review of safety data collected during the PHE as well as data from FAERS; our literature search; and information provided by advocacy groups, individuals, the Applicants, and the plaintiffs in the *Chelius v. Becerra* litigation, (b) (6) and (b) (6) have concluded that a REMS modification is necessary and should include the following changes:

- Removing the requirement under ETASU C that mifepristone be dispensed only in certain healthcare settings, specifically clinics, medical offices, and hospitals.
- Adding a requirement under ETASU B that pharmacies that dispense the drug be specially certified.

^{aa} Our current conclusion that the REMS would allow dispensing by mail order or specialty pharmacies is based on data received from Applicants relating to the periods when the in-person dispensing requirement was not enforced and mail-order pharmacies were used to dispense the product, as well as our analysis of postmarketing safety data and available literature. At this time we do not have data (from the Applicants or from other sources) to assess the certification of retail pharmacies under the REMS. We have not yet determined the details of pharmacy certification requirements, including whether any limitations on the types of pharmacies that may dispense the product are necessary.

(b) (6) and (b) (6) recommend the Applicants be issued a REMS Modification Notification Letter that requests submission within 120 days from the date of the letter.

6. References

- ¹ Grossman D, Raifman S, Morris N, et.al. Mail-order pharmacy dispensing of mifepristone for medication abortion after in-person clinical assessment. *Contraception* 2021; In press.
doi:<https://doi.org/10.1016/j.contraception.2021.09.008>
- ² Grossman D, Baba CF, Kaller S, et al. Medication Abortion With Pharmacist Dispensing of Mifepristone. *Obstet Gynecol* 2021;137:613–22.
- ³ Chong E, Shochet T, et al. Expansion of a direct-to-patient telemedicine abortion service in the United States and experience during the COVID-19 pandemic. *Contraception* 2021;104:43-48.
- ⁴ Kerestes C, Murayama S, et al. Provision of medication abortion in Hawai'i during COVID-19: Practical experience with multiple care delivery models. *Contraception* 2021 Jul;104(1):49-53.
doi:10.1016/j.contraception.2021.03.025. Epub 2021 Mar 28.
- ⁵ Aiken ARA, Lohr PA, et al. Effectiveness, safety and acceptability of no-test medical abortion (termination of pregnancy) provided via telemedicine: a national cohort study. *BJOG* 2021;128:1464–1474.
- ⁶ Rocca CH, Puri M, et al. Effectiveness and safety of early medication abortion provided in pharmacies by auxiliary nurse-midwives: A non-inferiority study in Nepal. *PLoS ONE* 2018 13(1): e0191174. <https://doi.org/10.1371/journal.pone.0191174>
- ⁷ Wiebe ER, Campbell M, et al. Comparing telemedicine to in-clinic medication abortions induced with mifepristone and misoprostol. *Contracept X*. 2020; 2: 100023.
- ⁸ National Abortion Federation 2020 Clinical Policy Guidelines for Abortion Care, available at https://5aa1b2xfmfh2e2mk03kk8rsx-wpengine.netdna-ssl.com/wp-content/uploads/2020_CPGs.pdf
- ⁹ American College of Obstetricians and Gynecologists Committee on Practice Bulletins—Gynecology and the Society of Family Planning. Simultaneously published as ACOG Bulletin

Number 225: Medication abortion up to 70 days of gestation. *Obstet Gynecol* 2020;136(4): e31-e47 and in *Contraception* 2020; 102:225-236.

¹⁰ Jones HE, O'Connell White K, Norman WV, Guilbert E, Lichtenberg ES, Paul M. First trimester medication abortion practice in the United States and Canada. *PLoS ONE* 2017; 12(10): e0186487. <https://doi.org/10.1371/journal.pone.0186487>

¹¹ Grossman D, Grindlay K, Altshuler AL, Schulkin J. Induced abortion provision among a national sample of obstetrician-gynecologists. *Obstet Gynecol* 2019;133:477-483.

¹² Daniel S, Schulkin J, Grossman D. Obstetrician-gynecologist willingness to provide medication abortion with removal of the in-person dispensing requirement for mifepristone. *Contraception*. 2021;104:73-76

¹³ (b) (6) Review of the one-year REMS assessment report for the Mifepristone REMS Program, December 16, 2021.

¹⁴ Upadhyay UD, Koenig LR, Meckstroth KR. Safety and Efficacy of Telehealth Medication Abortion in the US During the COVID-19 Pandemic. *JAMA Network Open*. 2021;4(8):e2122320. doi:10.1001/jamanetworkopen.2021.22320

¹⁵ Hyland P, Raymond EG, Chong E. A direct-to-patient telemedicine abortion service in Australia: Retrospective analysis of the first 18 months. *Aust N Z J Obstet Gynaecol* 2018;58: 335-340.

¹⁶ Raymond E, Chong E, et al. TelAbortion: evaluation of a direct to patient telemedicine abortion service in the United States. *Contraception* 2019;100:173-177

¹⁷ Anger HA, Raymond EG, et al. Clinical and service delivery implications of omitting ultrasound before medication abortion provided via direct-to-patient telemedicine and mail. *Contraception* 2021 Jul 28;S0010-7824(21)00342-5. doi: 10.1016/j.contraception.2021.07.108. Published online.

¹⁸ Reynolds-Wright JJ, Johnstone A, McCabe K, et al. Telemedicine medical abortion at home under 12 weeks' gestation: a prospective observational cohort study during the COVID-19 pandemic. *BMJ Sex Reprod Health* 2021;0:1–6. doi:10.1136/bmjsexrh-2020-200976

¹⁹ Aiken AR, Digon I, Trussell J, et al. Self reported outcomes and adverse events after medical abortion through online telemedicine: population based study in the Republic of Ireland and Northern Ireland. *BMJ* 2017;357:j2011.

²⁰ Norton H, Ilozumba O, Wilkinson J, et.al. 10-year evaluation of the use of medical abortion through telemedicine: a retrospective cohort study. *BJOG* 2021; <https://doi.org/10.1111/1471-0528.16765>.

²¹ Endler M, Beets L, Gemzell Danielsson K, Gomperts R. Safety and acceptability of medical abortion through telemedicine after 9 weeks of gestation: a population-based cohort study. *BJOG* 2019;126:609–618.

²² Raymond EG, Grossman D, Mark A, et.al. Commentary: No-test medication abortion: A sample protocol for increasing access during a pandemic and beyond. *Contraception* 2020;101:361-366

²³ Mark A, Foster A, Perritt J. The future of abortion is now: Mifepristone by mail and in-clinic abortion access in the United States. *Contraception* 2021;104:38-42

²⁴ Martin D, Miller A, Quesnel-Vallee, A, et al. Canada's global leadership on health 1. Canada's universal health care system: achieving its potential. *Lancet* 2018; 391:1718-35

7. Appendix A

References Cited in Letters from Plaintiffs

References cited in letter from <i>Chelius v. Becerra</i> Plaintiffs (September 29, 2021)	
References included in the REMS review	
Aiken A et al. BJOG 2021; 128 (9): 1464-1474	
Chong, et al. Contraception 2021; 104(1) 43-48	
Daniel S. et al. Contraception 2021; 104(1): 73-76	
References excluded from the REMS review	Rationale for Exclusion
Am. Coll. of Obstetricians & Gynecologists, <i>Position Statement: Improving Access to Mifepristone for Reproductive Health Indications</i> (June 2018), https://www.acog.org/clinical-information/policy-and-position-statements/position-statements/2018/improving-access-to-mifepristone-for-reproductive-health-indications	Policy/advocacy statement
House of Delegates, Am. Med. Ass'n., <i>Memorial Resolutions Adopted Unanimously No. 504 (2018)</i> https://www.ama-assn.org/sites/ama-assn.org/files/corp/media-browser/public/hod/a18-resolutions.pdf	Policy/advocacy statement
Cong. Of Delegates, Am. Acad. Of Fam. Physicians, <i>Resolution No. 506 (CoSponsored C) Removing Risk Evaluation and Mitigation Strategy (REMS) Categorization of Mifepristone</i> (May 24, 2018) https://www.reproductiveaccess.org/wp-content/uploads/2019/02/Resolution-No.-506-REMS.pdf	Policy/advocacy statement
Schummers L et al, Contraception 2020; 102(4): 273	Abstract
Upadhyay UD et al.) Obstet & Gynecol 2015; 125: 175	Published prior to March 29, 2016-July 26, 2021 timeframe for current literature review. We note that the extensive literature review conducted as part of the 2016 review, which was consistent with the division's standard approach for reviewing an efficacy supplement

	and encompassed 90 references, did not capture this publication. However, the authors' conclusion in this publication is consistent with our review of the safety data in 2016.
Kapp N et al. Best Pract Clin Obstet Gynaecol. 2020;63:37-44	Abstract. Also outside the scope of first trimester medical abortion.
Fuentes L et al. J Women's Health 2019; 28 (12): 1623, 1625 Bearak JM, Lancet Pub Health 2017 Nov;2(11): e493, e495-96 Cartwright A et al 20 J Med Internet Res 2018 20(5):e10235 Barr-Walker J, et al PLoS One 2019;14(4): e0209991 Grossman et al JAMA Network 2017;317(4):437, 437-438 Dobie S et al 31 Fam Plan Persp 1999; 31(5): 241-244 Shelton JD 8 Fam Plan Persp 1976; 8(6):260, 260-262 Norris AH et al Am J Pub Health 2020; 110 (8): 1228,1232 Upadhyay UD et al Am J Pub Health 2014; 104(9):1687, 1689	Focused on the logistics of accessing abortion care.
CDC MMWR Abortion Surveillance – United States, 2018 https://www.cdc.gov/mmwr/volumes/69/ss/ss6907a1.htm#T5 down	Contains primarily general statistics on abortion care by state.

References cited in appendix from <i>Chelius v. Becerra</i> Plaintiffs (September 29, 2021)
References included in the REMS review
None

References excluded from the REMS review	Rationale for Exclusion
Jones RK et al Guttmacher Institute Abortion Incidence and Service Availability in the United States, 2017 (2019) Guttmacher Inst, Induced Abortion in the United States (2019)	Contains primarily general statistics on abortion care and logistics of accessing abortion care.
University of Minnesota Healthy Youth Dev. Prevention Rsch Ctr, 2019 Minnesota Adolescent Sexual Health Report 3 (2019)	Not related specifically to abortion care.
Jerman J et al Guttmacher Inst, Characteristics of U.S. Abortion Patients in 2014 and Changes since 2008 (2016)	Contains figures on patient characteristics from 2008-2014.
Roberts CM et al Women's Health Issues 2014; 24:e211, e215	Focused on cost of abortion.
CDC MMWR Abortion Surveillance 2018 https://www.cdc.gov/mmwr/volumes/69/ss/ss6907a1.htm#T7 down (last updated Nov. 7, 2020)	Contains primarily statistics on number of abortions in the US.
Jones RK Persp on Sexual & Reprod Health 2017; 49:17, 20	Focused on abortion incidence and service availability.
Fuentes L et al (as above) Bearak JM et al (as above) Cartwright A et al (as above) Johns NE et al. BMC Health Serv Res 2017; 17: 287, 294	Focused on logistics of accessing abortion care.

References cited in letter from Society of Family Planning (August 11, 2021)
References included in the REMS review
Grossman D. Obstet Gynecol 2019;133 (3): 477-483

Grossman D et al. Obstet Gynecol 2021; 137 (4): 613-622.	
Winikoff B et al. Obstet Gynecol 2012; 120: 1070-1076 reviewed in 2016 clinical memo	
Chen MJ et al. Obstet Gynecol 2015;126(1):12-21 reviewed in 2016 memo	
Chong et al. Contraception 2021;104(1): 43-48	
Aiken A et al. BJOG 2021; 128 (9): 1464 -1474	
Hyland 2018 et al. Aust New Zeal J Obstet Gynaecol 2018; 58 (3): 335-340	
References excluded from the REMS review	Rationale for Exclusion
Schummers L et al. BMJ Sex Reprod Heal 2021;47(e1)	Abstract
Kapp et al. 2020 (as above)	Abstract
Upadhyay et al. 2015 (as above)	(See rationale above)
Srinivasulu et al. Contraception 2021; 104(1):92-97	Survey on clinician perspectives on access to mifepristone.
Calloway D et al. Contraception 2021; 104(1): 24-28	Primarily addresses provider stigma around abortion care.
Rasmussen et al. Contraception; 104(1): 98-103	Opinion/commentary
Cleland et al. Obstet Gynecol 2013;121(1):166-171	Published prior to March 29, 2016 - July 26, 2021 timeframe for current literature review. We note that the extensive literature search conducted as part of the 2016 clinical review, which was consistent with the division's standard approach for reviewing an efficacy supplement and encompassed 90 references, did not capture this publication. However, the authors' conclusion in this publication is consistent with our review of the safety data in 2016.
National Academy of Sciences, Engineering, and Medicine. Safety and Quality of Abortion Care in the US 2018	General information about abortion care in the US. Did not provide safety data relevant to the elements of the REMS
Raymond EG. Obstet Gynecol 2012; 119(2): 215-219	Does not separate out medical and surgical abortion.

Bartlett LA et al. Obstet Gynecol 2004; 103(4): 729-737	Focused on surgical abortion.
Jones RK, Jerman J. Time to appointment and delays in accessing care among U.S. abortion patients, Guttmacher 2016	Focused on logistics of accessing abortion care.
Foster DG et al. Perspect Sex Reprod Health 2013; 45(4):210-218	Focused on second trimester abortion.
Ely G et al. Heal Soc Work 2019;44(1):13-21	Focused on logistics of accessing abortion care.
Munro S et al. Ann Fam Med 2020; 18(5):413-421.	Survey on physician perspectives on implementing medical abortion with mifepristone.

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

(b) (6)

12/16/2021 10:22:50 AM

(b) (6)

12/16/2021 10:25:34 AM

(b) (6)

12/16/2021 10:30:08 AM

(b) (6)

12/16/2021 10:35:53 AM

(b) (6)

12/16/2021 10:49:46 AM

(b) (6)

12/16/2021 10:59:01 AM

(b) (6)

12/16/2021 11:08:07 AM

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

020687Orig1s025

OTHER REVIEW(S)

Review of Labeling

(b) (6)

NDA Number/Supplement	020687/025
Applicant	Danco Laboratories, LLC
Product Name	Mifeprex (mifepristone) tablets
Therapeutic Class	Progestin antagonist
Indication	For the medical termination of intrauterine pregnancy through 70 days gestation, in a regimen with misoprostol
Material Reviewed	Prescribing Information and Medication Guide received December 16, 2022
Date of Review	January 3, 2023
Reviewer	(b) (6)

This memorandum is the (b) (6) (Division's) review of the proposed revisions to the labeling and Medication Guide for Mifeprex submitted by Danco Laboratories (Applicant) on December 16, 2022. These revisions are to align the language in the Prescribing Information and the Medication Guide with the proposed modification to the Mifepristone single, shared system Risk Evaluation and Mitigation Strategy (REMS) (referred to as the Mifepristone REMS Program) submitted under NDA 020687/Supplement-025, as amended.

1. Background

Mifeprex (mifepristone), in a regimen with misoprostol, is indicated for the medical termination of intrauterine pregnancy through 70 days gestation. Mifeprex and its approved generic are subject to the Mifepristone REMS Program to mitigate the risk of serious complications associated with mifepristone. The Mifepristone REMS Program consists of elements to assure safe use (ETASU), an implementation system, and a timetable for submission of assessments of the REMS.

The goal of the Mifepristone REMS Program is to mitigate the risk of serious complications associated with mifepristone by:

- Requiring healthcare providers who prescribe mifepristone to be certified in the Mifepristone REMS Program (ETASU A).
- Ensuring that mifepristone is only dispensed in certain healthcare settings by or under the supervision of certified prescribers (ETASU C).
- Informing patients about the risk of serious complications associated with mifepristone (ETASU D).

On December 16, 2021, FDA sent REMS Modification Notification letters to the Applicants for Mifeprex and the approved generic version of Mifeprex, Mifepristone Tablets, 200 mg (Danco

Laboratories and GenBioPro, respectively). The letters informed the Applicants of the FDA's determination that the approved Mifepristone REMS Program must be modified to minimize the burden on the healthcare delivery system of complying with the REMS and to ensure that the benefits of the drug outweigh the risks by: (1) removal of the requirement that mifepristone be dispensed only in certain healthcare settings, specifically clinics, medical offices, and hospitals (i.e., the "in-person dispensing requirement") (ETASU C) and (2) to add the requirement for certification of pharmacies that dispense the drug (ETASU B).¹ For a detailed discussion of the modification recommendations refer to the REMS Modification Rationale Review, jointly completed by the (b) (6) and the (b) (6) on December 16, 2021.²

As proposed by the Applicants in their December 16, 2022, amendments to their pending supplements, the modified goal of the Mifepristone REMS Program would read as follows:

The goal of the REMS for mifepristone is to mitigate the risk of serious complications associated with mifepristone by:

- a) Requiring healthcare providers who prescribe mifepristone to be certified in the Mifepristone REMS Program.
- b) Ensuring that mifepristone is only dispensed by or under the supervision of certified prescribers, or by certified pharmacies on prescriptions issued by certified prescribers.
- c) Informing patients about the risk of serious complications associated with mifepristone.

The Division reviewed the Applicant's proposed revisions to the approved Mifeprex labeling under NDA 020687 Supplement 025, as amended, which provides REMS document and materials to align with the changes described in the REMS Modification Notification letters. Specifically, the requirement that mifepristone must be dispensed to patients in certain healthcare settings (ETASU C) was removed and the dispensing of mifepristone through specially certified pharmacies (ETASU B) was added. This REMS modification ensures that mifepristone is only dispensed by or under the supervision of certified prescribers, or by certified pharmacies on prescriptions issued by certified prescribers. There were no changes to ETASU A and ETASU D (prescribers must be specially certified and mifepristone must be dispensed to patients with evidence of safe use conditions, respectively).

On December 16, 2022, final labeling for the Mifeprex Prescribing Information (PI) and Medication Guide (MG) were received to align with the Mifepristone REMS Program modification. This submission is reviewed below. Review of the labeling changes made in the PI

¹ REMS Modification Notification letter dated December 16, 2021.

<https://darrrts.fda.gov/darrrts/faces/ViewDocument?documentId=090140af80633e9f>

² REMS memorandum dated December 16, 2021.

<https://darrrts.fda.gov/darrrts/faces/ViewDocument?documentId=090140af80633d74>

and in the MG for the approved generic (ANDA 091178) will be completed by the (b) (6)

2. Review of Labeling Changes

Prescribing Information and Medication Guide

The following two changes were made in both the PI and MG for Mifeprex:

1. Gender neutral edits were made throughout the PI and MG.

Reviewer comment: The Division agrees to the use of gender-neutral language to align with such changes previously approved for documents associated with this REMS.

2. Removal of instructions for patients to take the MG when they visit an emergency room or another healthcare provider who did not prescribe Mifeprex, so the provider knows that the patient is undergoing a medical abortion.

Reviewer comment: The Applicant requested this change. Removal of the instructions in the PI and MG aligns with the updated Patient Agreement Form submitted in Supplement S-025, which removes instructions for patients to take the MG to healthcare visits with providers who did not provide Mifeprex. Although these instructions were added to the MG a number of years ago, upon further consideration the Division agrees with removing them because patients seeking emergency medical care are not likely to carry a MG with them, the MG is readily available online, and information about medical conditions and previous treatments can be obtained at the point of care.

Prescribing Information

Additional changes were made in the PI to include pharmacy dispensing in Section 5.3 Mifepristone REMS Program and Section 17 Patient Counseling Information. Typographical errors were corrected in Sections 16 and 17.

1. Section 5.3 Mifepristone REMS Program

The Applicant proposed changes to one of the three bulleted ETASU requirements, to state “MIFEPREX must only be dispensed to patients by or under the supervision of a certified prescriber, or by certified pharmacies on prescriptions issued by certified prescribers.”

Reviewer comment: The revised language is consistent with the REMS modification to include certified pharmacy dispensing. The Division agrees with the proposed language.

2. Section 17 Patient Counseling Information

The Applicant proposed changes to one of the two bulleted ETASU requirements, to state “MIFEPREX is only dispensed by or under the supervision of certified prescribers or by certified pharmacies on prescriptions issued by certified prescribers.”

Reviewer comment: As above, the revised language for the PI for the approved MIFEPREX product is consistent with the REMS modification to include certified pharmacies and the Division agrees with the proposed language.

3. Section 16 How Supplied/Storage and Handling and Section 17 Patient Counseling Information

- a. Both Sections 16 and 17 refer to the “mifepristone REMS Program” which was edited to “Mifepristone REMS Program”.
- b. In Section 17, reinstatement was made of “a” before Patient Agreement Form in the bullet “Patients must sign a Patient Agreement Form”.

Reviewer comment: The Applicant will be informed of these minor edits.

Medication Guide

Additional changes were made in the MG within the section titled “How should I take Mifeprex?”, as follows:

1. Addition of obtaining Mifeprex at a pharmacy: “Mifeprex will be given to you by a healthcare provider or pharmacy.”
2. Removal of statement “Your healthcare provider will either give you or prescribe for you 4 misoprostol tablets to take 24 to 48 hours later.”

Reviewer comment: The proposed MG changes for Mifeprex are acceptable to (b) (6). The addition of Mifeprex to be dispensed at a pharmacy reflects the REMS modification change. The statement on misoprostol dispensing is removed because it is not necessary to specify in the Mifeprex labeling how misoprostol is dispensed. The directions to take misoprostol tablets 24 to 48 hours after taking mifepristone are maintained in the Mifeprex patient instructions.

3. Conclusion

We received the final proposed labeling revisions to the Prescribing Information and Medication Guide from the Applicant on December 16, 2022. The Prescribing Information and Medication Guide for Mifeprex were revised to align with the Mifepristone REMS Program modification, which include removing the requirement that Mifeprex be dispensed only in certain healthcare settings and adding the requirement for pharmacy certification. The proposed labeling revisions were reviewed and found to be acceptable with minor edits.

All labeling changes outlined above will be applied to the approved generic mifepristone (ANDA 091178). The (b) (6) will review labeling for the approved generic and ensure it mirrors the updated approved labeling for the Mifeprex product.

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

(b) (6)

01/03/2023 05:09:06 PM

(b) (6)

01/03/2023 05:10:03 PM

(b) (6)

01/03/2023 05:10:49 PM

(b) (6)

01/03/2023 05:11:23 PM

MEMORANDUM

DEPARTMENT OF HEALTH AND HUMAN SERVICES
PUBLIC HEALTH SERVICE
FOOD AND DRUG ADMINISTRATION
CENTER FOR DRUG EVALUATION AND RESEARCH

TO: FILE

FROM:

(b) (6)

(b) (6)

Digitally signed by (b) (6)
Date: 2022.12.23 13:11:43 -05'00'

DATE: December 23, 2022

SUBJECT: Review of Supplemental Drug Applications Proposing Modifications to the
Mifepristone REMS Program

FDA is currently reviewing a supplemental new drug application from Danco Laboratories, LLC (Danco) and a supplemental abbreviated new drug application from GenBioPro, Inc. (GBP) that propose to modify the Mifepristone Risk Evaluation and Mitigation Strategy (REMS) Program as approved under Danco's new drug application for Mifeprex (mifepristone) (NDA 020867) and GBP's abbreviated new drug application for Mifepristone Tablets 200 mg (ANDA 091178). Citing the Comstock Act, 18 U.S.C. §§ 1461, 1462, Plaintiffs in *Alliance for Hippocratic Medicine v. U.S. Food and Drug Administration*, No. 2:22-cv-00223-Z (N.D. Tex.), have alleged that FDA's actions regarding mifepristone do not comply with "federal laws that expressly prohibit the mailing or delivery by any letter carrier, express company, or other common carrier of any substance or drug intended for producing abortion" and also that FDA "failed to acknowledge and address" those laws. Complaint ¶¶ 22, 392 (Nov. 18, 2022). This memorandum notes that the Office of Legal Counsel of the United States Department of Justice, which provides controlling advice to Executive Branch officials on questions of law, has concluded that the Comstock Act provisions cited by Plaintiffs "[do] not prohibit the mailing of mifepristone or misoprostol where the sender lacks the intent that the recipient will use them unlawfully. And in light of the many lawful uses of mifepristone and misoprostol, the fact that these drugs are being mailed to a jurisdiction that significantly restricts abortion is not a sufficient basis for concluding that the mailing violates [these provisions]." Memorandum for Thomas J. Marshall, General Counsel, United States Postal Service, from Christopher H. Schroeder, Assistant Attorney General, Office of Legal Counsel, *Re: Application of the Comstock Act to the Mailing of Prescription Drugs That Can Be Used for Abortions*, at 15 (December 23, 2022).¹ Thus, even if the Comstock Act provisions bear on FDA's analysis of the pending supplemental drug applications, in light of the conclusions set forth by the Office of Legal Counsel, they pose no issue for FDA's approval of the applications.

¹ The Office of Legal Counsel's analysis applies to 18 U.S.C. § 1461 and § 1462. *See id.* at 1 n.3.

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

(b) (6)

12/23/2022 03:02:29 PM

I am placing this document in for

(b) (6)

This is for archiving purposes.

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

020687Orig1s025

**RISK ASSESSMENT and RISK MITIGATION
REVIEW(S)**

Risk Evaluation and Mitigation Strategy (REMS) Memorandum
REMS Modification: Removal of a Requirement and Addition of a Requirement
U.S. FOOD AND DRUG ADMINISTRATION
CENTER FOR DRUG EVALUATION AND RESEARCH

(b) (6)

(b) (6)

NDA/ANDA #s: NDA 20687, ANDA 91178
Products: Mifeprex, mifepristone, 200 mg tablets
APPLICANTS: Danco, GenBioPro
FROM: (b) (6)
(b) (6)
DATE: December 16, 2021

Mifepristone is a progestin antagonist indicated, in a regimen with misoprostol, for the medical termination of intrauterine pregnancy (IUP) through 70 days gestation. Mifeprex was approved on September 28, 2000, with a restricted distribution program under 21 CFR 314.520 (subpart H) and subsequently was deemed to have a REMS under section 505-1 of the Federal Food, Drug, and Cosmetic Act with the passage of the Food and Drug Administration Amendments Act (FDAAA) of 2007. The Mifeprex REMS with elements to assure safe use (ETASU) was approved on June 8, 2011 and a supplemental efficacy application and REMS modification was approved on March 29, 2016. The Mifepristone REMS Program (a single, shared system REMS that currently includes NDA 020687 and ANDA 91178) was approved on April 11, 2019.

The goal of the REMS for mifepristone is to mitigate the risk of serious complications associated with mifepristone. The current Mifepristone REMS Program includes elements to assure safe use (ETASU), an implementation system, and a timetable for submission of assessments. Elements to assure safe use include requirements for prescriber certification (ETASU A), that mifepristone be dispensed only in certain healthcare settings by or under the supervision of a certified prescriber (ETASU C), and that mifepristone be dispensed only with documentation of safe use conditions (ETASU D). Documentation of safe use conditions consists of a Patient Agreement Form between the prescriber and the patient, indicating that the patient has received counseling from the prescriber regarding the risk of serious complications associated with mifepristone.

The requirement under ETASU C that mifepristone be dispensed only in certain health care settings, specifically clinics, medical offices, and hospitals, is referred to as the “in-person dispensing requirement.”

On January 31, 2020 the Secretary of Health and Human Services (HHS) declared COVID-19 a public health emergency (PHE) as of January 27, 2020. During the COVID-19 pandemic, there have been periods when the in-person dispensing requirement has not been enforced. From July

13, 2020 until January 12, 2021, enforcement was barred by an injunction issued in the *ACOG v. FDA* litigation. More recently, on April 12, 2021, the Agency stated its intent to exercise enforcement discretion with respect to the in-person dispensing requirement during the COVID-19 PHE, which is still ongoing as of the date of this review. These circumstances have provided additional information regarding the in-person dispensing requirement as there have been periods when the in-person dispensing requirement was not enforced.

As part of the May 7, 2021, joint motion to stay the *Chelius v. Becerra* litigation, the Agency agreed to undertake a full review the Mifepristone REMS Program, in accordance with the REMS assessment provisions of the Federal Food, Drug, and Cosmetic Act (FD&C Act).¹

After consultations between the (b) (6), analyzing several different sources of information, including published literature, safety information collected during the COVID-19 PHE, FDA Adverse Event Reporting System (FAERS) reports, REMS assessment reports, and information provided by advocacy groups, individuals, the Applicants, and the plaintiffs in the *Chelius* litigation, we determined that the approved REMS for mifepristone could be modified without adversely impacting patient safety. Importantly, our review did not identify any differences in adverse events between periods when the in-person dispensing requirement was being enforced and periods when that requirement was not being enforced. The data suggested that the requirements for prescriber certification (ETASU A) and the Patient Agreement Form (ETASU D) should be maintained, while the in-person dispensing requirement (under ETASU C) could be removed, to reduce the burden imposed by the REMS. In determining that the in-person dispensing requirement could be removed, we concluded that a new requirement for pharmacy certification (ETASU B) is necessary to ensure the benefits of the product outweigh the risks.

(b) (6) and (b) (6) assessment and recommendations were (b) (6) (b) (6) on November 2, 2021. The (b) (6) unanimously agreed with our recommendations.

For more detailed information on the review and assessments of the information, refer to the REMS Modification Rationale Review, jointly completed by (b) (6) and (b) (6) on December 16, 2021.

In conclusion, provided all other conditions of the Mifepristone REMS Program are met and that the other elements of the REMS are maintained (ETASU A and D), the following are required:

1. Modification of the Mifepristone REMS Program to remove the requirement under ETASU C that mifepristone be dispensed only in certain healthcare settings, specifically clinics, medical offices, and hospitals. This would allow, for example, dispensing of mifepristone by mail, via certified prescribers or pharmacies, in addition to in-person

¹ Section 505-1(g)(2) of the FD&C Act (21 U.S.C. § 355-1(g)(2)).

dispensing in clinics, medical offices and hospitals as currently outlined in ETASU C . We find that this provision is no longer necessary to ensure that the benefits of the drug outweigh the risks and that removing it will help minimize the burden of complying with the REMS on the healthcare delivery system.

2. Modification of the Mifepristone REMS Program to add a requirement under ETASU B that pharmacies that dispense the drug be specially certified.

Based on the (b) (6) and (b) (6) determination that a modified REMS with the components described above is necessary to reduce the burden imposed by the REMS and ensure the benefits of mifepristone outweigh the risks, FDA is requiring submission of the proposed REMS modification within 120 days of the date of the notification letter.

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

(b) (6)

12/16/2021 11:49:15 AM

(b) (6) (b) (6)
(b) (6) (b) (6)
(b) (6) (b) (6)
Center for Drug Evaluation and Research (CDER)

Application Type	NDA and ANDA
Application Number	NDA 020687 and ANDA 91178
Supplement Number, Date Received	NDA Supplement-025 and ANDA Supplement-004 received June 22, 2022 (sequence 18 and 87 respectively) and amended October 19, 2022 (sequences 22 and 91 respectively), November 30, 2022, and December 9, 2022 (sequences 25 and 93 respectively)
Targeted Action Date	December 19, 2022
(b) (6) #	2022-1169
Reviewer Names	(b) (6)
(b) (6)	(b) (6)
(b) (6)	(b) (6)
Review Completion Date	December 14, 2022
Subject	Review of proposed Major REMS Modification
Established Name	Mifepristone REMS Program
Name of Applicant	Danco Laboratories, LLC and GenBioPro, Inc.
Therapeutic Class	Progestin antagonist
Formulation	Oral tablet

1. Introduction

Refer to the proposed modification to the single, shared system Risk Evaluation and Mitigation Strategy (REMS) for mifepristone 200 mg (hereafter referred to as the Mifepristone REMS Program) submitted by Danco Laboratories, LLC (Danco) for new drug application (NDA) 020687 and by GenBioPro, Inc. (GBP) for abbreviated new drug application (ANDA) 091178.

The Applicants submitted proposed modifications to the Mifepristone REMS Program on June 22, 2022 in response to REMS Modification Notification letters issued on December 16, 2021 to Danco and GBP, requiring the following modifications to minimize the burden on the healthcare delivery system of complying with the REMS and to ensure that the benefits of the drug outweigh the risks:

- removal of the requirement that mifepristone be dispensed only in certain healthcare settings, specifically clinics, medical offices, and hospitals (i.e., the “in-person dispensing requirement”)
- addition of certification of pharmacies that dispense the drug

The comments in this review focus on the Applicants’ amendments that were received on December 9, 2022, which included updates to the REMS Document and materials that were discussed and edited during a teleconference between the Agency and Applicants on December 8, 2022.

2. Comments to the Sponsor

General Comments

The documents have been revised for you per the comments below.

REMS Document

We have revisions in the REMS document in the following sections:

II. A.2.a.iv.2) I, the language must be updated to replace “provider” with “prescriber,” and to remove the phrase that (b) (4)

(b) (4)

II.A.2.c.ii A requirement was added to ensure prescribers previously certified in the Mifepristone REMS Program complete the new *Prescriber Agreement Form* (b) (4)

(b) (4)

II. A.3, we have the following edits to align with the statute. Mifepristone must be (b) (4) dispensed (b) (4) to patients with evidence or other documentation of safe use conditions as ensured by the certified prescriber in signing the *Prescriber Agreement Form*.

II.B.5 must be updated to include: Mifepristone Sponsors must audit their certified pharmacies within 180 calendar days after the pharmacy places its first order of mifepristone, and annually thereafter audit certified pharmacies that have ordered mifepristone in the previous 12 months to ensure that all processes and procedures are in place and functioning to support the requirements of the Mifepristone REMS Program. Mifepristone Sponsors will take steps to address their pharmacy compliance if noncompliance is identified.

REMS Supporting Document

We require clarification and revisions in several sections. A red-lined Supporting Document has been attached for reference.

The wording in the Supporting Document was not consistent with the REMS document and REMS materials. Throughout the Supporting Document, the document was revised to align with the wording in the REMS Document, *Prescriber Agreement Form*, *Patient Agreement Form*, and *Pharmacy Agreement Form*.

The Supporting Document is inconsistent with respect to grammar and punctuation, as well as with respect to capitalization, italicization, and terminology. Review and revise the Supporting Document for grammar and punctuation, and for consistency in capitalization, italicization, and terminology.

The term “(b) (4)” was replaced with “supervised healthcare providers” where applicable.

There was minimal information about account set up processes for the certified prescribers and pharmacies. Include explanation of how accounts will be set up and maintained and what information from the account set up process will be used for REMS assessments.

There was also minimal information about how Sponsors will communicate with each other when stakeholders report information that the other needs to know such as when stakeholders report patient deaths to the wrong Sponsor or when decertifications of pharmacies or prescribers take place. Clarify what actions require timely communication between Sponsors and the timeline in which these actions should be reported to the other Sponsor.

Prescriber Agreement Forms for Danco Laboratories, LLC and GenBioPro, Inc.

(b) (4). This data will be available through the account set up process.

Pharmacy Agreement Forms for Danco Laboratories, LLC and GenBioPro, Inc.

(b) (4) must be removed as it is redundant to the language later in the form. The confidentiality requirement later in the form, under the requirements that certified pharmacies must put processes and procedures in place to accomplish, must be further aligned with the confidentiality requirement added to the REMS Document (discussed above).

Patient Agreement Form

The Patient Agreement Form is acceptable.

Resubmission Instructions

Accept all track changes and submit the following revised REMS materials by 12/16/22. The next submission to the Gateway should include Clean Word, Tracked Word, and pdf formatted versions of the following documents and one clean compiled PDF file that includes the REMS Document and all REMS materials in their final format:

- REMS Document
- Prescriber Agreement Form for Danco Laboratories, LLC
- Prescriber Agreement Form for GenBioPro, Inc.
- Pharmacy Agreement Form for Danco Laboratories, LLC
- Pharmacy Agreement Form for GenBioPro, Inc.
- Patient Agreement Form
- REMS Supporting Document with the Assessment Plan

Appendix

- REMS Document
- Prescriber Agreement Form for Danco Laboratories, LLC
- Prescriber Agreement Form for GenBioPro, Inc.

Pharmacy Agreement Form for Danco Laboratories, LLC
Pharmacy Agreement Form for GenBioPro, Inc
REMS Supporting Document

30 Pages of Draft REMS Documents Have Been Withheld in Full as B4 (CCI/TS)
Immediately Following This Page

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

(b) (6)

12/14/2022 04:18:28 PM

(b) (6)

12/14/2022 04:22:00 PM

(b) (6)

12/14/2022 04:34:44 PM

(b) (6)
(b) (6)
(b) (6)

Center for Drug Evaluation and Research (CDER)

Application Type	NDA and ANDA
Application Number	NDA 020687 and ANDA 91178
Supplement Number, Date Received	NDA Supplement-025 and ANDA Supplement-004 received June 22, 2022 (sequence 18 and 87 respectively) and amended October 19, 2022 (sequences 22 and 91 respectively), November 30, 2022
(b) (6) #	2022-1169
(b) (6)	(b) (6)
(b) (6)	(b) (6)
Review Completion Date	December 8, 2022
Subject	Review of Proposed Major REMS Modification: Draft REMS Assessment Plan Comments
Established Name	Mifepristone REMS Program
Name of Applicant	Danco Laboratories, LLC and GenBioPro, Inc.
Therapeutic Class	Progestin antagonist
Formulation	Oral tablet

1. Introduction

Refer to the proposed modification to the single, shared system Risk Evaluation and Mitigation Strategy (REMS) for mifepristone 200 mg (hereafter referred to as the Mifepristone REMS Program) submitted by Danco Laboratories, LLC (Danco) for new drug application (NDA) 020687 and from GenBioPro Inc. (GBP) for abbreviated new drug application (ANDA) 091178.

The Applicants submitted proposed modifications to the Mifepristone REMS Program on June 22, 2022 in response to a REMS Modification Notification letter issued on December 16, 2021 to Danco and GBP, requiring the following modifications to minimize the burden on the healthcare delivery system of complying with the REMS and to ensure that the benefits of the drug outweigh the risks:

- removal of the requirement that mifepristone be dispensed only in certain healthcare settings, specifically clinics, medical offices, and hospitals (i.e., the “in-person dispensing requirement”)
- addition of certification of pharmacies that dispense the drug

The comments in this review focus on revisions to the REMS Assessment Plan.

2. Comments to the Applicant

We have provided a draft REMS Assessment Plan as discussed on the December 7, 2022 teleconference. The proposed Agency edits have been marked in tracked changes from the Mifepristone REMS Assessment Plan in the April 11, 2019, Supplement Approval Letter. Refer to the attached Draft REMS Assessment Plan with tracked changes. Review of the REMS proposal is ongoing; these comments should not be considered final.

We have the following comments:

1. Provide each assessment plan metric for the two previous, current, and cumulative reporting periods (if applicable) for both the NDA and ANDA unless otherwise noted.
2. The Assessment Plan Categories of 1) Program Implementation and Operations and 2) Overall Assessment of REMS Effectiveness were added.
3. A REMS Certification Statistics metric was added to capture the following:
 - a. Total number of certified, newly certified, and active prescribers along with a summary of the practice setting of the certified prescribers and the method in which they became certified.
 - b. Total number of certified, newly certified, and active certified pharmacies.
 - c. Total number of authorized, newly authorized, and active wholesaler/distributors.
4. A Utilization Data metric was added to capture wholesaler/distributor shipment and pharmacy data.
5. A REMS Compliance Data metric was added to capture stakeholder audit results and a summary of instances of non-compliance and actions taken to address non-compliance.

Resubmission Instructions

Submit the revised REMS Assessment Plan in the REMS Supporting Document with your 12/9/22 submission. Accept the tracked changes in the draft REMS Assessment Plan with which you agree and only indicate any new changes you propose as tracked changes in your next submission. The submission should include clean Word, tracked changes Word, and pdf formatted versions of the following document.

Appendix

REMS Assessment Plan

4 Pages of Draft REMS Documents Have Been Withheld in Full as B4 (CCI/TS)
Immediately Following This Page

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

(b) (6)

12/08/2022 03:27:22 PM

(b) (6)

12/08/2022 03:29:06 PM

(b) (6) (b) (6)
(b) (6) (b) (6)
(b) (6) (b) (6)
Center for Drug Evaluation and Research (CDER)

Application Type	NDA and ANDA
Application Number	NDA 020687 and ANDA 91178
Supplement Number, Date Received	NDA Supplement-025 and ANDA Supplement-004 received June 22, 2022 (sequence 18 and 87 respectively) and amended October 19, 2022 (sequences 22 and 91 respectively), November 30, 2022
PDUFA Date	December 19, 2022
(b) (6) #	2022-1169
Reviewer Names	(b) (6)
(b) (6)	(b) (6)
(b) (6)	(b) (6)
(b) (6)	(b) (6)
Review Completion Date	December 5, 2022
Subject	Review of proposed Major REMS Modification
Established Name	Mifepristone REMS Program
Name of Applicant	Danco Laboratories, LLC and GenBioPro, Inc.
Therapeutic Class	Progestin antagonist
Formulation	Oral tablet

1. Introduction

Refer to the proposed modification to the single, shared system Risk Evaluation and Mitigation Strategy (REMS) for mifepristone 200 mg (hereafter referred to as the Mifepristone REMS Program) submitted by Danco Laboratories, LLC (Danco) for new drug application (NDA) 020687 and by GenBioPro, Inc. (GBP) for abbreviated new drug application (ANDA) 091178.

The Applicants submitted proposed modifications to the Mifepristone REMS Program on June 22, 2022 in response to REMS Modification Notification letters issued on December 16, 2021 to Danco and GBP, requiring the following modifications to minimize the burden on the healthcare delivery system of complying with the REMS and to ensure that the benefits of the drug outweigh the risks:

- removal of the requirement that mifepristone be dispensed only in certain healthcare settings, specifically clinics, medical offices, and hospitals (i.e., the “in-person dispensing requirement”)
- addition of certification of pharmacies that dispense the drug

The comments in this review focus on the Applicants’ amendments that were received on October 19 and 20, 2022 and November 30, 2022 (REMS Document). The comments also reflect the REMS Document that was further discussed and edited during a teleconference between the Agency and Applicants on December 1, 2022.

2. Comments to the Sponsor

General Comments

We have updated the REMS Document as discussed during the December 1, 2022 teleconference and aligned the REMS materials. Your edits, where the Agency agrees, have been accepted and Agency edits have been marked in tracked changes. Refer to the attached, red-lined REMS Document and REMS Materials. Review of the REMS proposal is ongoing; these comments should not be considered final.

- The Agency has determined that further clarification that certified prescribers are responsible for overseeing implementation and compliance with the REMS Program is appropriate in the *Prescriber Agreement Form* and has been added. This clarification provides flexibility for certified prescribers in overseeing REMS implementation and compliance. The certified prescriber may do so in a manner that may include the use of a larger healthcare team and delegation of tasks. We have determined that to make additional changes to the prescriber certification requirements would constitute a substantive modification to the REMS that would go beyond the required REMS modifications set forth in the December 16, 2021 REMS Modification Notification letters. Adequate rationale is required to support a proposed REMS modification for the addition, modification, or removal of any goal or element of the REMS, as described in section 505-1(g)(4) of the FD&C Act. Your application does not include such adequate rationale.
- We agree that healthcare settings, such as medical offices, clinics, and hospitals, where mifepristone is dispensed by or under the supervision of a certified prescriber in the Mifepristone REMS Program do not require pharmacy certification and have added this clarification to the *Prescriber Agreement Forms* and *Pharmacy Agreement Forms*. These clarifications negate the need for further clarification in the REMS Document as you proposed in the footnote to section II.A.2.
- Moving forward, italicize all proper names of forms e.g. *Prescriber Agreement Form* in the REMS.

Prescriber Agreement Forms for Danco Laboratories, LLC and GenBioPro, Inc.

The instructions on the *Prescriber Agreement Forms* were revised for clarity. Prescriber information collected on the forms was updated to capture practice setting (b) (4) address, and an additional line for medical license state was added. Duplicative or unnecessary instructions have been removed.

Pharmacy Agreement Forms for Danco Laboratories, LLC and GenBioPro, Inc.

The titles of the *Pharmacy Agreement Forms* were edited to remove “(b) (4).” The REMS pharmacy requirements will dictate whether a pharmacy can participate in the Mifepristone REMS Program and therefore (b) (4). Additional information fields regarding authorized representatives and pharmacies were added and the term “(b) (4)” was replaced with “pharmacy.” Duplicative or unnecessary instructions have been removed.

Patient Agreement Form

The first paragraph of the form has been aligned with the currently approved Mifepristone REMS Program and clarification that signatures on the document can be written or electronic was added. Risk information regarding ectopic pregnancy has been reorganized. (b) (4)

Resubmission Instructions

Accept all track changes and submit the following revised REMS materials by 12/08/22. The next submission to the Gateway should include Clean Word, Tracked Word, and pdf formatted versions of the following documents and one clean compiled PDF file that includes the REMS Document and all REMS materials in their final format:

- REMS Document
- Prescriber Agreement Form for Danco Laboratories, LLC
- Prescriber Agreement Form for GenBioPro, Inc.
- Pharmacy Agreement Form for Danco Laboratories, LLC
- Pharmacy Agreement Form for GenBioPro, Inc
- Patient Agreement Form
- REMS Supporting Document with the Assessment Plan
- Updated prescription label and Medication Guide

Appendix

- REMS Document
- Prescriber Agreement Form for Danco Laboratories, LLC
- Prescriber Agreement Form for GenBioPro, Inc.
- Pharmacy Agreement Form for Danco Laboratories, LLC
- Pharmacy Agreement Form for GenBioPro, Inc
- Patient Agreement Form

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

(b) (6)

12/05/2022 08:49:01 PM

(b) (6)

12/05/2022 08:53:19 PM

(b) (6)

12/05/2022 09:08:48 PM

(b) (6)

12/05/2022 09:17:13 PM

(b) (6) (b) (6)
(b) (6) (b) (6)
(b) (6) (b) (6)
Center for Drug Evaluation and Research (CDER)

Application Type	NDA and ANDA
Application Number	NDA 020687 and ANDA 91178
Supplement Number, Date Received	NDA Supplement-025 and ANDA Supplement-004 received June 22, 2022 (sequence 18 and 87 respectively) and amended October 19, 2022 (sequences 22 and 91 respectively)
Action Date	December 19, 2022
(b) (6) #	2022-1169
Reviewer Names	(b) (6)
(b) (6)	(b) (6)
(b) (6)	(b) (6)
	(b) (6)
Review Completion Date	November 23, 2022
Subject	Review of proposed Major REMS Modification
Established Name	Mifepristone REMS Program
Name of Applicant	Danco Laboratories, LLC and GenBioPro, Inc.
Therapeutic Class	Progestin antagonist
Formulation	Oral tablet

1. Introduction

Refer to the proposed modification to the single, shared system Risk Evaluation and Mitigation Strategy (REMS) for mifepristone 200 mg (hereafter referred to as the Mifepristone REMS Program) submitted by Danco Laboratories, LLC (Danco) for new drug application (NDA) 020687 and from GenBioPro Inc. (GBP) for abbreviated new drug application (ANDA) 091178.

The Applicants submitted proposed modifications to the Mifepristone REMS Program on June 22, 2022 in response to a REMS Modification Notification letter issued on December 16, 2021 to Danco and GBP, requiring the following modifications to minimize the burden on the healthcare delivery system of complying with the REMS and to ensure that the benefits of the drug outweigh the risks:

- removal of the requirement that mifepristone be dispensed only in certain healthcare settings, specifically clinics, medical offices, and hospitals (i.e., the “in-person dispensing requirement”)
- addition of certification of pharmacies that dispense the drug

Danco amended their submission on October 19, 2022 and GBP amended their submission on October 20, 2022.

2. Comments to the Sponsor

General Comments

For clarity, we have used the approved REMS Document to provide edits. Your proposed edits and Agency edits have been marked in tracked changes. Refer to the attached, red-lined REMS Document. Review of the REMS proposal is ongoing; these comments should not be considered final.

We have additional questions that must be addressed. Refer to the red-lined REMS Document attached. You must address the following question with your next submission:

1. Refer to the distributor requirement on page five (in all markup), “Put processes and procedures in place to maintain a distribution system that is secure, confidential and follows all processes and procedures, including those for storage, handling, shipping, tracking package serial numbers, proof of delivery and controlled returns of mifepristone” (underline added for emphasis)

Clarify whether the packages that are tracked by the distributors will still be done by serial numbers or if tracking will only use the NDC and lot numbers.

REMS Goal

We have revised the goal to the following:

“The goal of the REMS for mifepristone is to mitigate the risk of serious complications associated with mifepristone by:

- a) Requiring healthcare providers who prescribe mifepristone to be certified in the Mifepristone REMS Program.
- b) Ensuring that mifepristone is only dispensed by or under the supervision of certified prescribers or by certified pharmacies.
- c) Informing patients about the risk of serious complications associated with mifepristone.”

REMS Document

We have provided edits to the REMS Document to reflect the requirements communicated in the December 16, 2021 letter and to incorporate additional requirements. We have determined that for this modification we will not be using the format for the REMS that is in the 2017 draft Guidance for Industry - Format and Content of a REMS Document, as not all requirements are easily transferred to the newer format and may result in creating unnecessary confusion to stakeholders. Note that all REMS Materials and the REMS Supporting Document must align with the REMS Document.

Prescriber requirements were edited for clarity, brevity and to align with certified prescriber qualifications in the currently approved REMS. Additional requirements were needed to address situations that may arise if a certified prescriber opts to dispense through a certified pharmacy.

Pharmacy requirements were revised: to ensure pharmacies are able to ship mifepristone using a shipping service that provides tracking information, to include the use of an authorized representative to coordinate REMS implementation on behalf of the pharmacy, to dispense mifepristone such that it is delivered to the patient within four calendar days of the date the pharmacy receives the prescription, and to confirm with the prescriber and document the appropriateness of dispensing mifepristone for patients who will not receive the drug within four calendar days of the date the pharmacy receives the prescription. Additional record-keeping and reporting requirements were also added.

Additional Sponsor requirements were added to support prescriber, pharmacy, and distributor REMS stakeholder requirements, and to ensure the REMS operates as intended.

REMS Supporting Document

The REMS Supporting Document must be included in your next submission and is necessary to help us understand how these changes will be implemented before we can take an action.

Resubmission Instructions

Submit the following revised REMS materials by 11/30/22. Accept the track changes in the REMS Document with which you agree in the Word newly redlined documents and only indicate any new changes you propose as redlined changes in your next submission. The next submission to the Gateway should include Clean Word, Tracked Word, and pdf formatted versions of the following documents:

- REMS Document
- REMS Supporting Document

Appendix

REMS Document

5 Pages of Draft REMS Documents Have Been Withheld in Full as B4 (CCI/TS)
Immediately Following This Page

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

(b) (6)
11/23/2022 11:42:12 AM

(b) (6)
11/23/2022 11:48:25 AM

(b) (6)
11/23/2022 11:51:57 AM

(b) (6)
11/23/2022 11:53:43 AM

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

020687Orig1s025

ADMINISTRATIVE and CORRESPONDENCE
DOCUMENTS



NDA 020687

REMS MODIFICATION NOTIFICATION

Danco Laboratories, LLC

(b) (4), (b) (6)

P.O. Box 4816
New York, NY 10185

Dear (b) (4), (b) (6) :

We refer to your new drug application (NDA) submitted under section 505(b) of the Federal Food, Drug, and Cosmetic Act (FDCA) for Mifeprex (mifepristone) Tablets.

RISK EVALUATION AND MITIGATION STRATEGY (REMS) REQUIREMENTS

The REMS for mifepristone was originally approved on June 8, 2011, and your single shared system REMS (SSS REMS) was approved on April 11, 2019. Your last SSS REMS modification was approved May 14, 2021. The SSS REMS consists of elements to assure safe use, an implementation system, and a timetable for submission of assessments of the REMS.

In accordance with section 505-1(g)(4)(B) of the Federal Food, Drug, and Cosmetic Act (FDCA), we have determined that your approved REMS for mifepristone must be modified to minimize the burden on the healthcare delivery system of complying with the REMS and to ensure that the benefits of the drug outweigh the risks.

This determination is based on a review of published literature, safety information collected during the COVID-19 PHE, FDA Adverse Event Reporting System (FAERS) reports, REMS assessment reports, and information provided by advocacy groups, individuals, the Applicants, and plaintiffs in ongoing litigation.

Your approved REMS must be modified as follows:

Elements to Assure Safe Use: We have determined that the requirement that mifepristone be dispensed only in certain healthcare settings, specifically clinics, medical offices, and hospitals (i.e., the “in-person dispensing requirement”) is no longer necessary to ensure the benefits of mifepristone outweigh the risks of serious complications associated with mifepristone that are listed in the labeling of the drug. Removal of the requirement for in-person dispensing will also minimize the burden on the healthcare delivery system of complying with the REMS.

Elements to Assure Safe Use: Pursuant to 505-1(f)(1), we have also determined that an additional element to assure safe use is necessary to mitigate the risk of serious

complications associated with mifepristone listed in the labeling of the drug. Modification of the Mifepristone REMS to allow dispensing of mifepristone by pharmacies requires the addition of certification of pharmacies that dispense the drug.

Your REMS must include elements to mitigate this risk, including at least the following:

- Healthcare providers have particular experience or training, or are specially certified
- Pharmacies, practitioners, or health care settings that dispense the drug are specially certified
- The drug is dispensed to patients with evidence or other documentation of safe use conditions.

The REMS must include an implementation system to monitor, evaluate, and work to improve the implementation of the elements to assure safe use (outlined above). Include an intervention plan to address any findings of non-compliance with the ETASU.

The proposed REMS must include a timetable for submission of assessments. The proposed REMS modification submission should include a new proposed REMS document and appended REMS materials, as appropriate, that show the complete previously approved REMS with all proposed modifications highlighted and revised REMS materials.

In addition, the submission should also include an update to the REMS supporting document that includes a description of all proposed modifications and their potential impact on other REMS elements. Revisions to the REMS supporting document should be submitted with all changes marked and highlighted.

Because we have determined that a REMS modification as described above is necessary to minimize the burden on the health care delivery system of complying with the REMS, and to ensure that the benefits of the drug outweigh the risks, you must submit a proposed REMS modification within 120 days of the date of this letter.

Submit the proposed modified REMS as a Prior Approval supplement (PAS) to your NDA.

Because FDA is requiring the REMS modifications in accordance with section 505-1(g)(4)(B), you are not required to submit an adequate rationale to support the proposed modifications, as long as the proposals are consistent with the modifications described in this letter. If the proposed REMS modification supplement includes changes that differ from the modifications described in this letter, an adequate rationale is required for those additional proposed changes in accordance with section 505-1(g)(4)(A).

Prominently identify the submission with the following wording in bold capital letters at the top of the first page of the submission:

**NEW SUPPLEMENT FOR NDA 020687/S-000
PRIOR APPROVAL SUPPLEMENT
PROPOSED MAJOR REMS MODIFICATION**

Prominently identify subsequent submissions related to the proposed REMS modification with the following wording in bold capital letters at the top of the first page of the submission:

**NDA 020687/S-000
PROPOSED REMS MODIFICATION-AMENDMENT**

To facilitate review of your submission, we request that you submit your proposed modified REMS and other REMS-related materials in Microsoft Word format. If certain documents, such as enrollment forms, are only in PDF format, they may be submitted as such, but the preference is to include as many as possible in Word format.

SUBMISSION OF REMS DOCUMENT IN SPL FORMAT

In addition to submitting the proposed modified REMS as described above, you can also submit the REMS document in Structured Product Labeling (SPL) format. If you intend to submit the REMS document in SPL format, include the SPL file with your proposed REMS modification submission.

For more information on submitting REMS in SPL format, please email FDAREMSwebsite@fda.hhs.gov.

If you have any questions, call (b) (6), at (b) (6).

Sincerely,

{See appended electronic signature page}

(b) (6)

Center for Drug Evaluation and Research

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

(b) (6)

12/16/2021 03:09:07 PM