

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

210274Orig1s000

LABELING

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use **VANCOMYCIN HYDROCHLORIDE FOR INJECTION** safely and effectively. See full prescribing information for **VANCOMYCIN HYDROCHLORIDE FOR INJECTION**.

VANCOMYCIN HYDROCHLORIDE for injection, for intravenous use or oral use.

Initial U.S. Approval: 1958

INDICATIONS AND USAGE

Vancomycin Hydrochloride for Injection is a glycopeptide antibacterial indicated in adult and pediatric patients less than 18 years of age as follows:

- Vancomycin Hydrochloride for Injection **administered intravenously** is indicated for the treatment of:
 - Septicemia (1.1)
 - Infective Endocarditis (1.2)
 - Skin and Skin Structure Infections (1.3)
 - Bone Infections (1.4)
 - Lower Respiratory Tract Infections (1.5)
- Vancomycin Hydrochloride for Injection **administered orally** is indicated for the treatment of:
 - *Clostridioides difficile*-associated diarrhea (1.6)
 - Enterocolitis caused by *Staphylococcus aureus* (including methicillin-resistant strains) (1.7)

Limitations of Use (1.8)

- Vancomycin Hydrochloride for Injection **administered intravenously** is not approved for the treatment of *C. difficile*-associated diarrhea and enterocolitis caused by susceptible isolates of *Staphylococcus aureus* because it is not effective.
- Vancomycin Hydrochloride for Injection **administered orally** is not approved for the treatment of septicemia, infective endocarditis, skin and skin structure infections, bone infections and lower respiratory tract infections because it is not effective.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Vancomycin Hydrochloride for Injection and other antibacterial drugs, Vancomycin Hydrochloride for Injection should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. (1.9)

DOSAGE AND ADMINISTRATION

- **Pharmacy Bulk Package. Not for direct intravenous infusion**
See full prescribing information for important information on intravenous and oral administration, preparation and storage instructions (2.1, 2.5, 2.8).
- **Recommended Dosage and Administration**
- **Intravenous Use:**
 - Administer Vancomycin Hydrochloride for Injection in a diluted solution by intravenous infusion over 60 minutes or greater to reduce the risk of infusion reactions.
 - Adult Patients: 2 grams (g) divided either as 0.5 g every 6 hours or 1 g every 12 hours (2.2)
 - Pediatric Patients (1 month and older): 10 mg/kg per dose given every 6 hours (2.2)
 - Pediatric Patients (younger than 1 month of age): See full prescribing information for recommended dosage (2.2)
 - Patients with Renal Impairment: See full prescribing information for recommended dosage in patients with renal impairment (2.4)
- **Oral use:**
Adult Patients (18 years of age or greater):
 - *C. difficile* Associated Diarrhea: 125 mg administered orally 4 times daily for 10 days (2.3).
 - Staphylococcal enterocolitis: 500 mg to 2 grams administered orally in 3 or 4 divided doses for 7 to 10 days (2.3)
Pediatric Patients (less than 18 years of age):
 - For both *C. difficile*-associated diarrhea and staphylococcal enterocolitis: 40 mg/kg in 3 or 4 divided doses for 7 days to 10 days. The total daily dosage should not exceed 2 g (2.3)

DOSAGE FORMS AND STRENGTHS

Vancomycin Hydrochloride for Injection: Pharmacy Bulk Package vial containing vancomycin hydrochloride equivalent to 5 gram or 10 gram of vancomycin base (3)

CONTRAINDICATIONS

Hypersensitivity to vancomycin, and polysorbate 80 (4)

WARNINGS AND PRECAUTIONS

- **Nephrotoxicity:** Systemic vancomycin exposure may result in acute kidney injury (AKI) including acute renal failure, mainly due to interstitial nephritis or less commonly acute tubular necrosis. Monitor serum vancomycin concentrations and renal function in patients receiving Vancomycin Hydrochloride for Injection intravenously. Monitor renal function in patients over 65 years of age receiving Vancomycin Hydrochloride for Injection orally. (5.1)
- **Ototoxicity:** Ototoxicity has occurred in patients administered vancomycin intravenously or orally. Monitor for signs and symptoms of ototoxicity during oral or intravenous therapy. Assessment of auditory function may be appropriate in some instances. (5.2)
- **Severe Dermatologic Reactions:** Discontinue Vancomycin Hydrochloride for Injection at the first appearance of skin rashes, mucosal lesions, or blisters (5.3).
- **Neutropenia:** This has been reported in patients administered vancomycin intravenously or orally. Periodically monitor leukocyte count. (5.4)
- **Infusion Reactions:** Hypotension, including shock and cardiac arrest, wheezing, dyspnea, urticaria, muscular, chest pain and vancomycin infusion reaction which manifests as pruritus and erythema that involves the face, neck and upper torso may occur with rapid intravenous administration. To reduce the risk of infusion reactions, administer Vancomycin Hydrochloride for Injection in a diluted solution over a period of 60 minutes or greater and also prior to intravenous anesthetic agents. (2.1, 5.5)
- **Phlebitis:** To reduce the risk of local irritation and phlebitis administer Vancomycin Hydrochloride for Injection by a secure intravenous route of administration. (5.6)
- ***Clostridioides difficile*-Associated Diarrhea:** Evaluate patients if diarrhea occurs. (5.7)
- **Development of Drug-Resistant Bacteria:** Prescribing Vancomycin Hydrochloride for Injection in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and increases the risk of the development of drug resistant bacteria. (5.9)

ADVERSE REACTIONS

The common adverse reactions following **intravenously, and orally administered** vancomycin were acute kidney injury, hearing loss, neutropenia, anaphylaxis, vancomycin infusion reaction. (6.1)

The most common adverse reaction of **orally administered** vancomycin (> 10%) were nausea, abdominal pain, and hypokalemia. (6.1)

To report **SUSPECTED ADVERSE REACTIONS**, contact Hikma Injectables USA Inc. at 1-877-845-0689 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- **Anesthetic Agents:** Concomitant administration of vancomycin and anesthetic agents has been associated with erythema and histamine-like flushing. (2.1, 7.1)
- **Piperacillin/Tazobactam:** Increased incidence of acute kidney injury in patients administered concomitant piperacillin/tazobactam and vancomycin as compared to vancomycin alone. Monitor kidney function in patients (7.2)

See 17 for **PATIENT COUNSELING INFORMATION**.

Revised: 1/2023

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 1.1 Septicemia
- 1.2 Infective Endocarditis
- 1.3 Skin and Skin Structure Infections
- 1.4 Bone Infections
- 1.5 Lower Respiratory Tract Infections
- 1.6 *Clostridioides difficile*-Associated Diarrhea
- 1.7 Enterocolitis Caused by *S. aureus* (including methicillin-resistant strains)
- 1.8 Limitations of Use
- 1.9 Usage

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Administration Instructions for Intravenous Use
- 2.2 Intravenous Dosage in Adult and Pediatric Patients with Normal Renal Function
- 2.3 Orally Administered Dosage in Adult and Pediatric Patients
- 2.4 Intravenous Dosage in Patients with Renal Impairment
- 2.5 Directions for the Preparation, Dilution, and Storage of the Pharmacy Bulk Package for Intravenous Use
- 2.6 Compatibility with Intravenous Fluids for Intravenous Use
- 2.7 Incompatibilities for Intravenous Use
- 2.8 Preparation of Vancomycin Hydrochloride for Injection for Oral Administration
- 2.9 Storage of Reconstituted and Diluted Vancomycin Hydrochloride for Injection for Intravenous or Oral Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Nephrotoxicity
- 5.2 Ototoxicity
- 5.3 Severe Dermatologic Reactions
- 5.4 Neutropenia

- 5.5 Infusion Reactions
- 5.6 Phlebitis and Adverse Reactions with Unapproved Routes of Administration
- 5.7 *Clostridioides difficile*-Associated Diarrhea (CDAD)
- 5.8 Hemorrhagic Occlusive Retinal Vasculitis (HORV)
- 5.9 Development of Drug-Resistant Bacteria
- 5.10 Potential for Systemic Absorption after Oral Administration

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Anesthetic Agents
- 7.2 Piperacillin-Tazobactam
- 7.3 Ototoxic and/or Nephrotoxic Drugs

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Septicemia

Vancomycin Hydrochloride for Injection **administered intravenously** is indicated in adults and pediatric patients less than 18 years of age for the treatment of septicemia due to:

- Susceptible isolates of methicillin-resistant *Staphylococcus aureus* (MRSA) and coagulase negative staphylococci.
- Methicillin-susceptible staphylococci in penicillin-allergic patients, or those patients who cannot receive or who have failed to respond to other drugs, including penicillins or cephalosporins.

1.2 Infective Endocarditis

Vancomycin Hydrochloride for Injection **administered intravenously** is indicated in adults and pediatric patients less than 18 years of age for the treatment of infective endocarditis due to:

- Susceptible isolates of MRSA.
- Viridans group streptococci *Streptococcus gallolyticus* (previously known as *Streptococcus bovis*), *Enterococcus* species and *Corynebacterium* species. For enterococcal endocarditis, use Vancomycin Hydrochloride for Injection in combination with an aminoglycoside.
- Methicillin-susceptible staphylococci in penicillin-allergic patients, or those patients who cannot receive or who have failed to respond to other drugs, including penicillins or cephalosporins.

Vancomycin Hydrochloride for Injection **administered intravenously** is indicated in adults and pediatric patients less than 18 years of age for the treatment of early-onset prosthetic valve endocarditis caused by *Staphylococcus epidermidis* in combination with rifampin and an aminoglycoside.

1.3 Skin and Skin Structure Infections

Vancomycin Hydrochloride for Injection **administered intravenously** is indicated in adults and pediatric patients less than 18 years of age for the treatment of skin and skin structure infections due to:

- Susceptible isolates of MRSA and coagulase negative staphylococci.
- Methicillin-susceptible staphylococci in penicillin-allergic patients, or those patients who cannot receive or who have failed to respond to other drugs, including penicillins or cephalosporins.

1.4 Bone Infections

Vancomycin Hydrochloride for Injection **administered intravenously** is indicated in adults and pediatric patients less than 18 years of age for the treatment of bone infections due to:

- Susceptible isolates of MRSA and coagulase negative staphylococci.
- Methicillin-susceptible staphylococci in penicillin-allergic patients, or those patients who cannot receive or who have failed to respond to other drugs, including penicillins or cephalosporins.

1.5 Lower Respiratory Tract Infections

Vancomycin Hydrochloride for Injection **administered intravenously** is indicated in adults and pediatric patients less than 18 years of age for the treatment of lower respiratory tract infections due to:

- Susceptible isolates of MRSA
- Methicillin-susceptible staphylococci in penicillin-allergic patients, or those patients who cannot receive or who have failed to respond to other drugs, including penicillins or cephalosporins.

1.6 *Clostridioides difficile*-Associated Diarrhea

Vancomycin Hydrochloride for Injection **administered orally** is indicated for the treatment of *C. difficile*-associated diarrhea (CDAD) in adult and pediatric patients less than 18 years of age.

1.7 Enterocolitis Caused by *S. aureus* (including methicillin-resistant strains)

Vancomycin Hydrochloride for Injection **administered orally** is indicated for the treatment of enterocolitis caused by susceptible isolates of *Staphylococcus aureus* in adults and pediatric patients less than 18 years of age.

1.8 Limitations of Use

Vancomycin Hydrochloride for Injection **administered intravenously** is not approved for the treatment of the following conditions because it is not effective:

- *C. difficile*-Associated Diarrhea
- Enterocolitis caused by susceptible isolates of *Staphylococcus aureus*

Vancomycin Hydrochloride for Injection **administered orally** is not approved for the treatment of the following conditions because it is not effective:

- Septicemia due to susceptible isolates of MRSA or methicillin-susceptible staphylococci
- Infective endocarditis due to susceptible isolates of MRSA, methicillin susceptible staphylococci, Viridans group streptococci *Streptococcus gallolyticus*, *Enterococcus* species and *Corynebacterium* species, or for the treatment of early-onset prosthetic valve endocarditis caused by *Staphylococcus epidermidis* in combination with rifampin and an aminoglycoside
- Skin and skin structure infections due to susceptible isolates of MRSA and methicillin susceptible staphylococci
- Bone infections due to susceptible isolates of MRSA and lower respiratory tract infections due to susceptible isolates of MRSA and methicillin susceptible staphylococci

1.9 Usage

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Vancomycin

Hydrochloride for Injection and other antibacterial drugs, Vancomycin Hydrochloride for Injection should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Instructions for Intravenous Use

- Vancomycin Hydrochloride for Injection is supplied in pharmacy bulk packages. A pharmacy bulk package is a sterile dosage form containing many single doses. The contents of Vancomycin Hydrochloride for Injection pharmacy bulk package are intended for use by a pharmacy admixture service for addition to suitable parenteral fluids in the preparation of admixtures for intravenous infusion. The Pharmacy Bulk Package is **NOT** for direct infusion. Further dilution is required before use [see *Dosage and Administration* (2.4)].
- To reduce the risk of infusion related adverse reactions, administer diluted Vancomycin Hydrochloride for Injection by intravenous infusion over 60 minutes or greater [see *Warnings and Precautions* (5.5) and *Adverse Reactions* (6.1)].
- Diluted Vancomycin Hydrochloride for Injection concentrations of no more than 5 mg/mL are recommended in adults [see *Dosage and Administration* (2.2)]. See also age-specific recommendations [see *Dosage and Administration* (2.2, 2.3)]. In selected patients in need of fluid restriction, a concentration up to 10 mg/mL may be used. Infusion related events may occur, however, at any rate or concentration [see *Warnings and Precautions* (5.5)].
- Other drugs should not be added to the Vancomycin Hydrochloride for Injection Pharmacy Bulk Package or the diluted Vancomycin Hydrochloride for Injection [see *Dosage and Administration* (2.7)].
- Administer diluted Vancomycin Hydrochloride for Injection prior to intravenous anesthetic agents to reduce the risk of infusion related adverse reactions [see *Warnings and Precautions* (5.5)].
- Administer diluted Vancomycin Hydrochloride for Injection by a secure intravenous route of administration to avoid local irritation and phlebitis reactions [see *Warnings and Precautions* (5.6)].

2.2 Intravenous Dosage in Adult and Pediatric Patients with Normal Renal Function

Dosage in Adult Patients

The usual daily intravenous dose is 2 grams (g) divided either as 500 mg every 6 hours or 1 g every 12 hours. Administer each dose by intravenous infusion over a period of 60 minutes or greater. Other patient factors, such as age or obesity, may call for modification of the usual intravenous daily dose.

Dosage in Pediatric Patients Aged 1 Month and Older

The usual intravenous dosage of vancomycin is 10 mg/kg per dose given every 6 hours. Each dose should be administered over a period of at least 60 minutes. Close monitoring of serum concentrations of vancomycin may be warranted in these patients.

Dosage in Pediatric Patients Younger than 1 Month of Age

In pediatric patients, up to the age of 1 month, the total daily intravenous dosage may be lower. In neonates, an initial dose of 15 mg/kg is suggested, followed by 10 mg/kg every 12 hours for pediatric patients in the 1st week of life and every 8 hours thereafter up to the age of 1 month. Each dose should be administered over 60 minutes. In premature infants, vancomycin clearance decreases as postconceptional age decreases. Therefore, longer dosing intervals may be necessary in premature infants. Close monitoring of serum concentrations of vancomycin is recommended in these patients.

2.3 Orally Administered Dosage in Adult and Pediatric Patients

Dosage in Adult Patients

- *C. difficile*-associated diarrhea: The recommended dose is 125 mg administered orally 4 times daily

for 10 days.

- Staphylococcal enterocolitis: Total daily dosage is 500 mg to 2 g administered orally in 3 or 4 divided doses for 7 days to 10 days.

Dosage in Pediatric Patients (Less than 18 years of age)

For both *C. difficile*-associated diarrhea and staphylococcal enterocolitis, the usual daily dosage is 40 mg/kg in 3 or 4 divided doses for 7 days to 10 days. The total daily dosage should not exceed 2 g.

2.4 Intravenous Dosage in Patients with Renal Impairment

Dosage adjustment must be made in adult and pediatric patients with renal impairment. The initial dose should be no less than 15 mg/kg, in adult patients with any degree of renal impairment.

In premature infants and the elderly, greater dosage reductions than expected may be necessary because of decreased renal function. Measure trough vancomycin serum concentrations to guide therapy, especially in seriously ill patients with changing renal function.

For functionally anephric patients, an initial dose of 15 mg/kg of body weight should be given to achieve prompt therapeutic serum concentration. A dose of 1.9 mg/kg/24 hr should be given after the initial dose of 15 mg/kg.

2.5 Directions for the Preparation, Dilution and Storage of the Pharmacy Bulk Package for Intravenous Use

Preparation

- Vancomycin Hydrochloride for Injection pharmacy bulk package is **not for direct intravenous infusion**. Prior to intravenous administration, contents of the Vancomycin Hydrochloride for Injection pharmacy bulk package **must be reconstituted and further diluted**. No preservative is present in this product. Aseptic technique must be used in the preparation of final IV solution.
- Prepare Vancomycin Hydrochloride for Injection pharmacy bulk packages for use in a pharmacy admixture service only in a suitable work area, such as a laminar flow hood. They should be hung by the integral hanger provided and suspended as a unit in the laminar flow hood.
- Using aseptic technique, penetrate the container closure only one time utilizing a suitable sterile dispensing set or transfer device which allows measured distribution dispensing of the contents.
- Use of a syringe and needle is not recommended as it may cause leakage. Swab bottle stopper with an antiseptic solution prior to inserting the dispensing set into the bottle using aseptic technique.
- Once the sterile dispensing set has been inserted into the container, withdrawal of the contents from the Vancomycin Hydrochloride for Injection pharmacy bulk package should be accomplished without delay immediately. However, if this is not possible, a maximum time of **4 hours** from the initial closure entry may be permitted to complete fluid transfer operations.
- Discard the container and any unused portion in the container no later than **4 hours** after initial closure puncture the container closure has been penetrated.

Reconstitution and Dilution

5 g Pharmacy Bulk Package Vial

At the time of use, reconstitute by adding 45 mL of Sterile Water for Injection to the 5 g vial of dry, sterile vancomycin powder. The resultant solution will contain vancomycin equivalent to about 100 mg/mL. After reconstitution, further dilution is required using one of the compatible intravenous diluents listed below [see *Dosage and Administration* (2.6)], prior to intravenous administration.

Reconstituted solutions of vancomycin (100 mg/mL) must be further diluted. A dose of 500 mg (5 mL) must be diluted with at least 100 mL of a suitable infusion. For doses of 1 gram (10 mL), at least 200 mL of solution must be used. The desired dose diluted in this manner should be administered by intermittent IV infusion over a period of at least 60 minutes.

10 g Pharmacy Bulk Package Vial

At the time of use, reconstitute by adding 90 mL of Sterile Water for Injection to the 10 g vial of dry, sterile vancomycin powder. The resultant solution will contain vancomycin equivalent to about 100 mg/mL. After reconstitution, further dilution is required using one of the compatible intravenous diluents listed below [see *Dosage and Administration* (2.6)], prior to intravenous administration.

Reconstituted solutions of vancomycin (100 mg/mL) must be further diluted. A dose of 500 mg (5 mL) must be diluted with at least 100 mL of a suitable infusion solution. For doses of 1 gram (10 mL), at least 200 mL of solution must be used. The desired dose diluted in this manner should be administered by intermittent IV infusion over a period of at least 60 minutes.

Parenteral drug products should be visually inspected for particulate matter and discoloration prior to administration, whenever solution and container permit.

2.6 Compatibility with Intravenous Fluids for Intravenous Use

The following diluents are physically and chemically compatible with approximately 4.5 mg/mL of vancomycin hydrochloride:

- 5% Dextrose Injection, USP
- 5% Dextrose Injection and 0.9% Sodium Chloride Injection, USP
- Lactated Ringer's Injection, USP
- 5% Dextrose and Lactated Ringer's Injection, USP
- 0.9% Sodium Chloride Injection, USP

2.7 Incompatibilities for Intravenous Use

Vancomycin Hydrochloride for Injection reconstituted solution has a low pH and may cause chemical or physical instability when it is mixed with other compounds.

Mixtures of solutions of vancomycin and beta-lactam antibacterial drugs have been shown to be physically incompatible. The likelihood of precipitation increases with higher concentrations of vancomycin. It is recommended to adequately flush the intravenous lines between the administration of these antibacterial drugs. It is also recommended to dilute solutions of vancomycin to 5 mg/mL or less.

2.8 Preparation of Vancomycin Hydrochloride for Injection for Oral Administration

Preparation of the Pharmacy Bulk Package for Oral Use

1. Reconstitute Vancomycin Hydrochloride for Injection to 100 mg/mL by adding the appropriate amount of sterile water for injection:

Table 1: Reconstitution-Volume of Sterile Water for Injection to be Added	
Vancomycin Hydrochloride for Injection Strength per Vial	Volume of Sterile Water for Injection to be added to achieve 100 mg/mL Vancomycin Hydrochloride Solution
5 g	45 mL
10 g	90 mL

2. Sweeteners may be added to this solution to improve the taste for oral administration following the steps outlined in *Instructions for How to Prepare and Add Sweeteners* below.

Instructions for How to Prepare and Add Sweeteners

1. Prepare the sweetening agent separately from the reconstituted Vancomycin hydrochloride solution in a plastic-container following one of the preparation options below:
 - Mix sucrose and sterile water for injection in a 1:2 ratio (e.g., 10 g sucrose and 20 mL sterile water for injection) OR
 - Mix agave nectar with an equal volume of sterile water for injection
2. Mix the sweetening agent with the reconstituted Vancomycin hydrochloride solution in a 3:1 ratio (e.g., 30 mL sweetening agent with 10 mL vancomycin solution) to obtain a sweetened solution containing 25 mg/mL of vancomycin that is suitable for oral administration.

The diluted solution may be administered orally or via a nasogastric tube [see *Dosage and Administration* (2.3)].

2.9 Storage of Reconstituted and Diluted Vancomycin Hydrochloride for Injection for Intravenous or Oral Administration

Intravenous Administration:

Administer the compounded mixtures within **4** hours after preparation when storing at room temperature and 6 days when stored refrigerated (14 days for solutions made with 0.9% sodium chloride and 5% dextrose).

Oral Administration:

Store the diluted solution for oral use in a plastic container refrigerated at 2 °C to 8 °C (36°F to 46°F) for up to 10 days.

3 DOSAGE FORMS AND STRENGTHS

Vancomycin Hydrochloride for Injection, in a Pharmacy Bulk Package is a sterile white, almost white to tan to brown spray-dried powder in vials each containing vancomycin hydrochloride equivalent to 5 g or 10 g vancomycin base.

4 CONTRAINDICATIONS

Vancomycin Hydrochloride for Injection is contraindicated in patients with known hypersensitivity to vancomycin and polysorbate 80.

5 WARNINGS AND PRECAUTIONS

5.1 Nephrotoxicity

Vancomycin Hydrochloride for Injection administered intravenously or orally can result in acute kidney injury (AKI), including acute renal failure, mainly due to interstitial nephritis or less commonly acute tubular necrosis. AKI is manifested by increasing blood urea nitrogen (BUN) and serum creatinine (Cr). The risk of AKI increases with higher vancomycin serum levels, prolonged exposure, concomitant administration of other nephrotoxic drugs, concomitant administration of piperacillin-tazobactam [see *Drug Interactions* (7.2)], volume depletion, pre-existing renal impairment and in critically ill patients and patients with co-morbid conditions that predispose to renal impairment.

Monitor serum vancomycin concentrations and renal function in all patients receiving Vancomycin Hydrochloride for Injection intravenously.

Nephrotoxicity (e.g., reports of renal failure, renal impairment, blood creatine increased) has occurred following oral vancomycin hydrochloride capsule therapy in randomized controlled clinical studies and can occur either during or after completion of therapy. The risk of nephrotoxicity is increased in patients >65 years of age [see *Adverse Reactions* (6.1) and *Use in Specific Populations* (8.5)]. In patients over 65 years of age, including those with normal renal function prior to treatment, renal function should be monitored during and following treatment with oral Vancomycin Hydrochloride for Injection therapy to detect potential vancomycin induced nephrotoxicity.

More frequent monitoring is recommended in patients with comorbidities that predispose to impairment in renal function or are concomitantly receiving other nephrotoxic drugs, in critically ill patients, in patients with changing renal function, and in patients requiring higher therapeutic vancomycin levels. If acute kidney injury occurs, discontinue Vancomycin Hydrochloride for Injection or reduce the dose.

5.2 Ototoxicity

Ototoxicity has occurred in patients administered vancomycin intravenously or orally. It may be transient or permanent. Ototoxicity manifests as tinnitus, hearing loss, dizziness or vertigo. The risk is higher in older patients, patients who are receiving higher doses, who have an underlying hearing loss, who are receiving concomitant therapy with another ototoxic agent, such as an aminoglycoside or who have underlying renal impairment. Monitor for signs and symptoms of ototoxicity during oral or intravenous

therapy with Vancomycin Hydrochloride for Injection. Discontinue Vancomycin Hydrochloride for Injection if ototoxicity occurs. Serial tests of auditory function may be helpful in order to minimize the risk of ototoxicity.

5.3 Severe Dermatologic Reactions

Severe dermatologic reactions such as toxic epidermal necrolysis (TEN), Stevens-Johnson syndrome (SJS), drug reaction with eosinophilia and systemic symptoms (DRESS), acute generalized exanthematous pustulosis (AGEP), and linear IgA bullous dermatosis (LABD) have been reported in association with the use of vancomycin. Cutaneous signs or symptoms reported include skin rashes, mucosal lesions, and blisters.

Discontinue Vancomycin Hydrochloride for Injection at the first appearance of signs and symptoms of TEN, SJS, DRESS, AGEP, or LABD.

5.4 Neutropenia

Reversible neutropenia has been reported in patients administered vancomycin intravenously or orally. Patients who will undergo prolonged therapy with Vancomycin Hydrochloride for Injection or those who are receiving concomitant drugs which may cause neutropenia should have periodic monitoring of the leukocyte count.

5.5 Infusion Reactions

Hypotension, including shock and cardiac arrest, wheezing, dyspnea, urticaria, muscular and chest pain may occur with rapid Vancomycin Hydrochloride for Injection intravenous administration. The reactions may be more severe in younger patients, particularly children, and in patients receiving concomitant muscle relaxant anesthetics. Rapid intravenous administration of Vancomycin Hydrochloride for Injection may also be associated with vancomycin infusion reaction, which manifests as pruritus and erythema that involves the face, neck and upper torso. Infusion-related adverse reactions are related to both the concentration and the rate of administration of Vancomycin Hydrochloride for Injection. Infusion-related adverse reactions may occur, however, at any rate or concentration.

Administer Vancomycin Hydrochloride for Injection in a diluted solution over a period of 60 minutes or greater to reduce the risk of infusion-related adverse reactions. In selected patients in need of fluid restriction, a concentration up to 10 mg/mL may be used; use of such higher concentrations may increase the risk of infusion-related adverse reactions [see *Nonclinical Toxicology* (13.2)]. Administer prior to intravenous anesthetic agents when feasible. Stop the infusion if a reaction occurs.

5.6 Phlebitis and Adverse Reactions with Unapproved Routes of Administration

Inflammation at the site of injection of vancomycin has been reported. Vancomycin is irritating to tissue and must be given by a secure intravenous route of administration to reduce the risk of local irritation and phlebitis. Thrombophlebitis may occur, the frequency and severity of which can be minimized by slow infusion of the drug and by rotation of venous access sites.

Administration of Vancomycin Hydrochloride for Injection by intramuscular (IM), intraperitoneal, intrathecal intraventricular, or intravitreal routes has not been approved and is not recommended. The safety and efficacy of vancomycin administered by these routes of administration have not been established by adequate and well controlled trials. Pain, tenderness, and necrosis occur with IM injection of vancomycin or with inadvertent extravasation. Intraperitoneal administration during continuous ambulatory peritoneal dialysis (CAPD) can result in chemical peritonitis. Manifestations range from cloudy dialysate alone to a cloudy dialysate accompanied by variable degrees of abdominal pain and fever. This syndrome appears to be resolved after discontinuation of intraperitoneal vancomycin.

About 60% of an intraperitoneal dose of vancomycin administered during peritoneal dialysis is absorbed systemically in 6 hours. Serum concentrations of about 10 mcg/mL are achieved by intraperitoneal injection of 30 mg/kg of vancomycin. However, the safety and efficacy of the intraperitoneal use of vancomycin has not been established in adequate and well-controlled trials.

5.7 *Clostridioides difficile*-Associated Diarrhea (CDAD)

Clostridioides difficile-associated diarrhea (CDAD) has been reported with use of nearly all antibacterial

agents, including with intravenous administration of vancomycin and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibacterial use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents. If CDAD is suspected or confirmed, ongoing antibacterial use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibacterial treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

Clinically significant serum concentrations of vancomycin have been reported in some patients being treated for active *C. difficile*-induced pseudomembranous colitis after multiple oral doses of vancomycin [see *Warnings and Precautions* (5.10)].

Prolonged use of Vancomycin Hydrochloride for Injection may result in the overgrowth of non-susceptible microorganisms. Careful observation of the patient is essential. If superinfection occurs during therapy, appropriate measures should be taken. There have been reports of pseudomembranous colitis due to *C. difficile* developing in patients who received intravenous vancomycin.

5.8 Hemorrhagic Occlusive Retinal Vasculitis (HORV)

Hemorrhagic occlusive retinal vasculitis, including permanent loss of vision, occurred in patients receiving intracameral or intravitreal administration of vancomycin during or after cataract surgery. The safety and efficacy of vancomycin administered by the intracameral or the intravitreal route have not been established by adequate and well-controlled trials. Vancomycin Hydrochloride for Injection (intravenously and orally administered) is not indicated for the prophylaxis of endophthalmitis.

5.9 Development of Drug-Resistant Bacteria

Prescribing Vancomycin Hydrochloride for Injection (intravenously and orally administered) in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

5.10 Potential for Systemic Absorption after Oral Administration

Clinically significant serum concentrations have been reported in some patients who have taken multiple oral doses of oral vancomycin for active *C. difficile*-associated diarrhea. Some patients with inflammatory disorders of the intestinal mucosa also may have significant systemic absorption of vancomycin. These patients may be at risk for the development of adverse reactions associated with higher doses of oral vancomycin; therefore, monitoring of serum concentrations of vancomycin may be appropriate in some instances, e.g., in patients with renal insufficiency and/or colitis or in those receiving concomitant therapy with an aminoglycoside antibacterial drug.

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Nephrotoxicity [see *Warnings and Precautions* (5.1)]
- Ototoxicity [see *Warnings and Precautions* (5.2)]
- Severe Dermatologic Reactions [see *Warnings and Precautions* (5.3)]
- Neutropenia [see *Warnings and Precautions* (5.4)]
- Infusion Reactions [see *Warnings and Precautions* (5.5)]
- *Clostridioides difficile*-Associated Diarrhea [see *Warnings and Precautions* (5.7)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse Reactions in Patients Receiving Intravenously and Orally Administered Vancomycin

The following adverse reactions associated with the use of intravenously and orally administered

vancomycin were identified in clinical trials:

Renal and urinary disorders: Acute kidney injury and interstitial nephritis [see *Warnings and Precautions* (5.1)]

Ear and labyrinth disorders: Tinnitus, hearing loss, vertigo [see *Warnings and Precautions* (5.2)]

Skin and subcutaneous tissue disorders: Erythema (especially of the face, neck and upper torso) and pruritus which are manifestations of rashes including exfoliative dermatitis, toxic epidermal necrolysis (TEN), Stevens-Johnson syndrome (SJS), Linear IgA bullous dermatosis (LABD) [see *Warnings and Precautions* (5.3)].

Blood and lymphatic system disorders: Agranulocytosis, neutropenia, pancytopenia, leukopenia, thrombocytopenia, eosinophilia [see *Warnings and Precautions* (5.4)]

Immune system disorders: Hypersensitivity reactions including anaphylaxis and vancomycin infusion reaction [see *Warnings and Precautions* (5.5)]

General disorders and administration site conditions: General discomfort, fever, chills, phlebitis, injection site irritation, injection site pain and necrosis following intramuscular injection, chemical peritonitis following intraperitoneal administration. Vancomycin Hydrochloride for Injection is not approved for intramuscular and intraperitoneal administration [see *Warnings and Precautions* (5.6)]

Gastrointestinal disorders: Pseudomembranous colitis [see *Warnings and Precautions* (5.7)]

Cardiac disorders: Cardiac arrest, chest pain

Laboratory abnormalities: Elevated blood urea nitrogen, elevated serum creatinine

Musculoskeletal and connective tissue disorders: Muscle pain

Nervous system disorders: Dizziness

Respiratory, thoracic and mediastinal disorders: Wheezing, dyspnea

Vascular disorders: Hypotension, shock, vasculitis

Adverse Reactions in Patients Receiving Oral Administration of Vancomycin Hydrochloride Capsules

The data described below reflect exposure to vancomycin hydrochloride capsules in 260 adult subjects in two Phase 3 clinical trials for the treatment of diarrhea associated with *C. difficile*. In both trials, subjects received vancomycin hydrochloride capsules 125 mg orally four times daily. The mean duration of treatment was 9.4 days. The median age of patients was 67, ranging between 19 and 96 years of age. Patients were predominantly Caucasian (93%) and 52% were male.

Adverse reactions occurring in $\geq 5\%$ of vancomycin hydrochloride capsules-treated subjects are shown in Table 2. The most common adverse reactions associated with vancomycin hydrochloride capsules ($\geq 10\%$) were nausea, abdominal pain, and hypokalemia.

Table 2: Common ($\geq 5\%$) Adverse Reactions^a for Vancomycin Hydrochloride Capsules Reported in Clinical Trials for Treatment of Diarrhea Associated with *C. difficile*

System/Organ Class	Adverse Reaction	Vancomycin Hydrochloride Capsule % (N=260)
Gastrointestinal disorders	Nausea	17
	Abdominal pain	15
	Vomiting	9
	Diarrhea	9
	Flatulence	8
General disorders and administration site conditions	Pyrexia	9
	Edema peripheral	6
	Fatigue	5
Infections and infestations	Urinary tract infection	8
Metabolism and nutrition disorders	Hypokalemia	13
Musculoskeletal and	Back pain	6

connective tissue disorders		
Nervous system disorders	Headache	7
^a Adverse reaction rates were derived from the incidence of treatment-emergent adverse events.		

Nephrotoxicity (e.g., reports of renal failure, renal impairment, blood creatinine increased) occurred in 5% of subjects treated with vancomycin hydrochloride capsules. Nephrotoxicity following vancomycin hydrochloride capsules typically first occurred within one week after completion of treatment (median day of onset was Day 16). Nephrotoxicity following vancomycin hydrochloride capsules occurred in 6% of subjects >65 years of age and 3% of subjects ≤65 years of age.

The incidences of hypokalemia, urinary tract infection, peripheral edema, insomnia, constipation, anemia, depression, vomiting, and hypotension were higher among subjects >65 years of age than in subjects ≤65 years of age.

Discontinuation of study drug due to adverse events occurred in 7% of subjects treated with vancomycin hydrochloride capsules. The most common adverse events leading to discontinuation of vancomycin hydrochloride capsules were *C. difficile* colitis (<1%), nausea (<1%), and vomiting (<1%).

6.2 Postmarketing Experience

The following adverse reactions have been identified during postmarketing use of vancomycin (administered orally and intravenously). Because these reactions are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Skin and Subcutaneous Tissue Disorders: Drug reaction with eosinophilia and systemic symptoms (DRESS), acute generalized exanthematous pustulosis (AGEP) [see *Warnings and Precautions* (5.3)].

Ototoxicity: Cases of hearing loss associated with intravenously administered vancomycin have been reported. Most of these patients had kidney dysfunction or a preexisting hearing loss or were receiving concomitant treatment with an ototoxic drug [see *Warnings and Precautions* (5.2)]. Vertigo, dizziness, and tinnitus have been reported.

Hematopoietic: Reversible neutropenia, usually starting 1 week or more after onset of intravenous therapy with vancomycin or after a total dose of more than 25 g, has been reported for several dozen patients. Neutropenia appears to be promptly reversible when vancomycin is discontinued. Thrombocytopenia has been reported.

Miscellaneous: Patients have been reported to have had anaphylaxis, drug fever, chills, nausea, eosinophilia, and cases of vasculitis in association with the administration of vancomycin. A condition has been reported with oral vancomycin that is similar to the intravenous vancomycin-induced syndrome with symptoms consistent with anaphylactoid reactions, including hypotension, wheezing, dyspnea, urticaria, pruritus, flushing of the upper body ("vancomycin infusion reaction"), pain and muscle spasm of the chest and back [see *Warnings and Precautions* (5.5)]. These reactions usually resolve within 20 minutes but may persist for several hours.

7 DRUG INTERACTIONS

7.1 Anesthetic Agents

Concomitant administration of vancomycin and anesthetic agents has been associated with erythema and histamine-like flushing [see *Warnings and Precautions* (5.5) and *Use in Specific Populations* (8.4)].

7.2 Piperacillin-Tazobactam

Studies have detected an increased incidence of acute kidney injury in patients administered concomitant piperacillin/tazobactam and vancomycin as compared to vancomycin alone. Monitor kidney function in patients receiving concomitant piperacillin/tazobactam and Vancomycin Hydrochloride for Injection. No pharmacokinetic interactions have been noted between piperacillin/tazobactam and vancomycin.

7.3 Ototoxic and/or Nephrotoxic Drugs

Concurrent and/or sequential systemic or topical use of other potentially, neurotoxic and/or nephrotoxic

drugs requires more frequent monitoring of renal function.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no available data on vancomycin use in pregnant women to inform a drug associated risk of major birth defects or miscarriage. Available published data on vancomycin use in pregnancy during the second and third trimesters have not shown an association with adverse pregnancy related outcomes (*see Data*). Vancomycin did not show adverse developmental effects when administered intravenously to pregnant rats and rabbits during organogenesis at doses less than or equal to the recommended maximum human dose (*see Data*). Systemic absorption of vancomycin is low following oral administration of vancomycin hydrochloride capsules; however, absorption may vary depending on various factors [*see Clinical Pharmacology (12.3)*].

The estimated background risk of major birth defects and miscarriage for the indicated population(s) is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Data

Human Data

A published study evaluated hearing loss and nephrotoxicity in infants of 10 pregnant intravenous drug users treated with intravenously administered vancomycin for suspected or documented methicillin-resistant *Staphylococcus-aureus* (MRSA) infections in the second or third trimester. The comparison groups were 10 non-intravenous drug-dependent patients who received no treatment, and 10 untreated intravenous drug-dependent patients who served as substance abuse controls. No infant in the vancomycin exposed group had abnormal sensorineural hearing at 3 months of age or nephrotoxicity.

A published prospective study assessed outcomes in 55 pregnant women with a positive Group B streptococcus (GBS) culture and a high-risk penicillin allergy with resistance to clindamycin or unknown sensitivity who were administered vancomycin intravenously at the time of delivery. Vancomycin dosing ranged from the standard 1 g intravenously every 12 hours to 20 mg/kg intravenous every 8 hours (maximum individual dose 2 g). No major adverse reactions were recorded either in the mothers or their newborns. None of the newborns had sensorineural hearing loss. Neonatal renal function was not examined, but all of the newborns were discharged in good condition.

Animal Data

Vancomycin did not cause fetal malformations when administered during organogenesis to pregnant rats (gestation days 6 to 15) and rabbits (gestation days 6 to 18) at the equivalent recommended maximum human dose (based on body surface area comparisons) of 200 mg/kg/day IV to rats or 120 mg/kg/day IV to rabbits.¹ No effects on fetal weight or development were seen in rats at the highest dose tested or in rabbits given 80 mg/kg/day (approximately 1 and 0.8 times the recommended maximum human dose based on body surface area, respectively). Maternal toxicity was observed in rats (at doses 120 mg/kg and above) and rabbits (at 80 mg/kg and above). There were no oral embryo-fetal toxicity studies conducted in animals with this Vancomycin Injection product.

8.2 Lactation

Risk Summary

Vancomycin is present in human milk following intravenous administration, however, there are insufficient data to inform the levels. There are no available data on vancomycin presence in milk following oral administration. Systemic absorption of vancomycin is low following oral administration of vancomycin hydrochloride capsules; however, absorption may vary depending on various factors [*see Clinical Pharmacology (12.3)*]. There are no data on the effects of vancomycin on the breastfed infant or milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Vancomycin Hydrochloride for Injection and any potential adverse effects on the breastfed infant from Vancomycin Hydrochloride for Injection or from the underlying

maternal condition.

8.4 Pediatric Use

Vancomycin Hydrochloride for Injection administered intravenously is indicated in pediatric patients less than 18 years of age for the treatment of septicemia, infective endocarditis, skin and skin structure infections, bone infections and lower respiratory tract infections [see *Indications and Usage* (1.1 to 1.5) and *Dosage and Administration* (2.2)].

Vancomycin Hydrochloride for Injection administered orally is indicated in pediatric patients less than 18 years of age for the treatment of *Clostridioides difficile*-associated diarrhea and enterocolitis caused by *Staphylococcus aureus* (including methicillin-resistant strains) [see *Indications and Usage* (1.6, 1.7) and *Dosage and Administration* (2.3)].

In pediatric patients, monitor vancomycin serum concentration and renal function when administering Vancomycin Hydrochloride for Injection intravenously [see *Dosage and Administration* (2.2) and *Warnings and Precautions* (5.1)]. More severe infusion related reactions related to Vancomycin Hydrochloride for Injection administration may occur in pediatric patients. Concomitant administration of vancomycin and intravenous anesthetic agents has been associated with erythema and histamine-like flushing in all patients including pediatric patients [see *Warnings and Precautions* (5.5)].

8.5 Geriatric Use

Clinical studies with another vancomycin hydrochloride drug product in *C. difficile*-associated diarrhea have demonstrated that geriatric subjects are at increased risk of developing nephrotoxicity following treatment with oral vancomycin hydrochloride, which may occur during or after completion of therapy. In patients over 65 years of age, including those with normal renal function prior to treatment, renal function should be monitored during and following treatment with Vancomycin Hydrochloride for Injection to detect potential vancomycin induced nephrotoxicity [see *Warnings and Precautions* (5.1)] and *Adverse Reactions* (6.1)].

Patients over 65 years of age may take longer to respond to therapy compared to patients 65 years of age and younger. Clinicians should be aware of the importance of appropriate duration of Vancomycin Hydrochloride for Injection treatment in patients over 65 years of age and not discontinue or switch to alternative treatment prematurely.

Vancomycin Hydrochloride for Injection is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection [see *Dosage and Administration* (2.4)], and it may be useful to monitor renal function [see *Warnings and Precautions* (5.1)].

8.6 Renal Impairment

Dosage adjustment must be made in patients with impaired renal function [see *Dosage and Administration* (2.3)]. Measure trough vancomycin serum concentrations to guide intravenous therapy, especially in seriously ill patients with changing renal function.

10 OVERDOSAGE


Supportive care is advised, with maintenance of glomerular filtration. Vancomycin is poorly removed by dialysis. Hemofiltration and hemoperfusion with polysulfone resin have been reported to result in increased vancomycin clearance.

For current information on the management of overdose, contact the National Poison Control Center at 1-800-222-1222 or www.poisson.org.

11 DESCRIPTION

Vancomycin Hydrochloride for Injection contains the hydrochloride salt of vancomycin, a tricyclic glycopeptide antibacterial derived from *Amycolatopsis orientalis* (formerly *Nocardia orientalis*). The chemical name for vancomycin hydrochloride is (Sa)-(3S,6R,7R,22R,23S, 26S,36R,38aR)-44-[[2-O-(3-Amino-2,3,6-trideoxy-3-C-methyl- α -L-lyxo-hexopyranosyl)- β -D-glucopyranosyl]oxy]-3-

(carbamoylmethyl)-10,19-dichloro-2,3,4,5,6,7,23,24,25,26,36,37,38,38atetradecahydro-7,22,28,30,32-pentahydroxy-6-[(2*R*)-4-methyl-2-(methylamino)valeramido]2,5,24,38,39-pentaoxo-22*H*-8,11:18,21-dietheno-23,36-(iminomethano)-13,16:31,35-dimetheno-1*H*,16*H*-[1,6,9]oxadiazacyclohexadecino[4,5-*m*][10,2,16]-benzoxadiazacyclotetracosine-26-carboxylic acid, monohydrochloride. The molecular formula is $C_{66}H_{75}Cl_2N_9O_{24} \cdot HCl$ and the molecular weight is 1,485.71. Vancomycin hydrochloride has the following structural formula:



Vancomycin Hydrochloride for Injection, 5 g and 10 g, in Pharmacy Bulk Package, is a sterile, preservative-free, white, almost white to tan to brown, spray-dried powder, for preparing intravenous (IV) infusions, in vials. Each 5 g vial contains 5 g vancomycin (equivalent to 5.13 g vancomycin hydrochloride), 1.25 g trehalose, 0.5 mg polysorbate 80, and hydrochloric acid used to adjust the pH. Each 10 g vial contains 10 g vancomycin (equivalent to 10.25 g vancomycin hydrochloride), 2.5 g trehalose, 1 mg polysorbate 80, and hydrochloric acid used to adjust the pH.

The spray-dried powder is reconstituted with Sterile Water for Injection which forms a clear solution and subsequently diluted prior to intravenous or oral administration [see *Dosage and Administration* (2.5, 2.8)].

A pharmacy bulk package is a sterile dosage form containing many single doses. The contents of this pharmacy bulk package are intended for use by a pharmacy admixture service for addition to suitable parenteral fluids in the preparation of admixtures for intravenous infusion. Vancomycin Hydrochloride for Injection is for intravenous or oral administration. Vancomycin Hydrochloride for Injection must be diluted with an appropriate diluent prior to intravenous administration [see *Dosage and Administration* (2.5)]. Vancomycin Hydrochloride for Injection may also be diluted with a sweetening agent for oral administration [see *Dosage and Administration* (2.8)].

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Vancomycin is an antibacterial drug [see *Microbiology* (12.4)].

12.2 Pharmacodynamics

The pharmacodynamics of vancomycin is unknown.

12.3 Pharmacokinetics

General Pharmacokinetics

In subjects with normal kidney function, multiple intravenous dosing of 1 g of vancomycin (15 mg/kg) infused over 60 minutes produces mean plasma concentrations of approximately 63 mcg/mL immediately after the completion of infusion, mean plasma concentrations of approximately 23 mcg/mL 2 hours after infusion, and mean plasma concentrations of approximately 8 mcg/mL 11 hours after the end of the infusion. Multiple dosing of 500 mg infused over 30 minutes produces mean plasma concentrations of about 49 mcg/mL at the completion of infusion, mean plasma concentrations of about 19 mcg/mL 2 hours after infusion, and mean plasma concentrations of about 10 mcg/mL 6 hours after

infusion. The plasma concentrations during multiple dosing are like those after a single dose.

Absorption

Vancomycin is poorly absorbed after oral administration. During multiple dosing of 250 mg every 8 hours for 7 doses, fecal concentrations of vancomycin in volunteers exceeded 100 mg/kg in the majority of samples. No blood concentrations were detected, and urinary recovery did not exceed 0.76%. In anephric subjects with no inflammatory bowel disease who received vancomycin oral solution 2 g for 16 days, blood concentrations of vancomycin were less than or equal to 0.66 mcg/mL in 2 of 5 subjects. No measurable blood concentrations were attained in the other 3 subjects. Following doses of 2 g daily, concentrations of drug were >3100 mg/kg in the feces and <1 mcg/mL in the serum of subjects with normal renal function who had *C. difficile*-associated diarrhea. After multiple-dose oral administration of vancomycin, measurable serum concentrations may occur in patients with active *C. difficile*-associated diarrhea, and, in the presence of renal impairment, the possibility of accumulation exists. It should be noted that the total systemic and renal clearances of vancomycin are reduced in the elderly [see *Use in Specific Populations* (8.5)].

Distribution

The volume of distribution ranges from 0.3 to 0.43 L/kg after intravenous administration. Vancomycin is approximately 55% serum protein bound as measured by ultrafiltration at vancomycin serum concentrations of 10 to 100 mcg/mL. After intravenous administration of vancomycin, inhibitory concentrations are present in pleural, pericardial, ascitic, and synovial fluids; in urine; in peritoneal dialysis fluid; and in atrial appendage tissue. Vancomycin does not readily diffuse across normal meninges into the spinal fluid; but, when the meninges are inflamed, penetration into the spinal fluid occurs.

Elimination

Mean plasma clearance is about 0.058 L/kg/h, and mean renal clearance is about 0.048 L/kg/h. The mean elimination half-life of vancomycin from plasma is 4 to 6 hours in subjects with normal renal function. In anephric patients, the mean elimination half-life is 7.5 days. Total body and renal clearance of vancomycin may be reduced in the elderly.

Metabolism

There is no *apparent* metabolism of vancomycin.

Excretion

In the first 24 hours after intravenous administration, about 75% of an administered dose of vancomycin is excreted in urine by glomerular filtration. Renal impairment slows excretion of vancomycin.

12.4 Microbiology

Mechanism of Action

The bactericidal action of vancomycin results primarily from inhibition of cell-wall biosynthesis. In addition, vancomycin alters bacterial-cell-membrane permeability and RNA synthesis.

Resistance

Vancomycin is not active *in vitro* against gram-negative bacilli, mycobacteria, or fungi. There is no cross-resistance between vancomycin and other antibacterials.

C. difficile

Isolates of *C. difficile* generally have vancomycin minimal inhibitory concentrations (MICs) of < 1 mcg/mL; however, vancomycin MICs ranging from 4 mcg/mL to 16 mcg/mL have been reported. The mechanism which mediates *C. difficile*'s decreased susceptibility to vancomycin has not been fully elucidated.

S. aureus

S. aureus isolates with vancomycin MICs as high as 1024 mcg/mL have been reported. The exact mechanism of this resistance is not clear but is believed to be due to cell wall thickening and potentially the transfer of genetic material.

Interaction with Other Antimicrobials

The combination of vancomycin and an aminoglycoside acts synergistically *in vitro* against many isolates of *Staphylococcus aureus*, *Streptococcus gallolyticus* (previously known as *Streptococcus bovis*), *Enterococcus species*, and the viridans group streptococci.

Antimicrobial Activity

Vancomycin has been shown to be active against most isolates of the following microorganisms, both *in vitro* and in clinical infections [see *Indications and Usage* (1)].

Aerobic bacteria

Gram-positive bacteria

Corynebacterium species

Enterococcus species (including *Enterococcus faecalis*)

Staphylococcus aureus (including methicillin-resistant and methicillin-susceptible isolates)

Coagulase negative staphylococci (including *S.epidermidis* and methicillin-resistant isolates)

Streptococcus gallolyticus (previously known as *Streptococcus bovis*)

Viridans group streptococci

Anaerobic bacteria

Gram-positive bacteria

Clostridioides difficile isolates associated with *C. difficile* associated diarrhea.

The following *in vitro* data are available, but their clinical significance is unknown. At least 90 percent of the following bacteria exhibit an *in vitro* minimum inhibitory concentration (MIC) less than or equal to the susceptible breakpoint for vancomycin against isolates of similar genus or organism group. However, the efficacy of vancomycin in treating clinical infections caused by these bacteria has not been established in adequate and well-controlled clinical trials.

Aerobic bacteria

Gram-positive bacteria

Listeria monocytogenes

Streptococcus pyogenes

Streptococcus pneumoniae

Streptococcus agalactiae

Anaerobic bacteria

Gram-positive bacteria

Actinomyces species

Lactobacillus species

Susceptibility Testing

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term carcinogenesis studies in animals have been conducted.

At concentrations up to 1000 mcg/mL, vancomycin had no mutagenic effect *in vitro* in the mouse lymphoma forward mutation assay or the primary rat hepatocyte unscheduled DNA synthesis assay. The concentrations tested *in vitro* were above the peak plasma vancomycin concentrations of 20 to 40 mcg/mL usually achieved in humans after slow infusion of the maximum recommended dose of 1 g. Vancomycin had no mutagenic effect *in vivo* in the Chinese hamster sister chromatid exchange assay (400 mg/kg IP) or the mouse micronucleus assay (800 mg/kg IP).

No definitive fertility studies have been conducted.

13.2 Animal Toxicology and/or Pharmacology

In animal studies, hypotension and bradycardia occurred in dogs receiving an intravenous infusion of vancomycin hydrochloride 25 mg/kg, at a concentration and infusion rate higher than recommended (concentration of 25 mg/mL and an infusion rate of 13.3 mL/min).

In a 4-week repeat-dose toxicokinetic study, electrocardiograms were obtained from dogs receiving Vancomycin Hydrochloride for Injection by the IV route, twice daily at 33 mg/kg (at a concentration of 10 mg/mL, over 30 minutes), for a total daily vancomycin dose of 66 mg/kg/day. No bradycardia was observed in the dogs when the pretest heart rates were compared to the heart rates observed 1 to 2 hours after the first daily dose during week 4.

15 REFERENCES

1. Byrd RA., Gries CL, Buening M.: Developmental Toxicology Studies of Vancomycin Hydrochloride Administered Intravenously to Rats and Rabbits. *Fundam Appl Toxicol* 1994; 23: 590-597.

16 HOW SUPPLIED/STORAGE AND HANDLING

How supplied

Vancomycin Hydrochloride for Injection is a white, almost white to tan to brown, sterile spray-dried powder in Pharmacy Bulk Package vials each containing vancomycin hydrochloride equivalent to 5 g or 10 g of vancomycin base.

The vial stopper is not made with natural rubber latex.

NDC No.	Strength Unit of Sale
0143-9163-01	Vancomycin Hydrochloride for Injection, equivalent to 5 g vancomycin in a Pharmacy Bulk Package vial, packaged individually.
0143-9164-01	Vancomycin Hydrochloride for Injection, equivalent to 10 g vancomycin in a Pharmacy Bulk Package vial, packaged individually.

Storage

Store at 20°C to 25°C (68°F to 77°F). Brief exposure to 15°C to 30°C (59°F to 86°F) is permitted [See USP Controlled Room Temperature]. Storage after reconstitution and dilution is described elsewhere in the labeling [see *Dosage and Administration* (2.5, 2.8)].

17 PATIENT COUNSELING INFORMATION

Acute Kidney Injury

Advise patients that Vancomycin Hydrochloride for Injection can result in kidney damage and that blood tests are required to monitor vancomycin blood levels and kidney function during therapy [see *Warnings and Precautions* (5.1)].

Hearing Loss or Balance Problems

Advise patients that Vancomycin Hydrochloride for Injection may result in decreased hearing and to report hearing loss or balance problems to their health care provider [see *Warnings and Precautions* (5.2)].

Severe Dermatologic Reactions

Advise patients about the signs and symptoms of serious skin manifestations. Instruct patients to stop Vancomycin Hydrochloride for Injection immediately and promptly seek medical attention at the first signs or symptoms of skin rash, mucosal lesions or blisters [see *Warnings and Precautions* (5.3)].

Infusion Reactions During or After Intravenous Use

Advise patients that generalized skin redness, skin rash, itching, flushing, muscle pain, chest pain,

shortness of breath, wheezing, or dizziness may occur during Vancomycin Hydrochloride for Injection infusion. [see *Warnings and Precautions* (5.5)].

Diarrhea

Diarrhea is a common problem caused by antibacterial drugs, including vancomycin, which usually ends when the antibacterial drug is discontinued. Sometimes after starting treatment with antibacterial drugs, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibacterial drug. If this occurs during treatment with Vancomycin Hydrochloride for Injection, patients should contact their physician as soon as possible [see *Warnings and Precautions* (5.7)].

Antibacterial Resistance

Patients should be counseled that antibacterial drugs including Vancomycin Hydrochloride for Injection should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Vancomycin Hydrochloride for Injection is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by Vancomycin Hydrochloride for Injection or other antibacterial drugs in the future.

Manufactured by:
Zhejiang Novus Pharmaceuticals Co., Ltd.
No. 125, Yuezhong Rd, Lihai Subdistrict,
Yuecheng District, Shaoxing, 312366, Zhejiang, China

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use **VANCOMYCIN HYDROCHLORIDE FOR INJECTION** safely and effectively. See full prescribing information for **VANCOMYCIN HYDROCHLORIDE FOR INJECTION**.

VANCOMYCIN HYDROCHLORIDE for injection, for intravenous use or oral use

Initial U.S. Approval: 1958

INDICATIONS AND USAGE

Vancomycin Hydrochloride for Injection is a glycopeptide antibacterial indicated in adult and pediatric patients less than 18 years of age as follows:

- Vancomycin Hydrochloride for Injection **administered intravenously** is indicated for the treatment of:
 - Septicemia (1.1)
 - Infective Endocarditis (1.2)
 - Skin and Skin Structure Infections (1.3)
 - Bone Infections (1.4)
 - Lower Respiratory Tract Infections (1.5)
- Vancomycin Hydrochloride for Injection **administered orally** is indicated for the treatment of:
 - *Clostridioides difficile*-associated diarrhea (1.6)
 - Enterocolitis caused by *Staphylococcus aureus* (including methicillin-resistant strains) (1.7)

Limitations of Use (1.8)

- Vancomycin Hydrochloride for Injection **administered intravenously** is not approved for the treatment of *C. difficile*-associated diarrhea and enterocolitis caused by susceptible isolates of *Staphylococcus aureus* because it is not effective.
- Vancomycin Hydrochloride for Injection **administered orally** is not approved for the treatment of septicemia, infective endocarditis, skin and skin structure infections, bone infections and lower respiratory tract infections because it is not effective.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Vancomycin Hydrochloride for Injection and other antibacterial drugs, Vancomycin Hydrochloride for Injection should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. (1.9)

DOSAGE AND ADMINISTRATION

See full prescribing information for important information on intravenous and oral administration, preparation and storage instructions (2.1, 2.5, 2.8).

Recommended Dosage and Administration

- **Intravenous Use:**
 - Administer Vancomycin Hydrochloride for Injection in a diluted solution by intravenous infusion over 60 minutes or greater to reduce the risk of infusion reactions.
 - **Adult Patients:** 2 grams (g) divided either as 0.5 g every 6 hours or 1 g every 12 hours (2.2)
 - **Pediatric Patients (1 month and older):** 10 mg/kg per dose given every 6 hours (2.2)
 - **Pediatric Patients (younger than 1 month of age):** See full prescribing information for recommended dosage (2.2)
 - **Patients with Renal Impairment:** See full prescribing information for recommended dosage in patients with renal impairment (2.4)
- **Oral use:**
 - **Adult Patients (18 years of age or greater):**
 - *C. difficile*-Associated Diarrhea: 125 mg administered orally 4 times daily for 10 days (2.3).
 - *Staphylococcal enterocolitis:* 500 mg to 2 grams administered orally in 3 or 4 divided doses for 7 to 10 days (2.3)
 - **Pediatric Patients (less than 18 years of age):**
 - For both *C. difficile*-associated diarrhea and staphylococcal enterocolitis: 40 mg/kg in 3 or 4 divided doses for 7 days to 10 days. The total daily dosage should not exceed 2 g (2.3)

DOSAGE FORMS AND STRENGTHS

Vancomycin Hydrochloride for Injection is a sterile spray-dried powder for injection in single-dose vials containing vancomycin hydrochloride equivalent to 500 mg or 1 gram of vancomycin base (3)

CONTRAINDICATIONS

Hypersensitivity to vancomycin, and polysorbate 80 (4)

WARNINGS AND PRECAUTIONS

- **Nephrotoxicity:** Systemic vancomycin exposure may result in acute kidney injury (AKI) including acute renal failure, mainly due to interstitial nephritis or less commonly acute tubular necrosis. Monitor serum vancomycin concentrations and renal function in patients receiving Vancomycin Hydrochloride for Injection intravenously. Monitor renal function in patients over 65 years of age receiving Vancomycin Hydrochloride for Injection orally. (5.1)
- **Ototoxicity:** Ototoxicity has occurred in patients administered vancomycin intravenously or orally. Monitor for signs and symptoms of ototoxicity during oral or intravenous therapy. Assessment of auditory function may be appropriate in some instances. (5.2)
- **Severe Dermatologic Reactions:** Discontinue Vancomycin Hydrochloride for Injection at the first appearance of skin rashes, mucosal lesions, or blisters (5.3).
- **Neutropenia:** This has been reported in patients administered vancomycin intravenously or orally. Periodically monitor leukocyte count. (5.4)
- **Infusion Reactions:** Hypotension, including shock and cardiac arrest, wheezing, dyspnea, urticaria, muscular, chest pain and vancomycin infusion reaction which manifests as pruritus and erythema that involves the face, neck and upper torso may occur with rapid intravenous administration. To reduce the risk of infusion reactions, administer Vancomycin Hydrochloride for Injection in a diluted solution over a period of 60 minutes or greater and also prior to intravenous anesthetic agents. (2.1, 5.5)
- **Phlebitis:** To reduce the risk of local irritation and phlebitis, administer Vancomycin Hydrochloride for Injection by a secure intravenous route of administration. (5.6)
- ***Clostridioides difficile*-Associated Diarrhea:** Evaluate patients if diarrhea occurs. (5.7)
- **Development of Drug-Resistant Bacteria:** Prescribing Vancomycin Hydrochloride for Injection in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and increases the risk of the development of drug resistant bacteria. (5.9)

ADVERSE REACTIONS

The common adverse reactions following **intravenously, and orally administered** vancomycin were acute kidney injury, hearing loss, neutropenia, anaphylaxis, vancomycin infusion reaction. (6.1)
The most common adverse reactions of **orally administered** vancomycin (> 10%) were nausea, abdominal pain, and hypokalemia (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact Hikma Injectables USA Inc. at 1-877-845-0689 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- **Anesthetic Agents:** Concomitant administration of vancomycin and anesthetic agents has been associated with erythema and histamine-like flushing. (2.1, 7.1)
- **Piperacillin/Tazobactam:** Increased incidence of acute kidney injury in patients administered concomitant piperacillin/tazobactam and vancomycin as compared to vancomycin alone. Monitor kidney function in patients (7.2)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 1/2023

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 1.1 Septicemia
- 1.2 Infective Endocarditis
- 1.3 Skin and Skin Structure Infections
- 1.4 Bone Infections
- 1.5 Lower Respiratory Tract Infections
- 1.6 *Clostridioides difficile*-Associated Diarrhea
- 1.7 Enterocolitis Caused by *S. aureus* (including methicillin-resistant strains)
- 1.8 Limitations of Use
- 1.9 Usage

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Administration Instructions for Intravenous Use
- 2.2 Intravenous Dosage in Adult and Pediatric Patients with Normal Renal Function
- 2.3 Orally Administered Dosage in Adult and Pediatric Patients
- 2.4 Intravenous Dosage in Patients with Renal Impairment
- 2.5 Directions for the Preparation, Dilution, and Storage of the Single-Dose Vials for Intravenous Use
- 2.6 Compatibility with Intravenous Fluids for Intravenous Use
- 2.7 Incompatibilities for Intravenous Use
- 2.8 Preparation of Vancomycin Hydrochloride for Injection for Oral Administration
- 2.9 Storage of Reconstituted and Diluted Vancomycin Hydrochloride for Injection for Intravenous or Oral Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Nephrotoxicity
- 5.2 Ototoxicity
- 5.3 Severe Dermatologic Reactions
- 5.4 Neutropenia

- 5.5 Infusion Reactions
- 5.6 Phlebitis and Adverse Reactions with Unapproved Routes of Administration
- 5.7 *Clostridioides difficile*-Associated Diarrhea (CDAD)
- 5.8 Hemorrhagic Occlusive Retinal Vasculitis (HORV)
- 5.9 Development of Drug-Resistant Bacteria
- 5.10 Potential for Systemic Absorption after Oral Administration

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Anesthetic Agents
- 7.2 Piperacillin-Tazobactam
- 7.3 Ototoxic and/or Nephrotoxic Drugs

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Septicemia

Vancomycin Hydrochloride for Injection **administered intravenously** is indicated in adults and pediatric patients less than 18 years of age for the treatment of septicemia due to:

- Susceptible isolates of methicillin-resistant *Staphylococcus aureus* (MRSA) and coagulase negative staphylococci.
- Methicillin-susceptible staphylococci in penicillin-allergic patients, or those patients who cannot receive or who have failed to respond to other drugs, including penicillins or cephalosporins.

1.2 Infective Endocarditis

Vancomycin Hydrochloride for Injection **administered intravenously** is indicated in adults and pediatric patients less than 18 years of age for the treatment of infective endocarditis due to:

- Susceptible isolates of MRSA.
- Viridans group streptococci *Streptococcus gallolyticus* (previously known as *Streptococcus bovis*), *Enterococcus* species and *Corynebacterium* species. For enterococcal endocarditis, use Vancomycin Hydrochloride for Injection in combination with an aminoglycoside.
- Methicillin-susceptible staphylococci in penicillin-allergic patients, or those patients who cannot receive or who have failed to respond to other drugs, including penicillins or cephalosporins.

Vancomycin Hydrochloride for Injection **administered intravenously** is indicated in adults and pediatric patients less than 18 years of age for the treatment of early-onset prosthetic valve endocarditis caused by *Staphylococcus epidermidis* in combination with rifampin and an aminoglycoside.

1.3 Skin and Skin Structure Infections

Vancomycin Hydrochloride for Injection **administered intravenously** is indicated in adults and pediatric patients less than 18 years of age for the treatment of skin and skin structure infections due to:

- Susceptible isolates of MRSA and coagulase negative staphylococci.
- Methicillin-susceptible staphylococci in penicillin-allergic patients, or those patients who cannot receive or who have failed to respond to other drugs, including penicillins or cephalosporins.

1.4 Bone Infections

Vancomycin Hydrochloride for Injection **administered intravenously** is indicated in adults and pediatric patients less than 18 years of age for the treatment of bone infections due to:

- Susceptible isolates of MRSA and coagulase negative staphylococci.
- Methicillin-susceptible staphylococci in penicillin-allergic patients, or those patients who cannot receive or who have failed to respond to other drugs, including penicillins or cephalosporins.

1.5 Lower Respiratory Tract Infections

Vancomycin Hydrochloride for Injection **administered intravenously** is indicated in adults and pediatric patients less than 18 years of age for the treatment of lower respiratory tract infections due to:

- Susceptible isolates of MRSA
- Methicillin-susceptible staphylococci in penicillin-allergic patients, or those patients who cannot receive or who have failed to respond to other drugs, including penicillins or cephalosporins.

1.6 *Clostridioides difficile*-Associated Diarrhea

Vancomycin Hydrochloride for Injection **administered orally** is indicated for the treatment of *C. difficile*-associated diarrhea (CDAD) in adult and pediatric patients less than 18 years of age.

1.7 Enterocolitis Caused by *S. aureus* (including methicillin-resistant strains)

Vancomycin Hydrochloride for Injection **administered orally** is indicated for the treatment of enterocolitis caused by susceptible isolates of *Staphylococcus aureus* in adults and pediatric patients less than 18 years of age.

1.8 Limitations of Use

Vancomycin Hydrochloride for Injection **administered intravenously** is not approved for the treatment of the following conditions because it is not effective:

- *C. difficile*-Associated Diarrhea
- Enterocolitis caused by susceptible isolates of *Staphylococcus aureus*

Vancomycin Hydrochloride for Injection **administered orally** is not approved for the treatment of the following conditions because it is not effective:

- Septicemia due to susceptible isolates of MRSA or methicillin-susceptible staphylococci
- Infective endocarditis due to susceptible isolates of MRSA, methicillin susceptible staphylococci, Viridans group streptococci *Streptococcus gallolyticus*, *Enterococcus* species and *Corynebacterium* species, or for the treatment of early-onset prosthetic valve endocarditis caused by *Staphylococcus epidermidis* in combination with rifampin and an aminoglycoside
- Skin and skin structure infections due to susceptible isolates of MRSA and methicillin susceptible staphylococci
- Bone infections due to susceptible isolates of MRSA and lower respiratory tract infections due to susceptible isolates of MRSA and methicillin susceptible staphylococci

1.9 Usage

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Vancomycin Hydrochloride for Injection and other antibacterial drugs, Vancomycin Hydrochloride for Injection should be used

only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Instructions for Intravenous Use

- To reduce the risk of infusion related adverse reactions, administer diluted Vancomycin Hydrochloride for Injection by intravenous infusion over 60 minutes or greater [see *Warnings and Precautions* (5.5) and *Adverse Reactions* (6.1)].
- Diluted Vancomycin Hydrochloride for Injection concentrations of no more than 5 mg/mL are recommended in adults [see *Dosage and Administration* (2.2)]. See also age-specific recommendations [see *Dosage and Administration* (2.2, 2.3)]. In selected patients in need of fluid restriction, a concentration up to 10 mg/mL may be used. Infusion related events may occur, however, at any rate or concentration [see *Warnings and Precautions* (5.5)].
- Other drugs should not be added to the diluted Vancomycin Hydrochloride for Injection [see *Dosage and Administration* (2.7)].
- Administer diluted Vancomycin Hydrochloride for Injection prior to intravenous anesthetic agents to reduce the risk of infusion related adverse reactions [see *Warnings and Precautions* (5.5)].
- Administer diluted Vancomycin Hydrochloride for Injection by a secure intravenous route of administration to avoid local irritation and phlebitis reactions [see *Warnings and Precautions* (5.6)].

2.2 Intravenous Dosage in Adult and Pediatric Patients with Normal Renal Function

Dosage in Adult Patients

The usual daily intravenous dose is 2 grams (g) divided either as 500 mg every 6 hours or 1 g every 12 hours. Administer each dose by intravenous infusion over a period of 60 minutes or greater. Other patient factors, such as age or obesity, may call for modification of the usual intravenous daily dose.

Dosage in Pediatric Patients Aged 1 Month and Older

The usual intravenous dosage of vancomycin is 10 mg/kg per dose given every 6 hours. Each dose should be administered over a period of at least 60 minutes. Close monitoring of serum concentrations of vancomycin may be warranted in these patients.

Dosage in Pediatric Patients Younger than 1 Month of Age

In pediatric patients, up to the age of 1 month, the total daily intravenous dosage may be lower. In neonates, an initial dose of 15 mg/kg is suggested, followed by 10 mg/kg every 12 hours for pediatric patients in the 1st week of life and every 8 hours thereafter up to the age of 1 month. Each dose should be administered over 60 minutes. In premature infants, vancomycin clearance decreases as postconceptional age decreases. Therefore, longer dosing intervals may be necessary in premature infants. Close monitoring of serum concentrations of vancomycin is recommended in these patients.

2.3 Orally Administered Dosage in Adult and Pediatric Patients

Dosage in Adult Patients

- *C. difficile*-associated diarrhea: The recommended dose is 125 mg administered orally 4 times daily for 10 days.
- Staphylococcal enterocolitis: Total daily dosage is 500 mg to 2 g administered orally in 3 or 4 divided doses for 7 days to 10 days.

Dosage in Pediatric Patients (Less than 18 years of age)

For both *C. difficile*-associated diarrhea and staphylococcal enterocolitis, the usual daily dosage is 40 mg/kg in 3 or 4 divided doses for 7 days to 10 days. The total daily dosage should not exceed 2 g.

2.4 Intravenous Dosage in Patients with Renal Impairment

Dosage adjustment must be made in adult and pediatric patients with renal impairment. The initial dose should be no less than 15 mg/kg, in adult patients with any degree of renal impairment.

In premature infants and the elderly, greater dosage reductions than expected may be necessary because of decreased renal function. Measure trough vancomycin serum concentrations to guide therapy, especially in seriously ill patients with changing renal function.

For functionally anephric patients, an initial dose of 15 mg/kg of body weight should be given to achieve prompt therapeutic serum concentration. A dose of 1.9 mg/kg/24 hr should be given after the initial dose of 15 mg/kg.

2.5 Directions for the Preparation, Dilution, and Storage of the Single-Dose Vials for Intravenous Use

Preparation

Vancomycin Hydrochloride for Injection must be **reconstituted and further diluted**. No preservative is present in this product. Aseptic technique must be used in the preparation of final IV solution.

Reconstitution of the Spray-dried Powder. At the time of use, reconstitute the vials of Vancomycin Hydrochloride for Injection with Sterile Water for Injection to a concentration of 50 mg of vancomycin/mL (see following table for volume of diluent).

Vancomycin Strength per Vial	Volume of Diluent
500 mg	10 mL
1 g	20 mL

After reconstitution, the vials may be stored at 2°C to 8°C (36°F to 46°F) for 96 hours without significant loss of potency.

Dilution

Reconstituted solutions of vancomycin (50 mg/mL) must be further diluted. A dose of 500 mg (10 mL) must be diluted with at least 100 mL of a suitable infusion solution. For doses of 1 gram (20 mL), at least 200 mL of solution must be used. The desired dose diluted in this manner should be administered by intermittent IV infusion over a period of at least 60 minutes.

Parenteral drug products should be visually inspected for particulate matter and discoloration prior to administration, whenever solution and container permit.

2.6 Compatibility with Intravenous Fluids for Intravenous Use

The following diluents are physically and chemically compatible with approximately 4.5 mg/mL of vancomycin hydrochloride:

5% Dextrose Injection, USP

5% Dextrose Injection and 0.9% Sodium Chloride Injection, USP

Lactated Ringer's Injection, USP

5% Dextrose and Lactated Ringer's Injection, USP

0.9% Sodium Chloride Injection, USP

2.7 Incompatibilities for Intravenous Use

Vancomycin Hydrochloride for Injection reconstituted solution has a low pH and may cause chemical or physical instability when it is mixed with other compounds.

Mixtures of solutions of vancomycin and beta-lactam antibacterial drugs have been shown to be physically incompatible. The likelihood of precipitation increases with higher concentrations of vancomycin. It is recommended to adequately flush the intravenous lines between the administration of these antibacterial drugs. It is also recommended to dilute solutions of vancomycin to 5 mg/mL or less.

2.8 Preparation of Vancomycin Hydrochloride for Injection for Oral Administration

Preparation of the Single-Dose Vial for Oral Use

1. Reconstitute Vancomycin Hydrochloride for Injection to 100 mg/mL by adding the appropriate amount of

sterile water for injection:

Table 1: Reconstitution-Volume of Sterile Water for Injection to be Added	
Vancomycin Hydrochloride for Injection Strength per Vial	Volume of Sterile Water for Injection to be added to achieve 100 mg/mL Vancomycin Hydrochloride Solution
500 mg	5 mL
1 g	10 mL

- Sweeteners may be added to the solution to improve the taste for oral administration following the steps outlined in *Instructions for How to Prepare and Add Sweeteners* below.

Instructions for How to Prepare and Add Sweeteners

- Prepare the sweetening agent separately from the reconstituted Vancomycin hydrochloride solution in a plastic container following one of the preparation options below:
 - Mix sucrose and sterile water for injection in a 1:2 ratio (e.g., 10 g sucrose and 20 mL sterile water for injection) OR
 - Mix agave nectar with an equal volume of sterile water for injection
- Mix the sweetening agent with the reconstituted Vancomycin hydrochloride solution in a 3:1 ratio (e.g., 30 mL sweetening agent with 10 mL vancomycin solution) to obtain a sweetened solution containing 25 mg/mL of vancomycin that is suitable for oral administration.

The diluted solution may be administered orally or via a nasogastric tube [see *Dosage and Administration* (2.3)].

2.9 Storage of Reconstituted and Diluted Vancomycin Hydrochloride for Injection for Intravenous or Oral Administration

Intravenous Administration:

Administer the compounded mixtures within **4** hours after preparation when storing at room temperature and 6 days when stored refrigerated (14 days for solutions made with 0.9% sodium chloride and 5% dextrose).

Oral Administration:

Store the diluted solution for oral use in a plastic container refrigerated at 2 °C to 8 °C (36°F to 46°F) for up to 10 days.

3 DOSAGE FORMS AND STRENGTHS

Vancomycin Hydrochloride for Injection is a sterile white, almost white to tan to brown, spray-dried powder in single-dose vials each containing vancomycin hydrochloride equivalent to 500 mg or 1 g vancomycin base.

4 CONTRAINDICATIONS

Vancomycin Hydrochloride for Injection is contraindicated in patients with known hypersensitivity to vancomycin and polysorbate 80.

5 WARNINGS AND PRECAUTIONS

5.1 Nephrotoxicity

Vancomycin Hydrochloride for Injection administered intravenously or orally can result in acute kidney injury (AKI), including acute renal failure, mainly due to interstitial nephritis or less commonly acute tubular necrosis. AKI is manifested by increasing blood urea nitrogen (BUN) and serum creatinine (Cr). The risk of AKI increases with higher vancomycin serum levels, prolonged exposure, concomitant administration of other nephrotoxic drugs, concomitant administration of piperacillin-tazobactam [see *Drug Interactions* (7.2)], volume depletion, pre-existing renal impairment and in critically ill patients and patients with co-morbid conditions that predispose to renal impairment.

Monitor serum vancomycin concentrations and renal function in all patients receiving Vancomycin Hydrochloride for Injection intravenously.

Nephrotoxicity (e.g., reports of renal failure, renal impairment, blood creatine increased) has occurred following oral vancomycin hydrochloride capsule therapy in randomized controlled clinical studies and can occur either during or after completion of therapy. The risk of nephrotoxicity is increased in patients >65 years of age [see *Adverse Reactions (6.1) and Use in Specific Populations (8.5)*]. In patients over 65 years of age, including those with normal renal function prior to treatment, renal function should be monitored during and following treatment with oral Vancomycin Hydrochloride for Injection therapy to detect potential vancomycin induced nephrotoxicity.

More frequent monitoring is recommended in patients with comorbidities that predispose to impairment in renal function or are concomitantly receiving other nephrotoxic drugs, in critically ill patients, in patients with changing renal function, and in patients requiring higher therapeutic vancomycin levels. If acute kidney injury occurs, discontinue Vancomycin Hydrochloride for Injection or reduce the dose.

5.2 Ototoxicity

Ototoxicity has occurred in patients administered vancomycin intravenously or orally. It may be transient or permanent. Ototoxicity manifests as tinnitus, hearing loss, dizziness or vertigo. The risk is higher in older patients, patients who are receiving higher doses, who have an underlying hearing loss, who are receiving concomitant therapy with another ototoxic agent, such as an aminoglycoside or who have underlying renal impairment. Monitor for signs and symptoms of ototoxicity during oral or intravenous therapy with Vancomycin Hydrochloride for Injection. Discontinue Vancomycin Hydrochloride for Injection if ototoxicity occurs. Serial tests of auditory function may be helpful in order to minimize the risk of ototoxicity.

5.3 Severe Dermatologic Reactions

Severe dermatologic reactions such as toxic epidermal necrolysis (TEN), Stevens-Johnson syndrome (SJS), drug reaction with eosinophilia and systemic symptoms (DRESS), acute generalized exanthematous pustulosis (AGEP), and linear IgA bullous dermatosis (LABD) have been reported in association with the use of vancomycin. Cutaneous signs or symptoms reported include skin rashes, mucosal lesions, and blisters.

Discontinue Vancomycin Hydrochloride for Injection at the first appearance of signs and symptoms of TEN, SJS, DRESS, AGEP, or LABD.

5.4 Neutropenia

Reversible neutropenia has been reported in patients administered vancomycin intravenously or orally. Patients who will undergo prolonged therapy with Vancomycin Hydrochloride for Injection or those who are receiving concomitant drugs which may cause neutropenia should have periodic monitoring of the leukocyte count.

5.5 Infusion Reactions

Hypotension, including shock and cardiac arrest, wheezing, dyspnea, urticaria, muscular and chest pain may occur with rapid Vancomycin Hydrochloride for Injection intravenous administration. The reactions may be more severe in younger patients, particularly children, and in patients receiving concomitant muscle relaxant anesthetics. Rapid intravenous administration of Vancomycin Hydrochloride for Injection may also be associated with vancomycin infusion reaction, which manifests as pruritus and erythema that involves the face, neck and upper torso. Infusion-related adverse reactions are related to both the concentration and the rate of administration of Vancomycin Hydrochloride for Injection. Infusion-related adverse reactions may occur, however, at any rate or concentration.

Administer Vancomycin Hydrochloride for Injection in a diluted solution over a period of 60 minutes or greater to reduce the risk of infusion-related adverse reactions. In selected patients in need of fluid restriction, a concentration up to 10 mg/mL may be used; use of such higher concentrations may increase the risk of infusion-related adverse reactions [see *Nonclinical Toxicology (13.2)*]. Administer prior to intravenous anesthetic agents when feasible. Stop the infusion if a reaction occurs.

5.6 Phlebitis and Adverse Reactions with Unapproved Routes of Administration

Inflammation at the site of injection of vancomycin has been reported. Vancomycin is irritating to tissue and must be given by a secure intravenous route of administration to reduce the risk of local irritation and phlebitis. Thrombophlebitis may occur, the frequency and severity of which can be minimized by slow infusion of the drug

and by rotation of venous access sites.

Administration of Vancomycin Hydrochloride for Injection by intramuscular (IM), intraperitoneal, intrathecal intraventricular, or intravitreal routes has not been approved and is not recommended. The safety and efficacy of vancomycin administered by these routes of administration have not been established by adequate and well-controlled trials. Pain, tenderness, and necrosis occur with IM injection of vancomycin or with inadvertent extravasation. Intraperitoneal administration during continuous ambulatory peritoneal dialysis (CAPD) can result in chemical peritonitis. Manifestations range from cloudy dialysate alone to a cloudy dialysate accompanied by variable degrees of abdominal pain and fever. This syndrome appears to be resolved after discontinuation of intraperitoneal vancomycin.

About 60% of an intraperitoneal dose of vancomycin administered during peritoneal dialysis is absorbed systemically in 6 hours. Serum concentrations of about 10 mcg/mL are achieved by intraperitoneal injection of 30 mg/kg of vancomycin. However, the safety and efficacy of the intraperitoneal use of vancomycin has not been established in adequate and well-controlled trials.

5.7 *Clostridioides difficile*-Associated Diarrhea (CDAD)

Clostridioides difficile-associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including with intravenous administration of vancomycin and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibacterial use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents. If CDAD is suspected or confirmed, ongoing antibacterial use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibacterial treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated. Clinically significant serum concentrations of vancomycin have been reported in some patients being treated for active *C. difficile*-induced pseudomembranous colitis after multiple oral doses of vancomycin [see *Warnings and Precautions* (5.10)].

Prolonged use of Vancomycin Hydrochloride for Injection may result in the overgrowth of non-susceptible microorganisms. Careful observation of the patient is essential. If superinfection occurs during therapy, appropriate measures should be taken. There have been reports of pseudomembranous colitis due to *C. difficile* developing in patients who received intravenous vancomycin.

5.8 Hemorrhagic Occlusive Retinal Vasculitis (HORV)

Hemorrhagic occlusive retinal vasculitis, including permanent loss of vision, occurred in patients receiving intracameral or intravitreal administration of vancomycin during or after cataract surgery. The safety and efficacy of vancomycin administered by the intracameral or the intravitreal route have not been established by adequate and well-controlled trials. Vancomycin Hydrochloride for Injection (intravenously and orally administered) is not indicated for the prophylaxis of endophthalmitis.

5.9 Development of Drug-Resistant Bacteria

Prescribing Vancomycin Hydrochloride for Injection (intravenously and orally administered) in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

5.10 Potential for Systemic Absorption after Oral Administration

Clinically significant serum concentrations have been reported in some patients who have taken multiple oral doses of oral vancomycin for active *C. difficile*-associated diarrhea. Some patients with inflammatory disorders of the intestinal mucosa also may have significant systemic absorption of vancomycin. These patients may be at risk for the development of adverse reactions associated with higher doses of oral vancomycin; therefore, monitoring of serum concentrations of vancomycin may be appropriate in some instances, e.g., in patients with renal insufficiency and/or colitis or in those receiving concomitant therapy with an aminoglycoside antibacterial drug.

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Nephrotoxicity [see *Warnings and Precautions* (5.1)]
- Ototoxicity [see *Warnings and Precautions* (5.2)]
- Severe Dermatologic Reactions [see *Warnings and Precautions* (5.3)]
- Neutropenia [see *Warnings and Precautions* (5.4)]
- Infusion Reactions [see *Warnings and Precautions* (5.5)]
- *Clostridioides difficile*-Associated Diarrhea [see *Warnings and Precautions* (5.7)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse Reactions in Patients Receiving Intravenously and Orally Administered Vancomycin

The following adverse reactions associated with the use of intravenously and orally administered vancomycin were identified in clinical trials:

Renal and urinary disorders: Acute kidney injury and interstitial nephritis [see *Warnings and Precautions* (5.1)]

Ear and labyrinth disorders: Tinnitus, hearing loss, vertigo [see *Warnings and Precautions* (5.2)]

Skin and subcutaneous tissue disorders: Erythema (especially of the face, neck and upper torso) and pruritus which are manifestations of rashes including exfoliative dermatitis, toxic epidermal necrolysis (TEN), Stevens-Johnson syndrome (SJS), Linear IgA bullous dermatosis (LABD) [see *Warnings and Precautions* (5.3)].

Blood and lymphatic system disorders: Agranulocytosis, neutropenia, pancytopenia, leukopenia, thrombocytopenia, eosinophilia [see *Warnings and Precautions* (5.4)]

Immune system disorders: Hypersensitivity reactions including anaphylaxis and vancomycin infusion reaction [see *Warnings and Precautions* (5.5)]

General disorders and administration site conditions: General discomfort, fever, chills, phlebitis, injection site irritation, injection site pain and necrosis following intramuscular injection, chemical peritonitis following intraperitoneal administration. Vancomycin Hydrochloride for Injection is not approved for intramuscular and intraperitoneal administration [see *Warnings and Precautions* (5.6)]

Gastrointestinal disorders: Pseudomembranous colitis [see *Warnings and Precautions* (5.7)]

Cardiac disorders: Cardiac arrest, chest pain

Laboratory abnormalities: Elevated blood urea nitrogen, elevated serum creatinine

Musculoskeletal and connective tissue disorders: Muscle pain

Nervous system disorders: Dizziness

Respiratory, thoracic and mediastinal disorders: Wheezing, dyspnea

Vascular disorders: Hypotension, shock, vasculitis

Adverse Reactions in Patients Receiving Oral Administration of Vancomycin Hydrochloride Capsules

The data described below reflect exposure to vancomycin hydrochloride capsules in 260 adult subjects in two Phase 3 clinical trials for the treatment of diarrhea associated with *C. difficile*. In both trials, subjects received vancomycin hydrochloride capsules 125 mg orally four times daily. The mean duration of treatment was 9.4 days. The median age of patients was 67, ranging between 19 and 96 years of age. Patients were predominantly Caucasian (93%) and 52% were male.

Adverse reactions occurring in $\geq 5\%$ of vancomycin hydrochloride capsules-treated subjects are shown in Table 2. The most common adverse reactions associated with vancomycin hydrochloride capsules ($\geq 10\%$) were nausea, abdominal pain, and hypokalemia.

Table 2: Common ($\geq 5\%$) Adverse Reactions^a for Vancomycin Hydrochloride Capsules Reported in

Clinical Trials for Treatment of Diarrhea Associated with *C. difficile*

System/Organ Class	Adverse Reaction	Vancomycin Hydrochloride Capsule % (N=260)
Gastrointestinal disorders	Nausea	17
	Abdominal pain	15
	Vomiting	9
	Diarrhea	9
	Flatulence	8
General disorders and administration site conditions	Pyrexia	9
	Edema peripheral	6
	Fatigue	5
Infections and infestations	Urinary tract infection	8
Metabolism and nutrition disorders	Hypokalemia	13
Musculoskeletal and connective tissue disorders	Back pain	6
Nervous system disorders	Headache	7
^a Adverse reaction rates were derived from the incidence of treatment-emergent adverse events.		

Nephrotoxicity (e.g., reports of renal failure, renal impairment, blood creatinine increased) occurred in 5% of subjects treated with vancomycin hydrochloride capsules. Nephrotoxicity following vancomycin hydrochloride capsules typically first occurred within one week after completion of treatment (median day of onset was Day 16). Nephrotoxicity following vancomycin hydrochloride capsules occurred in 6% of subjects >65 years of age and 3% of subjects ≤65 years of age.

The incidences of hypokalemia, urinary tract infection, peripheral edema, insomnia, constipation, anemia, depression, vomiting, and hypotension were higher among subjects >65 years of age than in subjects ≤65 years of age.

Discontinuation of study drug due to adverse events occurred in 7% of subjects treated with vancomycin hydrochloride capsules. The most common adverse events leading to discontinuation of vancomycin hydrochloride capsules were *C. difficile* colitis (<1%), nausea (<1%), and vomiting (<1%).

6.2 Postmarketing Experience

The following adverse reactions have been identified during postmarketing use of vancomycin (administered orally and intravenously). Because these reactions are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Skin and Subcutaneous Tissue Disorders: Drug reaction with eosinophilia and systemic symptoms (DRESS), acute generalized exanthematous pustulosis (AGEP) [see *Warnings and Precautions* (5.3)].

Ototoxicity: Cases of hearing loss associated with intravenously administered vancomycin have been reported. Most of these patients had kidney dysfunction or a preexisting hearing loss or were receiving concomitant treatment with an ototoxic drug [see *Warnings and Precautions* (5.2)]. Vertigo, dizziness, and tinnitus have been reported.

Hematopoietic: Reversible neutropenia, usually starting 1 week or more after onset of intravenous therapy with vancomycin or after a total dose of more than 25 g, has been reported for several dozen patients. Neutropenia appears to be promptly reversible when vancomycin is discontinued. Thrombocytopenia has been reported.

Miscellaneous: Patients have been reported to have had anaphylaxis, drug fever, chills, nausea, eosinophilia, and cases of vasculitis in association with the administration of vancomycin.

A condition has been reported with oral vancomycin that is similar to the intravenous vancomycin-induced syndrome with symptoms consistent with anaphylactoid reactions, including hypotension, wheezing, dyspnea, urticaria, pruritus, flushing of the upper body ("vancomycin infusion reaction"), pain and muscle spasm of the chest and back [see *Warnings and Precautions* (5.5)]. These reactions usually resolve within 20 minutes but may persist for several hours.

7 DRUG INTERACTIONS

7.1 Anesthetic Agents

Concomitant administration of vancomycin and anesthetic agents has been associated with erythema and histamine-like flushing [see *Warnings and Precautions (5.5) and Use in Specific Populations (8.4)*].

7.2 Piperacillin-Tazobactam

Studies have detected an increased incidence of acute kidney injury in patients administered concomitant piperacillin/tazobactam and vancomycin as compared to vancomycin alone. Monitor kidney function in patients receiving concomitant piperacillin/tazobactam and Vancomycin Hydrochloride for Injection. No pharmacokinetic interactions have been noted between piperacillin/tazobactam and vancomycin.

7.3 Ototoxic and/or Nephrotoxic Drugs

Concurrent and/or sequential systemic or topical use of other potentially, neurotoxic and/or nephrotoxic drugs requires more frequent monitoring of renal function.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no available data on vancomycin use in pregnant women to inform a drug associated risk of major birth defects or miscarriage. Available published data on vancomycin use in pregnancy during the second and third trimesters have not shown an association with adverse pregnancy related outcomes (*see Data*). Vancomycin did not show adverse developmental effects when administered intravenously to pregnant rats and rabbits during organogenesis at doses less than or equal to the recommended maximum human dose (*see Data*). Systemic absorption of vancomycin is low following oral administration of vancomycin hydrochloride capsules; however, absorption may vary depending on various factors [see *Clinical Pharmacology (12.3)*].

The estimated background risk of major birth defects and miscarriage for the indicated population(s) is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Data

Human Data

A published study evaluated hearing loss and nephrotoxicity in infants of 10 pregnant intravenous drug users treated with intravenously administered vancomycin for suspected or documented methicillin-resistant *Staphylococcus-aureus* (MRSA) infections in the second or third trimester. The comparison groups were 10 non-intravenous drug-dependent patients who received no treatment, and 10 untreated intravenous drug-dependent patients who served as substance abuse controls. No infant in the vancomycin exposed group had abnormal sensorineural hearing at 3 months of age or nephrotoxicity.

A published prospective study assessed outcomes in 55 pregnant women with a positive Group B streptococcus (GBS) culture and a high-risk penicillin allergy with resistance to clindamycin or unknown sensitivity who were administered vancomycin intravenously at the time of delivery. Vancomycin dosing ranged from the standard 1 g intravenously every 12 hours to 20 mg/kg intravenous every 8 hours (maximum individual dose 2 g). No major adverse reactions were recorded either in the mothers or their newborns. None of the newborns had sensorineural hearing loss. Neonatal renal function was not examined, but all of the newborns were discharged in good condition.

Animal Data

Vancomycin did not cause fetal malformations when administered during organogenesis to pregnant rats (gestation days 6 to 15) and rabbits (gestation days 6 to 18) at the equivalent recommended maximum human dose (based on body surface area comparisons) of 200 mg/kg/day IV to rats or 120 mg/kg/day IV to rabbits¹. No effects on fetal weight or development were seen in rats at the highest dose tested or in rabbits given 80

mg/kg/day (approximately 1 and 0.8 times the recommended maximum human dose based on body surface area, respectively). Maternal toxicity was observed in rats (at doses 120 mg/kg and above) and rabbits (at 80 mg/kg and above). There were no oral embryo-fetal toxicity studies conducted in animals with this Vancomycin Injection product.

8.2 Lactation

Risk Summary

Vancomycin is present in human milk following intravenous administration, however, there are insufficient data to inform the levels. There are no available data on vancomycin presence in milk following oral administration. Systemic absorption of vancomycin is low following oral administration of vancomycin hydrochloride capsules; however, absorption may vary depending on various factors [see *Clinical Pharmacology* (12.3)]. There are no data on the effects of vancomycin on the breastfed infant or milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Vancomycin Hydrochloride for Injection and any potential adverse effects on the breastfed infant from Vancomycin Hydrochloride for Injection or from the underlying maternal condition.

8.4 Pediatric Use

Vancomycin Hydrochloride for Injection administered intravenously is indicated in pediatric patients less than 18 years of age for the treatment of septicemia, infective endocarditis, skin and skin structure infections, bone infections and lower respiratory tract infections [see *Indications and Usage* (1.1 to 1.5) and *Dosage and Administration* (2.2)].

Vancomycin Hydrochloride for Injection administered orally is indicated in pediatric patients less than 18 years of age for the treatment of *Clostridioides difficile*-associated diarrhea and enterocolitis caused by *Staphylococcus aureus* (including methicillin-resistant strains) [see *Indications and Usage* (1.6, 1.7) and *Dosage and Administration* (2.3)].

In pediatric patients, monitor vancomycin serum concentration and renal function when administering Vancomycin Hydrochloride for Injection intravenously [see *Dosage and Administration* (2.2) and *Warnings and Precautions* (5.1)]. More severe infusion related reactions related to Vancomycin Hydrochloride for Injection administration may occur in pediatric patients. Concomitant administration of vancomycin and intravenous anesthetic agents has been associated with erythema and histamine-like flushing in all patients including pediatric patients [see *Warnings and Precautions* (5.5)].

8.5 Geriatric Use

Clinical studies with another vancomycin hydrochloride drug product in *C. difficile*-associated diarrhea have demonstrated that geriatric subjects are at increased risk of developing nephrotoxicity following treatment with oral vancomycin hydrochloride, which may occur during or after completion of therapy. In patients over 65 years of age, including those with normal renal function prior to treatment, renal function should be monitored during and following treatment with Vancomycin Hydrochloride for Injection to detect potential vancomycin induced nephrotoxicity [see *Warnings and Precautions* (5.1)] and *Adverse Reactions* (6.1)].

Patients over 65 years of age may take longer to respond to therapy compared to patients 65 years of age and younger. Clinicians should be aware of the importance of appropriate duration of Vancomycin Hydrochloride for Injection treatment in patients over 65 years of age and not discontinue or switch to alternative treatment prematurely.

Vancomycin Hydrochloride for Injection is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection [see *Dosage and Administration* (2.4)], and it may be useful to monitor renal function [see *Warnings and Precautions* (5.1)].

8.6 Renal Impairment

Dosage adjustment must be made in patients with impaired renal function [see *Dosage and Administration* (2.3)]. Measure trough vancomycin serum concentrations to guide intravenous therapy, especially in seriously ill patients with changing renal function.


10 OVERDOSAGE

Supportive care is advised, with maintenance of glomerular filtration. Vancomycin is poorly removed by dialysis. Hemofiltration and hemoperfusion with polysulfone resin have been reported to result in increased vancomycin clearance.

For current information on the management of overdose, contact the National Poison Control Center at 1-800-222-1222 or www.poisson.org.

11 DESCRIPTION

Vancomycin Hydrochloride for Injection contains the hydrochloride salt of vancomycin, a tricyclic glycopeptide antibacterial derived from *Amycolatopsis orientalis* (formerly *Nocardia orientalis*). The chemical name for vancomycin hydrochloride is (Sa)-(3S,6R,7R,22R,23S, 26S,36R,38aR)-44-[[2-O-(3-Amino-2,3,6-trideoxy-3-C-methyl- α -L-lyxo-hexopyranosyl)- β -D-glucopyranosyl]oxy]-3-(carbamoylmethyl)-10,19-dichloro-2,3,4,5,6,7,23,24,25,26,36,37,38,38a-tetradecahydro-7,22,28,30,32-pentahydroxy-6-[(2R)-4-methyl-2-(methylamino)valeramido]2,5,24,38,39-pentaoxo-22H-8,11:18,21-dietheno-23,36-(iminomethano)-13,16:31,35-dimetheno-1H,16H-[1,6,9]oxadiazacyclohexadecino[4,5-m][10,2,16]-benzoxadiazacyclotetracosine-26-carboxylic acid, monohydrochloride. The molecular formula is $C_{66}H_{75}Cl_2N_9O_{24} \cdot HCl$ and the molecular weight is 1,485.71. Vancomycin hydrochloride has the following structural formula:



Vancomycin Hydrochloride for Injection is a sterile, preservative-free, white, almost white to tan to brown, spray-dried powder, for preparing intravenous (IV) infusions, in single-dose vials each containing the equivalent of 500 mg or 1 g vancomycin base. Each 500 mg vial contains 500 mg vancomycin (equivalent to 513 mg vancomycin hydrochloride), 125 mg trehalose, 0.05 mg polysorbate 80, and hydrochloric acid used to adjust the pH. Each 1 g vial contains 1 g vancomycin (equivalent to 1.025 g vancomycin hydrochloride), 250 mg trehalose, 0.1 mg polysorbate 80, and hydrochloric acid used to adjust the pH.

The spray-dried powder is reconstituted with Sterile Water for Injection which forms a clear solution and subsequently diluted prior to intravenous or oral administration [see *Dosage and Administration* (2.5, 2.8)]. Vancomycin Hydrochloride for Injection may also be diluted with a sweetening agent for oral administration [see *Dosage and Administration* (2.8)].

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Vancomycin is an antibacterial drug [see *Microbiology* (12.4)].

12.2 Pharmacodynamics

The pharmacodynamics of vancomycin is unknown.

12.3 Pharmacokinetics

General Pharmacokinetics

In subjects with normal kidney function, multiple intravenous dosing of 1 g of vancomycin (15 mg/kg) infused over 60 minutes produces mean plasma concentrations of approximately 63 mcg/mL immediately after the completion of infusion, mean plasma concentrations of approximately 23 mcg/mL 2 hours after infusion, and mean plasma concentrations of approximately 8 mcg/mL 11 hours after the end of the infusion. Multiple dosing of 500 mg infused over 30 minutes produces mean plasma concentrations of about 49 mcg/mL at the completion of infusion, mean plasma concentrations of about 19 mcg/mL 2 hours after infusion, and mean plasma concentrations of about 10 mcg/mL 6 hours after infusion. The plasma concentrations during multiple dosing are like those after a single dose.

Absorption

Vancomycin is poorly absorbed after oral administration. During multiple dosing of 250 mg every 8 hours for 7 doses, fecal concentrations of vancomycin in volunteers exceeded 100 mg/kg in the majority of samples. No blood concentrations were detected, and urinary recovery did not exceed 0.76%. In anephric subjects with no inflammatory bowel disease who received vancomycin oral solution 2 g for 16 days, blood concentrations of vancomycin were less than or equal to 0.66 mcg/mL in 2 of 5 subjects. No measurable blood concentrations were attained in the other 3 subjects. Following doses of 2 g daily, concentrations of drug were >3100 mg/kg in the feces and <1 mcg/mL in the serum of subjects with normal renal function who had *C. difficile*-associated diarrhea. After multiple-dose oral administration of vancomycin, measurable serum concentrations may occur in patients with active *C. difficile*-associated diarrhea, and, in the presence of renal impairment, the possibility of accumulation exists. It should be noted that the total systemic and renal clearances of vancomycin are reduced in the elderly [see *Use in Specific Populations* (8.5)].

Distribution

The volume of distribution ranges from 0.3 to 0.43 L/kg after intravenous administration. Vancomycin is approximately 55% serum protein bound as measured by ultrafiltration at vancomycin serum concentrations of 10 to 100 mcg/mL. After intravenous administration of vancomycin, inhibitory concentrations are present in pleural, pericardial, ascitic, and synovial fluids; in urine; in peritoneal dialysis fluid; and in atrial appendage tissue. Vancomycin does not readily diffuse across normal meninges into the spinal fluid; but, when the meninges are inflamed, penetration into the spinal fluid occurs.

Elimination

Mean plasma clearance is about 0.058 L/kg/h, and mean renal clearance is about 0.048 L/kg/h. The mean elimination half-life of vancomycin from plasma is 4 to 6 hours in subjects with normal renal function. In anephric patients, the mean elimination half-life is 7.5 days. Total body and renal clearance of vancomycin may be reduced in the elderly.

Metabolism

There is no *apparent* metabolism of vancomycin.

Excretion

In the first 24 hours after intravenous administration, about 75% of an administered dose of vancomycin is excreted in urine by glomerular filtration. Renal impairment slows excretion of vancomycin.

12.4 Microbiology

Mechanism of Action

The bactericidal action of vancomycin results primarily from inhibition of cell-wall biosynthesis. In addition, vancomycin alters bacterial-cell-membrane permeability and RNA synthesis.

Resistance

Vancomycin is not active *in vitro* against gram-negative bacilli, mycobacteria, or fungi. There is no cross-resistance between vancomycin and other antibacterials.

C. difficile

Isolates of *C. difficile* generally have vancomycin minimal inhibitory concentrations (MICs) of <1 mcg/mL; however, vancomycin MICs ranging from 4 mcg/mL to 16 mcg/mL have been reported. The mechanism which mediates *C. difficile*'s decreased susceptibility to vancomycin has not been fully elucidated.

S. aureus

S. aureus isolates with vancomycin MICs as high as 1024 mcg/mL have been reported. The exact mechanism of this resistance is not clear but is believed to be due to cell wall thickening and potentially the transfer of genetic material.

Interaction with Other Antimicrobials

The combination of vancomycin and an aminoglycoside acts synergistically *in vitro* against many isolates of *Staphylococcus aureus*, *Streptococcus gallolyticus* (previously known as *Streptococcus bovis*), *Enterococcus* species, and the viridans group streptococci.

Antimicrobial Activity

Vancomycin has been shown to be active against most isolates of the following microorganisms, both *in vitro* and in clinical infections [see *Indications and Usage (1)*].

Aerobic bacteria

Gram-positive bacteria

Corynebacterium species

Enterococcus species (including *Enterococcus faecalis*)

Staphylococcus aureus (including methicillin-resistant and methicillin-susceptible isolates)

Coagulase negative staphylococci (including *S. epidermidis* and methicillin-resistant isolates)

Streptococcus gallolyticus (previously known as *Streptococcus bovis*)

Viridans group streptococci

Anaerobic bacteria

Gram-positive bacteria

Clostridioides difficile isolates associated with *C. difficile* associated diarrhea.

The following *in vitro* data are available, but their clinical significance is unknown. At least 90 percent of the following bacteria exhibit an *in vitro* minimum inhibitory concentration (MIC) less than or equal to the susceptible breakpoint for vancomycin against isolates of similar genus or organism group. However, the efficacy of vancomycin in treating clinical infections caused by these bacteria has not been established in adequate and well-controlled clinical trials.

Aerobic bacteria

Gram-positive bacteria

Listeria monocytogenes

Streptococcus pyogenes

Streptococcus pneumoniae

Streptococcus agalactiae

Anaerobic bacteria

Gram-positive bacteria

Actinomyces species

Lactobacillus species

Susceptibility Testing

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term carcinogenesis studies in animals have been conducted.

At concentrations up to 1000 mcg/mL, vancomycin had no mutagenic effect *in vitro* in the mouse lymphoma forward mutation assay or the primary rat hepatocyte unscheduled DNA synthesis assay. The concentrations tested *in vitro* were above the peak plasma vancomycin concentrations of 20 to 40 mcg/mL usually achieved in humans after slow infusion of the maximum recommended dose of 1 g. Vancomycin had no mutagenic effect *in vivo* in the Chinese hamster sister chromatid exchange assay (400 mg/kg IP) or the mouse micronucleus assay (800 mg/kg IP).

No definitive fertility studies have been conducted.

13.2 Animal Toxicology and/or Pharmacology

In animal studies, hypotension and bradycardia occurred in dogs receiving an intravenous infusion of vancomycin hydrochloride 25 mg/kg, at a concentration and infusion rate higher than recommended (concentration of 25 mg/mL and an infusion rate of 13.3 mL/min).

In a 4-week repeat-dose toxicokinetic study, electrocardiograms were obtained from dogs receiving Vancomycin Hydrochloride for Injection by the IV route, twice daily at 33 mg/kg (at a concentration of 10 mg/mL, over 30 minutes), for a total daily vancomycin dose of 66 mg/kg/day. No bradycardia was observed in the dogs when the pretest heart rates were compared to the heart rates observed 1 to 2 hours after the first daily dose during week 4.

15 REFERENCES

1. Byrd RA., Gries CL, Buening M.: Developmental Toxicology Studies of Vancomycin Hydrochloride Administered Intravenously to Rats and Rabbits. *Fundam Appl Toxicol* 1994; 23: 590-597.

16 HOW SUPPLIED/STORAGE AND HANDLING

How supplied

Vancomycin Hydrochloride for Injection is a white, almost white to tan to brown, sterile spray-dried powder in single-dose vials each containing vancomycin hydrochloride equivalent to 500 mg or 1 g of vancomycin base. The vial stopper is not made with natural rubber latex.

NDC No. Unit of sale	Strength Unit of Sale	NDC for Each Vial
0143-9161-25 Unit of 25	Vancomycin Hydrochloride for Injection, equivalent to 500 mg vancomycin in a flip-top vial, in packages of 25 vials.	0143-9161-01
0143-9162-10 Unit of 10	Vancomycin Hydrochloride for Injection, equivalent to 1 g vancomycin in a flip-top vial in packages of 10 vials.	0143-9162-01

Storage

Store at 20°C to 25°C (68°F to 77°F). Brief exposure to 15°C to 30°C (59°F to 86°F) is permitted [See USP Controlled Room Temperature]. Storage after reconstitution and dilution is described elsewhere in the labeling [see *Dosage and Administration* (2.5, 2.8)].

17 PATIENT COUNSELING INFORMATION

Acute Kidney Injury

Advise patients that Vancomycin Hydrochloride for Injection can result in kidney damage and that blood tests are required to monitor vancomycin blood levels and kidney function during therapy [see *Warnings and Precautions* (5.1)].

Hearing Loss or Balance Problems

Advise patients that Vancomycin Hydrochloride for Injection may result in decreased hearing and to report hearing loss or balance problems to their health care provider [see *Warnings and Precautions* (5.2)].

Severe Dermatologic Reactions

Advise patients about the signs and symptoms of serious skin manifestations. Instruct patients to stop Vancomycin Hydrochloride for Injection immediately and promptly seek medical attention at the first signs or symptoms of skin rash, mucosal lesions or blisters [see *Warnings and Precautions* (5.3)].

Infusion Reactions During or After Intravenous Use

Advise patients that generalized skin redness, skin rash, itching, flushing, muscle pain, chest pain, shortness of breath, wheezing, or dizziness may occur during Vancomycin Hydrochloride for Injection infusion [see *Warnings and Precautions* (5.5)].

Diarrhea

Diarrhea is a common problem caused by antibacterial drugs, including vancomycin, which usually ends when the antibacterial drug is discontinued. Sometimes after starting treatment with antibacterial drugs, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibacterial drug. If this occurs during treatment with Vancomycin Hydrochloride for Injection, patients should contact their physician as soon as possible [see *Warnings and Precautions* (5.7)].

Antibacterial Resistance

Patients should be counseled that antibacterial drugs including Vancomycin Hydrochloride for Injection should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Vancomycin Hydrochloride for Injection is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by Vancomycin Hydrochloride for Injection or other antibacterial drugs in the future.

Manufactured by:

Zhejiang Novus Pharmaceuticals Co., Ltd.

No. 125, Yuezhong Rd, Lihai Subdistrict,

Yuecheng District, Shaoxing, 312366, Zhejiang,

China

53mm

149mm

NDC 0143-9164-01 Rx Only

**Vancomycin
Hydrochloride**
for Injection

Equivalent to

10 g*/Pharmacy Bulk Package

Vancomycin

For intravenous infusion after dilution
For Oral Use, see prescribing information

**PHARMACY
BULK PACKAGE**
NOT For Direct Infusion

**READ INSERT FOR PRECAUTIONS AND DIRECTIONS
BEFORE USE.**

Sterile. Spray dried. Preservative Free

*Each vial contains: Vancomycin hydrochloride
equivalent to 10 g vancomycin.

Recommended Dosage: See prescribing information.

IMPORTANT: Read package insert for **PRECAUTIONS**
and **DIRECTIONS FOR USE.**

Prior to reconstitution, store at 20°C to 25°C
(68°F to 77°F) [See USP Controlled Room Temperature].

After reconstitution, use promptly. (Discard vial
within 4 hours after entry). See package insert for
full information.

DIRECTIONS FOR USE:

Reconstitute with 90 mL of Sterile Water for Injection to
the 10 g vial of dry, sterile vancomycin powder.

The resultant solution will contain vancomycin
100 mg/mL.

**AFTER RECONSTITUTION, FURTHER DILUTION IS
REQUIRED.**

This pharmacy bulk package is for use in a pharmacy
admixtute service.

Dispense aliquots from the vial using a suitable sterile
dispensing set into infusion fluids under a laminar flow
hood using aseptic technique.

**DISCARD VIAL NO LATER THAN 4 HOURS AFTER
INITIAL CLOSURE PUNCTURE.** (See Package Insert).

Vial stoppers are not made with natural rubber latex.

Manufactured by:
Zhejiang Novus Pharmaceuticals Co., Ltd., No. 125, Yuezhong
Rd, Lihai Subdistrict, Yuecheng District, Shaoxing, 312366
Zhejiang, China

BOTTLE ENTRY

DATE:

TIME:

Lot

Exp YYYY-MM-DD

Linear barcode

(01)103XXXXXXX

Rev. 1/2023

Equivalent to 10 g Vancomycin

VANCOMYCIN HYDROCHLORIDE FOR INJECTION

124mm

42mm

NDC 0143-9163-01 Rx Only

Vancomycin Hydrochloride for Injection

Equivalent to

5 g*/Pharmacy Bulk Package

Vancomycin

For Intravenous Infusion after dilution
For Oral Use, see prescribing information

**PHARMACY
BULK PACKAGE**
NOT For Direct Infusion

READ INSERT FOR PRECAUTIONS AND DIRECTIONS BEFORE USE.

Sterile. Spray dried. Preservative Free.

*Each vial contains: Vancomycin hydrochloride equivalent to 5 g vancomycin.

Recommended Dosage: See prescribing information.

IMPORTANT: Read package insert for PRECAUTIONS and DIRECTIONS FOR USE

Prior to reconstitution, store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].

After reconstitution, use promptly. (Discard vial within 4 hours after entry). See package insert for full information.

DIRECTIONS FOR USE:

Reconstitute with 45 mL of Sterile Water for Injection to the 5 g vial of dry sterile vancomycin powder.

The resultant solution will contain vancomycin 100 mg/mL.

AFTER RECONSTITUTION, FURTHER DILUTION IS REQUIRED.

This pharmacy bulk package is for use in a pharmacy admixture service.

Dispense aliquots from the vial using a suitable sterile dispensing set into infusion fluids under a laminar flow hood using aseptic technique.

DISCARD VIAL NO LATER THAN 4 HOURS AFTER INITIAL CLOSURE PUNCTURE. (See Package Insert).

Vial stoppers are not made with natural rubber latex.
Manufactured by
Zhejiang Novus Pharmaceuticals Co., Ltd. No. 125 Yuezhong Rd. Lihai Subdistrict, Yuecheng District, Shaoxing 312366 Zhejiang, China

Rev. 1/2023

Equivalent to 5 g Vancomycin

VANCOMYCIN HYDROCHLORIDE FOR INJECTION

BOTTLE ENTRY

DATE

TIME

Lot

Exp. YYYY-MM-DD

Linear barcode

001103000000000000

32mm

86mm

NDC 0143 9162 01 Rx Only

Vancomycin Hydrochloride

for Injection

Equivalent to

1 g*/vial

Vancomycin

For Oral Use,

see prescribing Information.

For Intravenous Use

After Reconstitution Must Be

Further Diluted Before Use

READ INSERT FOR PRECAUTIONS AND
DIRECTIONS BEFORE USE.

Sterile. Spray dried. Preservative Free.

*Each single-dose vial contains Vancomycin
hydrochloride equivalent to 1 g vancomycin.

Recommended Dosage: See prescribing information.
Reconstitute with 20 mL Sterile Water for Injection. The
resultant solution will contain vancomycin 50 mg/mL.

AFTER RECONSTITUTION FURTHER DILUTION IS
REQUIRED.

Prior to reconstitution, store at 20°C to 25°C
(68°F to 77°F) [See USP Controlled Room

Temperature]. After reconstitution, this vial

may be stored in a refrigerator at 2°C to 8°C (36°F to

46°F) for 96 hours without significant loss of potency.

Single-Dose Vial-Discard Unused Portion.

Manufactured by

Zhejiang Novus Pharmaceuticals Co., Ltd.

No. 125 Yuezhong Rd, Lihai Subdistrict, Yuecheng

District, Shaoxing 312366 Zhejiang, China

Rev. 1/2023

Lot

Exp YYY-MM-DD

Linear barcode

(01)1030XXXXXXX

Reference ID: 5113379

103 mm

57 mm

un varnish area for batch details and 2D
barcode (to be printed online)
20 mm × 50 mm

Vancomycin Hydrochloride for Injection

NDC 0143-9163-01 Rx Only

READ INSERT FOR PRECAUTIONS AND DIRECTIONS BEFORE USE.

DIRECTIONS FOR USE:

Reconstitute with 90 mL of Sterile Water for Injection to the 10 g vial of dry, sterile vancomycin powder.

The resultant solution will contain vancomycin 100 mg/mL.

AFTER RECONSTITUTION, FURTHER DILUTION IS REQUIRED.

This pharmacy bulk package is for use in a pharmacy admixture service.

Dispense aliquots from the vial using a suitable sterile dispensing set into infusion fluids under a laminar flow hood using aseptic technique.

DISCARD VIAL NO LATER THAN 4 HOURS AFTER INITIAL CLOSURE PUNCTURE. (See Package Insert).

Rev. 1/2023

Sterile. Spray dried. **Preservative Free.**

*Each vial contains: Vancomycin hydrochloride equivalent to 10 g vancomycin.

Recommended Dosage: See prescribing information.

Important: Read package insert for PRECAUTIONS and DIRECTIONS FOR USE.

Prior to reconstitution, store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].

After reconstitution, use promptly. (Discard vial within 4 hours after entry). See package insert for full information.

Vial stoppers are not made with natural rubber latex.

Manufactured by:
Zhejiang Novus Pharmaceuticals Co., Ltd.
No. 125, Yuezhong Rd, Lihai Subdistrict, Yuecheng District, Shaoxing, 312366, Zhejiang, China

NDC 0143-9164-01 Rx Only

Vancomycin Hydrochloride for Injection

Equivalent to

10 g*/Pharmacy Bulk Package

Vancomycin


For intravenous infusion after dilution
For Oral Use, see prescribing information


**PHARMACY
BULK PACKAGE
NOT For
Direct Infusion**


Linear barcode

(01)103XXXXXXX

V10







This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

PETER W KIM
01/20/2023 08:19:48 PM