

**CENTER FOR DRUG EVALUATION AND  
RESEARCH**

*APPLICATION NUMBER:*

**219083Orig1s000**

**PROPRIETARY NAME REVIEW(S)**

---

PROPRIETARY NAME REVIEW

Division of Medication Error Prevention and Analysis 2 (DMEPA 2)  
Office of Medication Error Prevention and Risk Management (OMEPRM)  
Office of Surveillance and Epidemiology (OSE)  
Center for Drug Evaluation and Research (CDER)

\*\*\* This document contains proprietary information that cannot be released to the public\*\*\*

---

|                                                           |                                                        |
|-----------------------------------------------------------|--------------------------------------------------------|
| Date of This Review:                                      | December 6, 2024                                       |
| Application Type and Number:                              | NDA 219083                                             |
| Product Name, Dosage Form, and Strength:                  | Myqorzo (aficamten) Tablets, 5 mg, 10 mg, 15 mg, 20 mg |
| Product Type:                                             | Single Ingredient Product                              |
| Rx or OTC:                                                | Prescription (Rx)                                      |
| Applicant/Sponsor Name:                                   | Cytokinetics, Incorporated (Cytokinetics)              |
| PNR ID #:                                                 | 2024-1044726034                                        |
| DMEPA 2 Safety Evaluator:                                 | Julie Neshiewat, PharmD, MS, BCPS, CPH                 |
| DMEPA 2 Team Leader:                                      | Nicole Iverson, PharmD, BCPS                           |
| DMEPA 2 Associate Director for Nomenclature and Labeling: | Hina Mehta, PharmD                                     |

---

## Table of Contents

|       |                                                                                                 |    |
|-------|-------------------------------------------------------------------------------------------------|----|
| 1     | INTRODUCTION .....                                                                              | 3  |
| 1.1   | REGULATORY HISTORY .....                                                                        | 3  |
| 1.2   | PRODUCT INFORMATION .....                                                                       | 3  |
| 2     | DISCUSSION .....                                                                                | 5  |
| 2.1   | MISBRANDING ASSESSMENT .....                                                                    | 5  |
| 2.2   | SAFETY ASSESSMENT .....                                                                         | 5  |
| 2.2.1 | United States Adopted Names (USAN) Search .....                                                 | 5  |
| 2.2.2 | Components of the Proposed Proprietary Name .....                                               | 6  |
| 2.2.3 | Comments from Other Review Disciplines at Initial Review .....                                  | 6  |
| 2.2.4 | FDA Prescription Simulation Studies .....                                                       | 6  |
| 2.2.5 | Phonetic and Orthographic Computer Analysis (POCA) Search Results .....                         | 6  |
| 2.2.6 | Names Retrieved for Review Organized by Name Pair Similarity .....                              | 9  |
| 2.2.7 | Safety Analysis of Names with Potential Orthographic, Spelling, and Phonetic Similarities ..... | 9  |
| 2.2.8 | Communication of DMEPA's Determination .....                                                    | 9  |
| 3     | CONCLUSION .....                                                                                | 9  |
| 3.1   | Comments to Cytokinetics, Incorporated .....                                                    | 9  |
| 4     | REFERENCES .....                                                                                | 10 |
| 5     | APPENDICES .....                                                                                | 11 |

## 1 INTRODUCTION

This review evaluates the proposed proprietary name, Myqorzo, from a safety and misbranding perspective. The sources and methods used to evaluate the proposed proprietary name are outlined in the Reference section and Appendix A, respectively. Cytokinetics submitted an external name study, conducted (b) (4) for this proposed proprietary name. The submitted external name study was previously reviewed under IND 138814 (see Section 1.1 below).

### 1.1 REGULATORY HISTORY

Cytokinetics previously submitted the proposed proprietary name, Myqorzo, under IND 138814 on February 28, 2023, and we found the name conditionally acceptable on July 5, 2023.<sup>b</sup>

### 1.2 PRODUCT INFORMATION

The following product information is provided in the proprietary name submission received on September 26, 2024<sup>c</sup> and the prescribing information received on August 14, 2024<sup>d</sup>.

| Table 1. Relevant Product Information for Myqorzo |                                                                                     |
|---------------------------------------------------|-------------------------------------------------------------------------------------|
| Intended pronunciation:                           | mye kor' zoe                                                                        |
| Active ingredient:                                | aficamten                                                                           |
| Indication:                                       | Treatment of adults with symptomatic obstructive hypertrophic cardiomyopathy (oHCM) |
| Dosage Form:                                      | Tablets                                                                             |
| Strength:                                         | 5 mg, 10 mg, 15 mg, 20 mg                                                           |
| Route of administration:                          | Oral                                                                                |

<sup>a</sup> Proprietary Name Safety Summary for Myqorzo. (b) (4) 2024 SEP 26. Available from: <\\CDSESUB1\EVSPROD\nda219083\0004\m1\us\118-proprietary-names\proprietary-names.pdf>.

<sup>b</sup> Topper, C. Proprietary Name Review for Myqorzo (IND 138814). Silver Spring (MD): FDA, CDER, OSE, DMEPA 2 (US); 2023 JUL 05. PNR ID No. 2023-1044725017.

<sup>c</sup> Proprietary Names (NDA 219083 Aficamten). South San Francisco (CA): Cytokinetics, Incorporated; 2024 SEP 26. Available from: <\\CDSESUB1\EVSPROD\nda219083\0004\m1\us\118-proprietary-names\proprietary-names.pdf>.

<sup>d</sup> Proposed prescribing information for Myqorzo. South San Francisco (CA): Cytokinetics, Incorporated; 2024 AUG 14. Available from: <\\CDSESUB1\EVSPROD\nda219083\0002\m1\us\114-labeling\draft\labeling\draft-labeling-text.pdf>.

**Table 1. Relevant Product Information for Myqorzo****Dose and frequency:****Recommended Dosage and Administration**

The recommended starting dose of MYQORZO is 5 mg orally once daily. Increase the dose every 2 to 8 weeks by 5 mg until a maintenance dose or the maximum recommended dose of 20 mg is achieved. The maintenance dose of MYQORZO is individualized based on the patient's LVEF and LVOT-G. Recommendations for dosing based on LVEF and LVOT-G criteria are in Table 1.

**Table 1: Dose Adjustment of MYQORZO**

| LVEF          | Valsalva LVOT-G | Dose Adjustment                                                    |
|---------------|-----------------|--------------------------------------------------------------------|
| ≥55%          | ≥ 30 mmHg       | Increase dose by 5 mg (up to the maximum dose of 20 mg once daily) |
| ≥55%          | < 30 mmHg       | Maintain Dose                                                      |
| <55% and ≥50% | Any             | Maintain Dose                                                      |
| <50% and ≥40% | Any             | Decrease dose by 5 mg* (b) (4)                                     |
| <40%          | Any             | Interrupt treatment for at least 7 days.                           |

\* Dose decrease as follows: 20 mg → 15 mg; 15 mg → 10 mg; 10 mg → 5mg

(b) (4)

After the maintenance dose has been established, assess LVEF and Valsalva LVOT-G every 6 months, or every 3 months in patients with LVEF <55% to ≥50%. Consider monitoring LVEF and adjust dose per Table 1, as needed, in patients (b) (4) intercurrent illness (e.g., severe infection or COVID-19), new arrhythmia (e.g., new or uncontrolled atrial fibrillation or other uncontrolled tachyarrhythmia) or any other conditions that may impair systolic function. Do not increase dose until intercurrent illness or new arrhythmia has resolved or stabilized.

(b) (4) should be taken once daily with or without meals at about the same time every day. (b) (4)

(b) (4)

Swallow tablets whole

(b) (4)

(b) (4)

| Table 1. Relevant Product Information for Myqorzo |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |                   |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |
|---------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|--------------------|-----------------------|-------------------|--------------------------|-----------------------|-------------------|--------------------------|-----------------------|-------------------|--------------------|-----------------------|-------------------|
|                                                   | (b) (4)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                   |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |
| <b>How Supplied:</b>                              | <p>MYQORZO is supplied as purple, film-coated tablets containing 5 mg, 10 mg, 15 mg, or 20 mg aficamten. The tablets are debossed on one side with “CK” and the other side with “5”, “10”, “15” and “20” for the 5 mg, 10 mg, 15 mg, and 20 mg strength, respectively.</p> <p>Tablets are supplied as follows:</p> <table border="1"> <tr> <td>5 mg round tablets</td><td>Bottles of 30 tablets</td><td>NDC: 82112-105-30</td></tr> <tr> <td>10 mg triangular tablets</td><td>Bottles of 30 tablets</td><td>NDC: 82112-110-30</td></tr> <tr> <td>15 mg pentagonal tablets</td><td>Bottles of 30 tablets</td><td>NDC: 82112-115-30</td></tr> <tr> <td>20 mg oval tablets</td><td>Bottles of 30 tablets</td><td>NDC: 82112-120-30</td></tr> </table> |                   | 5 mg round tablets | Bottles of 30 tablets | NDC: 82112-105-30 | 10 mg triangular tablets | Bottles of 30 tablets | NDC: 82112-110-30 | 15 mg pentagonal tablets | Bottles of 30 tablets | NDC: 82112-115-30 | 20 mg oval tablets | Bottles of 30 tablets | NDC: 82112-120-30 |
| 5 mg round tablets                                | Bottles of 30 tablets                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | NDC: 82112-105-30 |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |
| 10 mg triangular tablets                          | Bottles of 30 tablets                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | NDC: 82112-110-30 |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |
| 15 mg pentagonal tablets                          | Bottles of 30 tablets                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | NDC: 82112-115-30 |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |
| 20 mg oval tablets                                | Bottles of 30 tablets                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | NDC: 82112-120-30 |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |
| <b>Storage:</b>                                   | <p>Store at 20°C to 25°C (68°F to 77°F) (b) (4) excursions permitted between 15°C (b) (4) 30°C (b) (4) 59°F (b) (4) 86°F [see USP controlled Room Temperature].</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |                   |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |

## 2 DISCUSSION

The following sections provide information obtained and considered in the overall evaluation of the proposed proprietary name, Myqorzo.

### 2.1 MISBRANDING ASSESSMENT

The Office of Prescription Drug Promotion (OPDP) determined that Myqorzo would not misbrand the proposed product. The Division of Medication Error Prevention and Analysis 2 (DMEPA 2) concurred with the findings of OPDP’s assessment for Myqorzo. The Division of Cardiology and Nephrology (DCN) concurred with the findings of OPDP’s assessment for Myqorzo.

### 2.2 SAFETY ASSESSMENT

The following aspects were considered in the safety evaluation of the proposed proprietary name, Myqorzo.

#### 2.2.1 UNITED STATES ADOPTED NAMES (USAN) SEARCH

There is no USAN stem present in the proposed proprietary name.<sup>e</sup>

<sup>e</sup> USAN stem search conducted on October 21, 2024.

2.2.2 COMPONENTS OF THE PROPOSED PROPRIETARY NAME

Cytokinetics did not provide a derivation or intended meaning for the proposed proprietary name, Myqorzo, in their submission. This proprietary name is comprised of a single word that contains the letter strings “OR”, which is the medical abbreviation for “operating room”. Although we typically discourage the inclusion of medical abbreviations in proprietary names, we determined that the location of the abbreviation “OR” in the middle of the proposed proprietary name, Myqorzo, makes it unlikely for the “OR” letter string to be separated from the surrounding letters in a manner that could lead to confusion in this case. Thus, we do not object to the inclusion of the “OR” letter string in this case.

Beyond this abbreviation, we note that Myqorzo does not contain any other additional components (e.g., a modifier, route of administration, dosage form, etc.) that are misleading or can contribute to medication error.

2.2.3 COMMENTS FROM OTHER REVIEW DISCIPLINES AT INITIAL REVIEW

The Division of Cardiology and Nephrology (DCN) did not forward any comments or concerns relating to Myqorzo at the initial phase of the review.

2.2.4 FDA PRESCRIPTION SIMULATION STUDIES

Sixty-eight practitioners participated in DMEPA’s prescription simulation studies for Myqorzo. The responses did not overlap with any currently marketed products nor did the responses sound or look similar to any currently marketed products or any products in the pipeline. Appendix B contains the results from the prescription simulation studies.

2.2.5 PHONETIC AND ORTHOGRAPHIC COMPUTER ANALYSIS (POCA) SEARCH RESULTS

Our POCA search<sup>f</sup> identified 21 names with a combined score of ≥55% or an individual orthographic or phonetic score of ≥70%. We had identified and evaluated some of the names in our previous proprietary name review. We re-evaluated the previously identified names of concern considering any lessons learned from recent post-marketing experience, which may have altered our previous conclusion regarding the acceptability of the name. We also evaluated previously identified names taking into account the changes in product characteristics listed in Table 2:

| Table 2. Changes in Product Characteristics |                                          |                                                                                     |
|---------------------------------------------|------------------------------------------|-------------------------------------------------------------------------------------|
|                                             | Myqorzo (Previous IND 138814 submission) | Myqorzo (Current NDA 219083 submission)                                             |
| Indication                                  | (b) (4)                                  | Treatment of adults with symptomatic obstructive hypertrophic cardiomyopathy (oHCM) |

<sup>f</sup> POCA search conducted on October 21, 2024 in version 5.9.

| Dose and Frequency | (b) (4)         | <p><b>Recommended Dosage and Administration</b></p> <p>The recommended starting dose of MYQORZO is 5 mg orally once daily. Increase the dose every 2 to 8 weeks by 5 mg until a maintenance dose or the maximum recommended dose of 20 mg is achieved. The maintenance dose of MYQORZO is individualized based on the patient's LVEF and LVOT-G. Recommendations for dosing based on LVEF and LVOT-G criteria are in Table 1.</p> <p><b>Table 1: Dose Adjustment of MYQORZO</b></p> <table border="1"> <thead> <tr> <th>LVEF</th><th>Valsalva LVOT-G</th><th>Dose Adjustment</th></tr> </thead> <tbody> <tr> <td>≥55%</td><td>≥ 30 mmHg</td><td>Increase dose by 5 mg (up to the maximum dose of 20 mg once daily)</td></tr> <tr> <td>≥55%</td><td>&lt; 30 mmHg</td><td>Maintain Dose</td></tr> <tr> <td>&lt;55% and ≥50%</td><td>Any</td><td>Maintain Dose</td></tr> <tr> <td>&lt;50% and ≥40%</td><td>Any</td><td>Decrease dose by 5 mg*</td></tr> <tr> <td>&lt;40%</td><td>Any</td><td>Interrupt treatment for at least 7 days.</td></tr> </tbody> </table> <p>* Dose decrease as follows: 20 mg → 15 mg; 15 mg → 10 mg; 10 mg → 5mg</p> <p>(b) (4)</p> <p>After the maintenance dose has been established, assess LVEF and Valsalva LVOT-G every 6 months, or every 3 months in patients with LVEF &lt;55% to ≥50%. Consider monitoring LVEF and adjust dose per Table 1, as needed, in patients (b) (4) intercurrent illness (e.g., severe infection or COVID-19), new arrhythmia (e.g., new or uncontrolled atrial fibrillation or other uncontrolled tachyarrhythmia) or any other conditions that may impair systolic function. Do not increase dose until intercurrent illness or new arrhythmia has resolved or stabilized.</p> | LVEF | Valsalva LVOT-G | Dose Adjustment | ≥55% | ≥ 30 mmHg | Increase dose by 5 mg (up to the maximum dose of 20 mg once daily) | ≥55% | < 30 mmHg | Maintain Dose | <55% and ≥50% | Any | Maintain Dose | <50% and ≥40% | Any | Decrease dose by 5 mg* | <40% | Any | Interrupt treatment for at least 7 days. |
|--------------------|-----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----------------|-----------------|------|-----------|--------------------------------------------------------------------|------|-----------|---------------|---------------|-----|---------------|---------------|-----|------------------------|------|-----|------------------------------------------|
| LVEF               | Valsalva LVOT-G | Dose Adjustment                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |      |                 |                 |      |           |                                                                    |      |           |               |               |     |               |               |     |                        |      |     |                                          |
| ≥55%               | ≥ 30 mmHg       | Increase dose by 5 mg (up to the maximum dose of 20 mg once daily)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |      |                 |                 |      |           |                                                                    |      |           |               |               |     |               |               |     |                        |      |     |                                          |
| ≥55%               | < 30 mmHg       | Maintain Dose                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |      |                 |                 |      |           |                                                                    |      |           |               |               |     |               |               |     |                        |      |     |                                          |
| <55% and ≥50%      | Any             | Maintain Dose                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |      |                 |                 |      |           |                                                                    |      |           |               |               |     |               |               |     |                        |      |     |                                          |
| <50% and ≥40%      | Any             | Decrease dose by 5 mg*                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |      |                 |                 |      |           |                                                                    |      |           |               |               |     |               |               |     |                        |      |     |                                          |
| <40%               | Any             | Interrupt treatment for at least 7 days.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |      |                 |                 |      |           |                                                                    |      |           |               |               |     |               |               |     |                        |      |     |                                          |



|                          |                            |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |
|--------------------------|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|-----------------------|-------------------|--------------------------|-----------------------|-------------------|--------------------------|-----------------------|-------------------|--------------------|-----------------------|-------------------|
|                          |                            | <div>(b) (4)</div> should be taken once daily with or without meals at about the same time every day. <div>(b) (4)</div> <div>(b) (4)</div> Swallow tablets whole <div>(b) (4)</div> <div>(b) (4)</div>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |
| How Supplied             | Bottles <div>(b) (4)</div> | <p>MYQORZO is supplied as purple, film-coated tablets containing 5 mg, 10 mg, 15 mg, or 20 mg aficamten. The tablets are debossed on one side with “CK” and the other side with “5”, “10”, “15” and “20” for the 5 mg, 10 mg, 15 mg, and 20 mg strength, respectively.</p> <p>Tablets are supplied as follows:</p> <table><tr><td>5 mg round tablets</td><td>Bottles of 30 tablets</td><td>NDC: 82112-105-30</td></tr><tr><td>10 mg triangular tablets</td><td>Bottles of 30 tablets</td><td>NDC: 82112-110-30</td></tr><tr><td>15 mg pentagonal tablets</td><td>Bottles of 30 tablets</td><td>NDC: 82112-115-30</td></tr><tr><td>20 mg oval tablets</td><td>Bottles of 30 tablets</td><td>NDC: 82112-120-30</td></tr></table> | 5 mg round tablets | Bottles of 30 tablets | NDC: 82112-105-30 | 10 mg triangular tablets | Bottles of 30 tablets | NDC: 82112-110-30 | 15 mg pentagonal tablets | Bottles of 30 tablets | NDC: 82112-115-30 | 20 mg oval tablets | Bottles of 30 tablets | NDC: 82112-120-30 |
| 5 mg round tablets       | Bottles of 30 tablets      | NDC: 82112-105-30                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |
| 10 mg triangular tablets | Bottles of 30 tablets      | NDC: 82112-110-30                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |
| 15 mg pentagonal tablets | Bottles of 30 tablets      | NDC: 82112-115-30                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |
| 20 mg oval tablets       | Bottles of 30 tablets      | NDC: 82112-120-30                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |
| Storage                  | <div>(b) (4)</div>         | <p>Store at 20°C to 25°C (68°F to 77°F) <div>(b) (4)</div> excursions permitted between 15°C <div>(b) (4)</div> 30°C ( <div>(b) (4)</div> 59°F <div>(b) (4)</div> 86°F) [see USP controlled Room Temperature].</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                    |                       |                   |                          |                       |                   |                          |                       |                   |                    |                       |                   |

Our evaluation has not altered our previous conclusion regarding the acceptability of the proposed proprietary name, Myqorzo. Therefore, we identified one name not previously analyzed. This name is included in Table 3 below.

## 2.2.6 NAMES RETRIEVED FOR REVIEW ORGANIZED BY NAME PAIR SIMILARITY

Table 3 provides the one name retrieved from our POCA search. This name pair is organized as highly similar, moderately similar, or low similarity for further evaluation.

| Table 3. Names Retrieved for Review Organized by Name Pair Similarity                       |                 |
|---------------------------------------------------------------------------------------------|-----------------|
| Similarity Category                                                                         | Number of Names |
| Highly similar name pair:<br>combined match percentage score $\geq 70\%$                    | 0               |
| Moderately similar name pair:<br>combined match percentage score $\geq 55\%$ to $\leq 69\%$ | 1               |
| Low similarity name pair:<br>combined match percentage score $\leq 54\%$                    | 0               |

## 2.2.7 SAFETY ANALYSIS OF NAMES WITH POTENTIAL ORTHOGRAPHIC, SPELLING, AND PHONETIC SIMILARITIES

Our analysis of the one name contained in Table 3 determined none of the names will pose a risk for confusion with Myqorzo as described in Appendices C through H.

## 2.2.8 COMMUNICATION OF DMEPA'S DETERMINATION

On December 6, 2024, DMEPA 2 communicated our determination to the Division of Cardiology and Nephrology (DCN).

## 3 CONCLUSION

The proposed proprietary name, Myqorzo, is conditionally acceptable.

### 3.1 COMMENTS TO CYTOKINETICS, INCORPORATED

We have completed our review of the proposed proprietary name, Myqorzo, and have concluded that this proprietary name is conditionally acceptable.

If any of the proposed product characteristics as stated in your submission, received on September 26, 2024 are altered prior to approval of the marketing application, the proprietary name must be resubmitted for review.

## 4 REFERENCES

### 1. *United States Adopted Names (USAN) Stems*

USAN Stems List contains all the recognized USAN stems, available at <https://www.ama-assn.org/about/united-states-adopted-names/united-states-adopted-names-approved-stems>.

### 2. *Phonetic and Orthographic Computer Analysis (POCA)*

POCA is a system that FDA designed. As part of the name similarity assessment, POCA is used to evaluate proposed names via a phonetic and orthographic algorithm. The proposed proprietary name is converted into its phonemic representation before it runs through the phonetic algorithm. Likewise, an orthographic algorithm exists that operates in a similar fashion. POCA is publicly accessible.

### *Drugs@FDA*

Drugs@FDA is an FDA Web site that contains most of the drug products approved in the United States since 1939. The majority of labels, approval letters, reviews, and other information are available for drug products approved from 1998 to the present. Drugs@FDA contains official information about FDA-approved brand name and generic drugs; therapeutic biological products, prescription and over-the-counter human drugs; and discontinued drugs (see Drugs@FDA Glossary of Terms, available at <https://www.fda.gov/drugs/drug-approvals-and-databases/drugsfda-glossary-terms>).

### *RxNorm*

RxNorm contains the names of prescription and many OTC drugs available in the United States. RxNorm includes generic and branded:

- Clinical drugs – pharmaceutical products given to (or taken by) a patient with therapeutic or diagnostic intent
- Drug packs – packs that contain multiple drugs, or drugs designed to be administered in a specified sequence

Radiopharmaceuticals, contrast media, food, dietary supplements, and medical devices, such as bandages and crutches, are all out of scope for RxNorm (<http://www.nlm.nih.gov/research/umls/rxnorm/overview.html>).

### *Purple Book*

The Purple Book is an online database that contains information about biological products, including biosimilar and interchangeable biological products, licensed (approved) by the FDA under the Public Health Service (PHS) Act. See Purple Book: Lists of Licensed Biological Products with Reference Product Exclusivity and Biosimilarity or Interchangeability Evaluations, available at <https://www.fda.gov/drugs/therapeutic-biologics-applications-bla/purple-book-lists-licensed-biological-products-reference-product-exclusivity-and-biosimilarity-or>.

### *Division of Medication Errors Prevention and Analysis pending proprietary name requests*

This is a list of proposed and pending names that is generated by the Division of Medication Error Prevention and Analysis.

## 5 APPENDICES

### APPENDIX A. FDA'S PROPRIETARY NAME RISK ASSESSMENT EVALUATES PROPOSED PROPRIETARY NAMES FOR MISBRANDING AND SAFETY CONCERNS

1. **Misbranding Assessment:** For prescription drug products, the Office of Prescription Drug Promotion (OPDP) assesses the name for misbranding concerns. For over-the-counter (OTC) drug products, the misbranding assessment of the proposed name is conducted by the Office of Non-Prescription Drugs (ONPD). OPDP or ONPD evaluates proposed proprietary names to determine if the name is false or misleading, such as by making misrepresentations with respect to safety or efficacy. For example, a fanciful proprietary name may misbrand a product by suggesting that it has some unique effectiveness or composition when it does not (21 CFR 201.10(c)(3)). OPDP or ONPD provides their opinion to DMEPA for consideration in the overall acceptability of the proposed proprietary name.
2. **Safety Assessment:** The safety assessment is conducted by DMEPA, and includes the following:
  - a. **Preliminary Assessment:** We consider inclusion of USAN stems or other characteristics that when incorporated into a proprietary name may cause or contribute to medication errors (i.e., dosing interval, dosage form/route of administration, medical or product name abbreviations, names that include or suggest the composition of the drug product, etc.) See prescreening checklist below in Table 3\*. DMEPA defines a medication error as any preventable event that may cause or lead to inappropriate medication use or patient harm while the medication is in the control of the health care professional, patient, or consumer.<sup>9</sup>

| *Table 3. Prescreening Checklist for Proposed Proprietary Name                                                                                                                                    |                                                                                                                                                                                                                                                   |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Answer the questions in the checklist below. Affirmative answers to any of these questions indicate a potential area of concern that should be carefully evaluated as described in this guidance. |                                                                                                                                                                                                                                                   |
| Y/N                                                                                                                                                                                               | <b>Is the proposed name obviously similar in spelling and pronunciation to other names?</b>                                                                                                                                                       |
|                                                                                                                                                                                                   | Proprietary names should not be similar in spelling or pronunciation to proprietary names, established names, or ingredients of other products.                                                                                                   |
| Y/N                                                                                                                                                                                               | <b>Are there inert or inactive ingredients referenced in the proprietary name?</b>                                                                                                                                                                |
|                                                                                                                                                                                                   | Proprietary names should not incorporate any reference to an inert or inactive ingredient in a way that might create an impression that the ingredient's value is greater than its true functional role in the formulation (21 CFR 201.10(c)(4)). |
| Y/N                                                                                                                                                                                               | <b>Does the proprietary name include combinations of active ingredients?</b>                                                                                                                                                                      |
|                                                                                                                                                                                                   | Proprietary names of fixed combination drug products should not include or suggest the name of one or more, but not all, of its active ingredients (see 21 CFR 201.6(b)).                                                                         |
| Y/N                                                                                                                                                                                               | <b>Is there a United States Adopted Name (USAN) stem in the proprietary name?</b>                                                                                                                                                                 |
|                                                                                                                                                                                                   | Proprietary names should not incorporate a USAN stem in the position that USAN designates for the stem.                                                                                                                                           |

<sup>9</sup> National Coordinating Council for Medication Error Reporting and Prevention. <https://www.nccmerp.org/about-medication-errors>. Last accessed 10/05/2020.

| *Table 3. Prescreening Checklist for Proposed Proprietary Name                                                                                                                                    |                                                                                                                                                                 |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Answer the questions in the checklist below. Affirmative answers to any of these questions indicate a potential area of concern that should be carefully evaluated as described in this guidance. |                                                                                                                                                                 |
| Y/N                                                                                                                                                                                               | Is this proprietary name used for another product that does not share at least one common active ingredient?                                                    |
|                                                                                                                                                                                                   | Drug products that do not contain at least one common active ingredient should not use the same (root) proprietary name.                                        |
| Y/N                                                                                                                                                                                               | Is this a proprietary name of a discontinued product?                                                                                                           |
|                                                                                                                                                                                                   | Proprietary names should not use the proprietary name of a discontinued product if that discontinued drug product does not contain the same active ingredients. |

- b. Phonetic and Orthographic Computer Analysis (POCA): Following the preliminary screening of the proposed proprietary name, DMEPA staff evaluates the proposed name against potentially similar names. In order to identify names with potential similarity to the proposed proprietary name, DMEPA enters the proposed proprietary name in POCA and queries the name against the following drug reference databases, Drugs@FDA, Cerner RxNorm, Purple Book, and names in the review pipeline using a 55% threshold in POCA. DMEPA reviews the combined orthographic and phonetic matches and group the names into one of the following three categories:

- Highly similar pair: combined match percentage score  $\geq 70\%$ .
- Moderately similar pair: combined match percentage score  $\geq 55\%$  to  $\leq 69\%$ .
- Low similarity: combined match percentage score  $\leq 54\%$ .

Using the criteria outlined in the check list (Table 4-6) that corresponds to each of the three categories (highly similar pair, moderately similar pair, and low similarity), DMEPA evaluates the name pairs to determine the acceptability or non-acceptability of a proposed proprietary name. The intent of these checklists is to increase the transparency and predictability of the safety determination of whether a proposed name is vulnerable to confusion from a look-alike or sound-alike perspective. Each bullet below corresponds to the name similarity category cross-references the respective table that addresses criteria that DMEPA uses to determine whether a name presents a safety concern from a look-alike or sound-alike perspective.

- For highly similar names, differences in product characteristics often cannot mitigate the risk of a medication error, including product differences such as strength and dose. Thus, proposed proprietary names that have a combined score of  $\geq 70$  percent are at risk for a look-alike sound-alike confusion which is an area of concern (See Table 4).
- Moderately similar names are further evaluated to identify the presence of attributes that are known to cause name confusion.
  - o Name attributes: We note that the beginning of the drug name plays a significant role in contributing to confusion. Additionally, drug name pairs that start with the same first letter and contain a shared letter string of at least 3 letters in both names are major contributing factor in

the confusion of drug names<sup>h</sup>. We evaluate all moderately similar names retrieved from POCA to identify the above attributes. These names are further evaluated to identify overlapping or similar strengths or doses.

- o Product attributes: Moderately similar names of products that have overlapping or similar strengths or doses represent an area for concern for FDA. The dose and strength information is often located in close proximity to the drug name itself on prescriptions and medication orders, and the information can be an important factor that either increases or decreases the potential for confusion between similarly named drug pairs. The ability of other product characteristics to mitigate confusion (e.g., route, frequency, dosage form) may be limited when the strength or dose overlaps. DMEPA reviews such names further, to determine whether sufficient differences exist to prevent confusion. (See Table 5).
  - Names with low similarity that have no overlap or similarity in strength and dose are generally acceptable (See Table 6) unless there are data to suggest that the name might be vulnerable to confusion (e.g., prescription simulation study suggests that the name is likely to be misinterpreted as a marketed product). In these instances, we would reassign a low similarity name to the moderate similarity category and review according to the moderately similar name pair checklist.
- c. FDA Prescription Simulation Studies: DMEPA also conducts prescription simulation studies using FDA health care professionals.

Four separate studies are conducted within the Centers of the FDA for the proposed proprietary name to determine the degree of confusion of the proposed proprietary name with marketed U.S. drug names (proprietary and established) due to similarity in visual appearance with handwritten prescriptions, verbal pronunciation of the drug name or during computerized provider order entry. The studies employ healthcare professionals (pharmacists, physicians, and nurses), and attempts to simulate the prescription ordering process. The primary Safety Evaluator uses the results to identify vulnerability of the proposed name to be misinterpreted by healthcare practitioners during written, verbal, or electronic prescribing.

In order to evaluate the potential for misinterpretation of the proposed proprietary name during written, verbal, or electronic prescribing of the name, written inpatient medication orders, written outpatient prescriptions, verbal orders, and electronic orders are simulated, each consisting of a combination of marketed and unapproved drug products, including the proposed name.
- d. Comments from Other Review Disciplines: DMEPA requests the Office of New Drugs (OND), Office of Generic Drugs (OGD), and/or Office of Pharmaceutical Quality (OPQ) for their comments or concerns with the proposed proprietary name, ask for any clinical issues that may impact the DMEPA review during the initial phase of the name review.

---

<sup>h</sup> Shah, M, Merchant, L, Characteristics That May Help in the Identification of Potentially Confusing Proprietary Drug Names. Therapeutic Innovation & Regulatory Science, September 2016

Additionally, when applicable, at the same time DMEPA requests concurrence/non-concurrence with OPDP's decision on the name. The primary Safety Evaluator addresses any comments or concerns in the Safety Evaluator's assessment.

The OND/OGD Regulatory Division is contacted a second time following our analysis of the proposed proprietary name. At this point, DMEPA conveys their decision to accept or reject the name.

Additionally, other review disciplines opinions such as OPQ may be considered depending on the proposed proprietary name.

When provided, DMEPA considers external proprietary name studies conducted by or for the Applicant/Sponsor and incorporates the findings of these studies into the overall risk assessment.

The DMEPA primary Safety Evaluator assigned to evaluate the proposed proprietary name is responsible for considering the collective findings, and provides an overall risk assessment of the proposed proprietary name.

| Table 4. Highly Similar Name Pair Checklist (i.e., combined Orthographic and Phonetic score is $\geq$ 70%). |                                                                                                                                                                                         |                    |                                                                                                        |
|-------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|--------------------------------------------------------------------------------------------------------|
| Orthographic Checklist                                                                                      |                                                                                                                                                                                         | Phonetic Checklist |                                                                                                        |
| Y/N                                                                                                         | Do the names begin with different first letters?<br><i>Note that even when names begin with different first letters, certain letters may be confused with each other when scripted.</i> | Y/N                | Do the names have different number of syllables?                                                       |
| Y/N                                                                                                         | Are the lengths of the names dissimilar* when scripted?<br><i>*FDA considers the length of names different if the names differ by two or more letters.</i>                              | Y/N                | Do the names have different syllabic stresses?                                                         |
| Y/N                                                                                                         | Considering variations in scripting of some letters (such as z and f), is there a different number or placement of upstroke/downstroke letters present in the names?                    | Y/N                | Do the syllables have different phonologic processes, such vowel reduction, assimilation, or deletion? |
| Y/N                                                                                                         | Is there different number or placement of cross-stroke or dotted letters present in the names?                                                                                          | Y/N                | Across a range of dialects, are the names consistently pronounced differently?                         |
| Y/N                                                                                                         | Do the infixes of the name appear dissimilar when scripted?                                                                                                                             |                    |                                                                                                        |

|     |                                                               |  |  |
|-----|---------------------------------------------------------------|--|--|
| Y/N | Do the suffixes of the names appear dissimilar when scripted? |  |  |
|-----|---------------------------------------------------------------|--|--|

| Table 5: Moderately Similar Name Pair Checklist (i.e., combined score is $\geq 55\%$ to $\leq 69\%$ ). |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |                                                                                                                                                                                                                                                                                                                                                                                                                             |
|--------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Step 1                                                                                                 | <p>Review the DOSAGE AND ADMINISTRATION and HOW SUPPLIED/STORAGE AND HANDLING sections of the prescribing information (or for OTC drugs refer to the Drug Facts label) to determine if strengths and doses of the name pair overlap or are very similar. Different strengths and doses for products whose names are moderately similar may decrease the risk of confusion between the moderately similar name pairs. Name pairs that have overlapping or similar strengths or doses have a higher potential for confusion and should be evaluated further (see Step 2). Because the strength or dose could be used to express an order or prescription for a particular drug product, overlap in one or both of these components would be reason for further evaluation.</p> <p>For single strength products, also consider circumstances where the strength may not be expressed.</p> <p>For any i.e., drug products comprised of more than one active ingredient, consider whether the strength or dose may be expressed using only one of the components.</p> <p>To determine whether the strengths or doses are similar to your proposed product, consider the following list of factors that may increase confusion:</p> <ul style="list-style-type: none"> <li>• Alternative expressions of dose: 5 mL may be listed in the prescribing information, but the dose may be expressed in metric weight (e.g., 500 mg) or in non-metric units (e.g., 1 tsp, 1 tablet/capsule). Similarly, a strength or dose of 1000 mg may be expressed, in practice, as 1 g, or vice versa.</li> <li>• Trailing or deleting zeros: 10 mg is similar in appearance to 100 mg which may potentiate confusion between a name pair with moderate similarity.</li> <li>• Similar sounding doses: 15 mg is similar in sound to 50 mg</li> </ul> |                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Step 2                                                                                                 | <p>Answer the questions in the checklist below. Affirmative answers to some of these questions suggest that the pattern of orthographic or phonetic differences in the names may reduce the likelihood of confusion for moderately similar names with overlapping or similar strengths or doses.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |                                                                                                                                                                                                                                                                                                                                                                                                                             |
|                                                                                                        | <p>Orthographic Checklist (Y/N to each question)</p> <ul style="list-style-type: none"> <li>• Do the names begin with different first letters?</li> <li>• Note that even when names begin with different first letters, certain letters may be confused with each other when scripted.</li> <li>• Are the lengths of the names dissimilar* when scripted?</li> <li>• *FDA considers the length of names different if the names differ by two or more letters.</li> <li>• Considering variations in scripting of some letters (such as z and f), is</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | <p>Phonetic Checklist (Y/N to each question)</p> <ul style="list-style-type: none"> <li>• Do the names have different number of syllables?</li> <li>• Do the names have different syllabic stresses?</li> <li>• Do the syllables have different phonologic processes, such vowel reduction, assimilation, or deletion?</li> <li>• Across a range of dialects, are the names consistently pronounced differently?</li> </ul> |



**Table 5: Moderately Similar Name Pair Checklist (i.e., combined score is  $\geq 55\%$  to  $\leq 69\%$ ).**



|  |                                                                                                                                                                                                                                                                                                                                                                                                      |  |
|--|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|
|  | <p>there a different number or placement of upstroke/downstroke letters present in the names?</p> <ul style="list-style-type: none"> <li>• Is there different number or placement of cross-stroke or dotted letters present in the names?</li> <li>• Do the infixes of the name appear dissimilar when scripted?</li> <li>• Do the suffixes of the names appear dissimilar when scripted?</li> </ul> |  |
|--|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|

**Table 6. Low Similarity Name Pair Checklist (i.e., combined score is  $\leq 54\%$ ).**

Names with low similarity are generally acceptable unless there are data to suggest that the name might be vulnerable to confusion (e.g., prescription simulation study suggests that the name is likely to be misinterpreted as a marketed product). In these instances, we would reassign a low similarity name to the moderate similarity category and review according to the moderately similar name pair checklist.

## APPENDIX B. PRESCRIPTION SIMULATION SAMPLES AND RESULTS

Figure 1. Myqorzo Study (Conducted on 10/11/2024)

| Handwritten Medication Order Prescription                                                                            | Verbal Prescription                                           |
|----------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------|
| <u>Medication Order:</u><br>        | Myqorzo 5 mg<br>One tablet by mouth once daily<br>Dispense 30 |
| <u>Outpatient Prescription:</u><br> |                                                               |
| CPOE Study Sample (displayed as sans-serif, 12-point, bold font)<br><b>Myqorzo</b>                                   |                                                               |

| FDA Prescription Simulation Responses (Aggregate Report) |           |            |       |      |       |
|----------------------------------------------------------|-----------|------------|-------|------|-------|
| Study Name: Myqorzo                                      |           |            |       |      |       |
| People Received Study: 187                               |           |            |       |      |       |
| People Responded: 68                                     |           |            |       |      |       |
| Total                                                    | 19        | 21         | 10    | 18   | 68    |
| INTERPRETATION                                           | INPATIENT | OUTPATIENT | VOICE | CPOE | TOTAL |
| MICORSO                                                  | 0         | 0          | 1     | 0    | 1     |
| MYCORSO                                                  | 0         | 0          | 1     | 0    | 1     |
| MYCORZO                                                  | 0         | 0          | 8     | 0    | 8     |
| MYGARZO                                                  | 1         | 0          | 0     | 0    | 1     |
| MYGORYO                                                  | 0         | 2          | 0     | 0    | 2     |
| MYGORZA                                                  | 1         | 1          | 0     | 0    | 2     |
| MYGORZO                                                  | 3         | 5          | 0     | 0    | 8     |
| MYGOZO                                                   | 0         | 1          | 0     | 0    | 1     |
| MYQORGO                                                  | 0         | 1          | 0     | 0    | 1     |
| MYQORYO                                                  | 0         | 1          | 0     | 0    | 1     |
| MYQORZ                                                   | 0         | 1          | 0     | 0    | 1     |

|         |    |   |   |    |    |
|---------|----|---|---|----|----|
| MYQORZA | 0  | 1 | 0 | 0  | 1  |
| MYQORZO | 14 | 8 | 0 | 18 | 40 |

APPENDIX C. HIGHLY SIMILAR NAMES (e.g., combined POCA score is  $\geq 70\%$ )

|     |                                                                                                                                                                                          |                |                                                                                                                                                                                                   |
|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| No. | Proposed name: Myqorzo<br>Pronunciation: mye kor' zoe<br>Established name: aficamten<br>Dosage form: Tablets<br>Strengths: 5 mg, 10 mg, 15 mg, 20 mg<br>Dosage: 5 mg to 20 mg once daily | POCA Score (%) | Orthographic and/or phonetic differences in the names sufficient to prevent confusion<br><br>Other prevention of failure mode expected to minimize the risk of confusion between these two names. |
|     | N/A                                                                                                                                                                                      |                |                                                                                                                                                                                                   |

APPENDIX D. MODERATELY SIMILAR NAMES (e.g., combined POCA score is  $\geq 55\%$  to  $\leq 69\%$ ) with no overlap or numerical similarity in Strength and/or Dose

| No. | Name        | POCA Score (%) |
|-----|-------------|----------------|
| 1.  | (b) (4) *** | 58             |

APPENDIX E. MODERATELY SIMILAR NAMES (e.g., combined POCA score is  $\geq 55\%$  to  $\leq 69\%$ ) with overlap or numerical similarity in Strength and/or Dose

|     |                                                                                                                                                                                            |                |                                                                                                                                                                                  |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| No. | Proposed name: Myqorzo<br>Pronunciation: mye kor' zoe<br>Established name: aficamten<br>Dosage form: Tablets<br>Strength(s): 5 mg, 10 mg, 15 mg, 20 mg<br>Dosage: 5 mg to 20 mg once daily | POCA Score (%) | Prevention of Failure Mode<br><br>In the conditions outlined below, the following combination of factors, are expected to minimize the risk of confusion between these two names |
|     | N/A                                                                                                                                                                                        |                |                                                                                                                                                                                  |

APPENDIX F. LOW SIMILARITY NAMES (e.g., combined POCA score is  $\leq 54\%$ )

| No. | Name | POCA Score (%) |
|-----|------|----------------|
|     | N/A  |                |

APPENDIX G. Names not likely to be confused or not used in usual practice settings for the reasons described.

| No. | Name | POCA Score (%) | Failure preventions |
|-----|------|----------------|---------------------|
|     | N/A  |                |                     |

APPENDIX H. Names not likely to be confused due to absence of attributes that are known to cause name confusion<sup>i</sup>.

| No. | Name | POCA Score (%) |
|-----|------|----------------|
|     | N/A  |                |

---

<sup>i</sup> Shah, M, Merchant, L, Chan, I, and Taylor, K. Characteristics That May Help in the Identification of Potentially Confusing Proprietary Drug Names. Therapeutic Innovation & Regulatory Science, September 2016

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

JULIE V NESHIEWAT  
12/06/2024 10:09:02 AM

NICOLE F IVERSON  
12/06/2024 03:26:22 PM

HINA S MEHTA  
12/09/2024 12:11:53 PM