

**CENTER FOR DRUG EVALUATION AND RESEARCH**

**Application Number 75-072**

**FINAL PRINTED LABELING**



100322B0499


# VERAPAMIL HYDROCHLORIDE Extended-Release Tablets, USP

R only

### DESCRIPTION

Verapamil hydrochloride is a calcium ion influx inhibitor (slow-channel blocker or calcium ion antagonist). The tablets are designed for extended-release of the drug in the gastrointestinal tract. Extended-release characteristics are not altered when the bisected tablets are divided in half.

The structural formula of verapamil hydrochloride is:



C<sub>20</sub>H<sub>28</sub>N<sub>2</sub>O<sub>4</sub> · HCl  
Benzenecarboxitrile  
M.W. = 491.07

α-[3-[[2-(3,4-dimethoxyphenyl)ethyl] methylamino] propyl]-3,4-dimethoxy-α-(1-methyl) hydrochloride  
Verapamil hydrochloride is an almost white, crystalline powder, practically free of odor, with a bitter taste. It is soluble in water, chloroform and methanol. Verapamil hydrochloride is not chemically related to other cardiovascular drugs.  
Verapamil hydrochloride extended-release tablets contain the following inactive ingredients: hydroxypropyl methylcellulose, magnesium stearate, methacrylic acid microcrystalline cellulose, polyethylene glycol, polyisobutyl 80, polyvinyl, sodium alginate and titanium dioxide. The following are the color additives per tablet strength:

Strength (mg)	Color Additives
120	F&D;C Red #40 Aluminum Lake
240	Yellow #10 Iron Oxide Red #30 Iron Oxide

The *in-vitro* USP Drug Release Test<sub>1</sub> for the Verapamil hydrochloride extended-release 120 mg tablets is pending.

The Verapamil hydrochloride extended-release tablets USP, 240 mg meet USP Drug Release Test 2.

### CLINICAL PHARMACOLOGY

Verapamil is a calcium ion influx inhibitor (slow-channel blocker or calcium ion antagonist) that exerts its pharmacologic effects by modulating the influx of ionic calcium across the cell membrane of the arterial smooth muscle as well as in conductive and contractile myocardial cells.

#### Mechanism of Action

**Essential Hypertension.** Verapamil exerts antihypertensive effects by decreasing systemic vascular resistance, usually without orthostatic decreases in blood pressure or reflex tachycardia. Bradycardia (rate less than 50 beats/min) is uncommon (1.4%). During isometric or dynamic exercise verapamil does not alter systemic cardiac function in patients with normal ventricular function.

Verapamil does not alter total serum calcium levels. However, one report suggested that calcium levels above the normal range may alter the therapeutic effect of verapamil.

Other pharmacologic actions of verapamil include the following:

Verapamil dilates the main coronary arteries and coronary arterioles both in normal and ischemic regions and is a potent inhibitor of coronary artery spasm, whether spontaneous or ergonovine-induced. This property increases myocardial oxygen delivery in patients with coronary artery spasm and is responsible for the effectiveness of verapamil in vasospastic (Prinzmetal's or variant) as well as unstable angina at rest. Whether this effect plays any role in classical effort angina is not clear, but studies of exercise tolerance have not shown an increase in the maximum exercise rate-pressure product, a widely accepted measure of oxygen utilization. This suggests that in general relief of spasm or dilation of coronary arteries is not an important factor in classical angina. Verapamil regularly reduces the total systemic resistance (afterload) against which the heart works both at rest and at a given level of exercise by dilating peripheral arterioles. Electrical activity through the AV node depends to a significant degree upon calcium ions through the slow channel. By decreasing the influx of calcium, verapamil prolongs the effective refractory period within the AV node and slows AV conduction in a rate-related manner.

Normal sinus rhythm is usually not affected, but in patients with sick sinus syndrome verapamil may interfere with sinus node impulse generation and may induce sinus arrest or sinoatrial block. Atrioventricular block can occur in patients without preexisting conduction defects (see WARNINGS).

Verapamil does not alter the normal atrial action potential or intraventricular conduction time but depresses amplitude, velocity of depolarization and conduction in depressed atrial fibers. Verapamil may shorten the antegrade effective refractory period of the accessory bypass tract. Acceleration of ventricular rate and/or ventricular fibrillation has been reported in patients with atrial flutter or atrial fibrillation and a coexisting accessory AV pathway following administration of verapamil (see WARNINGS).

Verapamil has a local anesthetic action that is 1.6 times that of procaine on an equimolar basis. It is not known whether this action is important at the doses used in man.

**Pharmacokinetics and Metabolism.** With the immediate-release formulation, more than 90% of the orally administered dose of verapamil hydrochloride is absorbed. Because of rapid biotransformation of verapamil hydrochloride during its first pass through the portal circulation, bioavailability ranges from 20% to 35%. Peak plasma concentrations are reached between 1 and 2 hours after oral administration. Chronic oral administration of 120 mg of verapamil hydrochloride every 6 hours resulted in plasma levels of verapamil ranging from 125 to 400 ng/mL, with higher values reported occasionally. A nonlinear correlation between the verapamil dose administered and verapamil plasma levels does exist.

In early dose titration with verapamil a relationship exists between verapamil plasma concentration and prolongation of the PR interval. However, during chronic administration the relationship may disappear. The mean elimination half-life in single-dose studies ranged from 2.8 to 7.4 hours. In these same studies, after repetitive dosing, the half-life increased to a range from 4.5 to 12.0 hours (after less than 10 consecutive doses given 6 hours apart). Half-life of verapamil may increase during titration. No relationship has been established between the plasma concentration of verapamil and a reduction in blood pressure. Agents may affect the pharmacokinetics of verapamil. Elimination half-life may be prolonged in the elderly.

In multiple dose studies under fasting conditions the bioavailability measured by AUC of verapamil hydrochloride extended-release tablets was similar to verapamil hydrochloride immediate release; rates of absorption were of course, different.

In a randomized, single-dose, crossover study using healthy volunteers, administration of verapamil hydrochloride extended-release tablets with food produced lower peak concentrations, delayed time to peak, and lesser total absorption (AUC), than when the product was administered to fasting subjects. Similar results were demonstrated for plasma norverapamil. Food thus produces decreased bioavailability (AUC) but a narrower peak-to-trough ratio. Good correlation of dose and response is not available, but controlled studies of extended-release verapamil have shown effectiveness of doses similar to the effective doses of immediate-release verapamil.

In healthy man, orally administered verapamil hydrochloride undergoes extensive metabolism in the liver. Twelve metabolites have been identified in plasma, all except norverapamil are present in trace amounts only. Norverapamil can reach steady-state plasma concentrations approximately equal to those of verapamil itself. The cardiovascular activity of norverapamil appears to be approximately 20% that of verapamil. Approximately 70% of an administered dose is excreted as metabolites in the urine and 16% or more in the feces within 5 days. About 3% to 4% is excreted in the urine as unchanged drug. Approximately 90% is bound to plasma proteins. In patients with hepatic insufficiency, metabolism of immediate-release verapamil is delayed and elimination half-life prolonged up to 14 to 16 hours (see PRECAUTIONS); the volume of distribution is increased and plasma clearance reduced to about 30% of normal. Verapamil clearance values suggest that patients with liver dysfunction may attain therapeutic verapamil plasma concentrations with one third of the oral daily dose required for patients with normal liver function.

After four weeks of oral dosing (120 mg q.i.d.), verapamil and norverapamil levels were noted in the cerebrospinal fluid with estimated partition coefficient of 0.06 for verapamil and 0.04 for norverapamil.

In ten healthy males, administration of oral verapamil (80 mg every 8 hours for 6 days) and a single oral dose of ethanol (0.8 g/kg) resulted in a 17% increase in mean peak ethanol concentrations (106.45±21.40 to 124.23±24.74 mg/ml/3L) compared to placebo. The area under the blood ethanol concentration versus time curve (AUC over 12 hours) increased by 30% (567.99±52 to 475.07±97.24 mg·hr/dL). Verapamil AUCs were positively correlated (r=0.71) to increased ethanol blood AUC values. (See PRECAUTIONS: Drug Interactions.)

**Hemodynamics and Myocardial Metabolism:** Verapamil hydrochloride reduces afterload and myocardial contractility. Improved left ventricular diastolic function in patients with IHSS and those with coronary heart disease has also been observed with verapamil hydrochloride. In most patients, including those with organic cardiac disease, the negative inotropic action of verapamil is countered by reduction of afterload, and cardiac index is usually not reduced. However, in patients with severe left ventricular dysfunction (e.g., pulmonary wedge pressure above 20 mm Hg or ejection fraction less than 30%), or in patients taking beta-adrenergic blocking agents or other cardiodepressant drugs, deterioration of ventricular function may occur (see Drug Interactions).

**Pulmonary Function:** Verapamil does not induce bronchoconstriction and hence does not impair ventilatory function.

### INDICATIONS AND USAGE

Verapamil hydrochloride extended-release tablets are indicated for the management of essential hypertension.

### CONTRAINDICATIONS

- Verapamil hydrochloride extended-release tablets are contraindicated in:
  - Severe left ventricular dysfunction (see WARNINGS)
  - Hypotension (systolic pressure less than 90 mm Hg) or cardiogenic shock
  - Sick sinus syndrome (except in patients with a functioning artificial ventricular pacemaker)
  - Second- or third-degree AV block (except in patients with a functioning artificial ventricular pacemaker)
  - Patients with atrial flutter or atrial fibrillation and an accessory bypass tract (e.g., Wolff-Parkinson-White, Lown-Ganong-Levine syndrome) (see WARNINGS)
  - Patients with known hypersensitivity to verapamil hydrochloride

### WARNINGS

**Heart Failure:** Verapamil has a negative inotropic effect which in most patients is compensated by its afterload reducing (decreased systemic vascular resistance) properties without a net impairment of ventricular performance. In clinical experience with 4,954 patients, 87 (1.8%) developed congestive heart failure or pulmonary edema. Verapamil should be avoided in patients with severe left ventricular dysfunction (e.g., ejection fraction less than 30%) or moderate to severe symptoms of cardiac failure and in patients with any degree of ventricular dysfunction if they are receiving a beta-adrenergic blocker (see Drug Interactions). Patients with milder ventricular dysfunction should, if possible, be controlled with optimum doses of digitalis and/or diuretics before verapamil treatment. (Note interactions with digoxin under PRECAUTIONS.)

**Hypotension:** Occasionally the pharmacologic action of verapamil may produce a decrease in blood pressure below normal levels which may result in dizziness or symptomatic hypotension. The incidence of hypotension observed in 4,954 patients enrolled in clinical trials was 2.5%. In hypertensive patients, decreases in blood pressure below normal are unusual. Tilt-table testing (60 degrees) was not able to induce orthostatic hypotension.

**Elevated Liver Enzymes:** Elevations of transaminases with and without concomitant elevations in alkaline phosphatase and bilirubin have been reported. Such elevations have sometimes been transient and may disappear even in the face of continued verapamil treatment. Several cases of hepatocellular injury related to verapamil have been proven by rechallenge. Half of these had clinical symptoms (malaise, fever, and/or right upper quadrant pain) in addition to elevations of SGPT, SGPT and alkaline phosphatase. Periodic monitoring of liver function in patients receiving verapamil is therefore prudent.

**Accessory Bypass Tract (Wolff-Parkinson-White or Lown-Ganong-Levine).** Some patients with paroxysmal and/or chronic atrial fibrillation or atrial flutter and a coexisting accessory AV pathway have developed increased antegrade conduction across the accessory pathway bypassing the AV node, producing a very rapid ventricular response or ventricular fibrillation after receiving intravenous verapamil (or digitalis). Although a risk of this occurring with oral verapamil has

not been established, such patients receiving oral verapamil may be at risk and its use in these patients is contraindicated (see CONTRAINDICATIONS).

Treatment is usually DC-cardioversion. Cardioversion has been used safely and effectively after oral verapamil.

**Arrhythmogenic Block:** The effect of verapamil on AV conduction and the SA node may cause asymptomatic first-degree AV block and transient bradycardia, sometimes accompanied by nodal escape rhythms. PR-interval prolongation is correlated with verapamil plasma concentrations especially during the early titration phases of therapy. Higher degrees of AV block, however, were infrequently (0.8%) observed. Marked first-degree block or progressive development to second- or third-degree AV block requires a reduction in dosage or, in rare instances, discontinuation of verapamil hydrochloride and institution of appropriate therapy, depending upon the clinical situation. Patients with hypertrophic cardiomyopathy (IHSS): In 120 patients with hypertrophic cardiomyopathy (most of them refractory or intolerant to propranolol) who received therapy with verapamil at doses up to 720 mg/day, a variety of serious adverse effects were seen. Three patients died in pulmonary edema; all had severe left ventricular outflow obstruction and a past history of left ventricular dysfunction. Eight other patients had pulmonary edema and/or severe hypotension, abnormally high (greater than 20 mm Hg) pulmonary wedge pressure and a marked left ventricular outflow obstruction were present in most of these patients. Concomitant administration of quinidine (see Drug Interactions) preceded the severe hypotension in 3 of the 8 patients (2 of whom developed pulmonary edema). Sinus bradycardia occurred in 11% of the patients, second-degree AV block in 4%, and sinus arrest in 2%. It must be appreciated that this group of patients had a serious disease with a high mortality rate. Most adverse effects responded well to dose reduction, and only rarely did verapamil use have to be discontinued.

### PRECAUTIONS

#### General

**Use in Patients with Impaired Hepatic Function:** Since verapamil is highly metabolized by the liver, it should be administered cautiously to patients with impaired hepatic function. Severe liver dysfunction prolongs the elimination half-life of immediate-release verapamil to about 14 to 16 hours; hence, approximately 30% of the dose given to patients with normal liver function should be administered to these patients. Careful monitoring for abnormal prolongation of the PR interval or other signs of excessive pharmacologic effects (see OVERDOSAGE) should be carried out.

**Use in Patients with Attenuated (Decreased) Neuromuscular Transmission:** It has been reported that verapamil decreases neuromuscular transmission in patients with Duchenne's muscular dystrophy, and that verapamil prolongs recovery from the neuromuscular blocking agent vecuronium. It may be necessary to decrease the dosage of verapamil when it is administered to patients with attenuated neuromuscular transmission.

**Use in Patients with Impaired Renal Function:** About 70% of an administered dose of verapamil is excreted as metabolites in the urine. Verapamil is not removed by hemodialysis. Until further data are available, verapamil should be administered cautiously to patients with impaired renal function. These patients should be carefully monitored for abnormal prolongation of the PR interval or other signs of overdosage (see OVERDOSAGE).

#### Drug Interactions

**Beta-Blockers:** Concomitant therapy with beta-adrenergic blockers and verapamil may result in additive negative effects on heart rate, atrioventricular conduction, and/or cardiac contractility. The combination of beta-blockers, verapamil and beta-adrenergic blocking agents has not been studied. However, there have been reports of excessive bradycardia and AV block, including complete heart block, when the combination has been used for the treatment of hypertension. For hypertensive patients, the risks of combined therapy may outweigh the potential benefits. The combination should be used only with caution and close monitoring.

Asymptomatic bradycardia (36 beats/min) with a wandering atrial pacemaker has been observed in a patient receiving concomitant timolol (a beta-adrenergic blocker), eyedrops and oral verapamil.

A decrease in metoprolol and propranolol clearance has been observed when either drug is administered concomitantly with verapamil. A variable effect has been seen when verapamil and atenolol were given together.

**Digitalis:** Clinical use of verapamil in digitalized patients has shown the combination to be well tolerated if digoxin doses are properly adjusted. Chronic verapamil treatment can increase serum digoxin levels by 50% to 75% during the first week of therapy, and this can result in digitalis toxicity. In patients with hepatic cirrhosis the influence of verapamil on digoxin kinetics is magnified. Verapamil may reduce total body clearance and extrarenal clearance of digoxin by 27% and 29%, respectively. Maintenance digitalization doses should be reduced when verapamil is administered, and the patient should be carefully monitored to avoid over- or underdigitalization. Whenever overdigitalization is suspected, the daily dose of digitalis should be reduced or temporarily discontinued. Upon discontinuation of verapamil hydrochloride, the patient should be reassessed to avoid underdigitalization.

**Antihypertensive Agents:** Verapamil administered concomitantly with oral antihypertensive agents (e.g., vasodilators, angiotensin-converting enzyme inhibitors, diuretics, beta-blockers) will usually have an additive effect on lowering blood pressure. Patients receiving these combinations should be appropriately monitored. Concomitant use of agents that attenuate alpha-adrenergic function with verapamil may result in a reduction in blood pressure that is excessive in some patients. Such an effect was observed in one study following the concomitant administration of verapamil and prazosin.

#### Antiarrhythmic Agents

**Disopyramide:** Until data on possible interactions between verapamil and disopyramide phosphate are obtained, disopyramide should not be administered within 48 hours before or 24 hours after verapamil administration.

**Flecainide:** A study in healthy volunteers showed that the concomitant administration of flecainide and verapamil may have additive effects on myocardial contractility AV conduction and repolarization. Concomitant therapy with flecainide and verapamil may result in additive negative inotropic effect and prolongation of atrioventricular conduction.

**Quinidine:** In a small number of patients with hypertrophic cardiomyopathy (IHSS) concomitant use of verapamil and quinidine resulted in significant hypotension. Until further data are obtained, combined therapy of verapamil and quinidine in patients with hypertrophic cardiomyopathy should probably be avoided.

The electrophysiologic effects of quinidine and verapamil on AV conduction were studied in 8 patients. Verapamil significantly

USP  
Extended-Release Tablets  
VERAPAMIL  
HYDROCHLORIDE



00322B

VERAPAMIL  
HYDROCHLORIDE  
Extended-Release Tablets,  
USP

MAY 25 1999

has been a report of increased quinidine levels during verapamil therapy.

**Warnings:** Verapamil has been given concomitantly with short- and long-acting nitrates without any undesirable drug interactions. The pharmacologic profile of both drugs and the clinical experience suggest beneficial interactions.

**Other:**

**Alcohol:** Verapamil has been found to significantly inhibit ethanol elimination resulting in elevated blood ethanol concentrations that may prolong the intoxicating effects of alcohol. (see CLINICAL PHARMACOLOGY, Pharmacokinetics and Metabolism)

**Cimetidine:** The interaction between cimetidine and chronically administered verapamil has not been studied. Variable results on clearance have been obtained in acute studies of healthy volunteers; clearance of verapamil was either reduced or unchanged.

**Lithium:** Increased sensitivity to the effects of lithium (neurotoxicity) has been reported during concomitant verapamil-lithium therapy. Lithium levels have been observed sometimes to increase, sometimes to decrease, and sometimes to be unchanged. Patients receiving both drugs must be monitored carefully.

**Carbamazepine:** Verapamil therapy may increase carbamazepine concentrations during combined therapy. This may produce carbamazepine side effects such as diplopia, headache, ataxia, or dizziness.

**Rifampin:** Therapy with rifampin may markedly reduce oral verapamil bioavailability.

**Phenobarbital:** Phenobarbital therapy may increase verapamil clearance.

**Cyclosporin:** Verapamil therapy may increase serum levels of cyclosporin.

**Theophylline:** Verapamil may inhibit the clearance and increase the plasma levels of theophylline.

**Inhalation Anesthetics:** Animal experiments have shown that inhalation anesthetics depress cardiovascular activity by decreasing the inward movement of calcium ions. When used concomitantly, inhalation anesthetics and calcium antagonists, such as verapamil, should each be titrated carefully to avoid excessive cardiovascular depression.

**Neuromuscular Blocking Agents:** Clinical data and animal studies suggest that verapamil may potentiate the activity of neuromuscular blocking agents (curare-like and depolarizing). It may be necessary to decrease the dose of verapamil and/or the dose of the neuromuscular blocking agent when the drugs are used concomitantly.

**Carcinogenesis, Mutagenesis, Impairment of Fertility:** An 18-month toxicity study in rats, at a low multiple (6-fold) of the maximum recommended human dose, and not the maximum tolerated dose, did not suggest a tumorigenic potential. There was no evidence of a carcinogenic potential of verapamil administered in the diet of rats for two years at doses of 10, 35, and 120 mg/kg/day or approximately 1x, 3.5x, and 12x respectively the maximum recommended human daily dose (480 mg/day or 9.6 mg/kg/day).

Verapamil was not mutagenic in the Ames test in 5 test strains at 3 mg per plate with or without metabolic activation.

Studies in female rats at daily dietary doses up to 5.5 times (55 mg/kg/day) the maximum recommended human dose did not show impaired fertility. Effects on male fertility have not been determined.

**Pregnancy:** Pregnancy Category C. Reproduction studies have been performed in rabbits and rats at oral doses up to 1.5 (15 mg/kg/day) and 6 (60 mg/kg/day) times the human oral daily dose, respectively, and have revealed no evidence of teratogenicity in the rat. However, this inhibition of the human dose was embryocidal and retarded fetal growth and development probably because of adverse maternal effects reflected in reduced weight gains of the dams. This oral dose has also been shown to cause hypotension in rats. There are no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed. Verapamil crosses the placental barrier and can be detected in umbilical vein blood at delivery.

**Labor and Delivery:** It is not known whether the use of verapamil during labor or delivery has immediate or delayed adverse effects on the fetus, or whether it prolongs the duration of labor or increases the need for forceps delivery or other obstetric intervention. Such adverse experiences have not been reported in the literature, despite a long history of use of verapamil in Europe in the treatment of cardiac side effects of beta-adrenergic agonist agents used to treat premature labor.

**Nursing Mothers:** Verapamil is excreted in human milk. Because of the potential for adverse reactions in nursing infants from verapamil, nursing should be discontinued while verapamil is administered.

**Pediatric Use:** Safety and efficacy of verapamil hydrochloride extended-release tablets in pediatric patients below the age of 18 years have not been established.

**Animal Pharmacology and/or Animal Toxicology:** In chronic animal toxicology studies verapamil caused lenticular and/or suture line changes at 30 mg/kg/day or greater and frank cataracts at 62.5 mg/kg/day or greater in the beagle dog but not in the rat. Development of cataracts due to verapamil has not been reported in man.

**ADVERSE REACTIONS**

Serious adverse reactions are uncommon when verapamil therapy is initiated with upward dose titration within the recommended single and total daily dose. See WARNINGS for discussion of heart failure, hypotension, elevated liver enzymes, AV block, and rapid ventricular response. Reversible (upon discontinuation of verapamil) non-obstructive paralytic ileus has been infrequently reported in association with the use of verapamil. The following reactions to orally administered verapamil occurred at rates greater than 1.0% or occurred at lower rates but appeared clearly drug-related in clinical trials in 4,954 patients:

Constipation	7.3%
Dizziness	3.3%
Nausea	2.7%
Hypotension	2.5%
Headache	2.2%
Edema	1.9%
CHF Pulmonary Edema	1.8%
Fatigue	1.7%
Dyspnea	1.4%
Bradycardia (HR <50/min)	1.4%
AV Block total (1° 2° 3°)	1.2%
2° and 3°	0.8%
Rash	1.2%
Flushing	0.6%

Elevated Liver Enzymes (see WARNINGS)

digoxinized patients who had atrial fibrillation or atrial flutter, ventricular rates below 50/min at rest occurred in 15% of patients and asymptomatic hypotension occurred in 5% of patients.

The following reactions, reported in 1% or less of patients, occurred under conditions (open trials, marketing experience) where a causal relationship is uncertain; they are listed to alert the physician to a possible relationship:

**Cardiovascular:** angina pectoris, sinoventricular dissociation, chest pain, claudication, myocardial infarction, palpitations, purpura (vasculitis), syncope.

**Digestive System:** diarrhea, dry mouth, gastrointestinal distress, gingival hyperplasia.

**Hemic and Lymphatic:** ecchymosis or bruising.

**Nervous System:** cerebrovascular accident, confusion, equilibrium disorders, insomnia, muscle cramps, paresthesia, psychotic symptoms, shakiness, somnolence.

**Skin:** arthralgia and rash, exanthema, hair loss, hyperkeratosis, maculae, sweating, urticaria, Stevens-Johnson syndrome, erythema multiforme.

**Special Senses:** blurred vision, tinnitus.

**Urogenital:** gynecomastia, galactorrhea/hyperprolactinemia, increased urination, spotty menstruation, impotence.

**Treatment of Acute Cardiovascular Adverse Reactions:** The frequency of cardiovascular adverse reactions that require therapy is rare; hence, experience with their treatment is limited. Whenever severe hypotension or complete AV block occurs following oral administration of verapamil, the appropriate emergency measures should be applied immediately; e.g., intravenously administered isoproterenol hydrochloride, norepinephrine bitartrate, atropine sulfate (all in the usual doses), or calcium gluconate (10% solution). In patients with hypertrophic cardiomyopathy (HSC), alpha-adrenergic agents (phenylephrine hydrochloride, metaraminol bitartrate or methoxamine hydrochloride) should be used to maintain blood pressure, and isoproterenol and norepinephrine should be avoided. If further support is necessary, dopamine hydrochloride or dobutamine hydrochloride may be administered. Actual treatment and dosage should depend on the severity of the clinical situation and the judgment and experience of the treating physician.

**OVERDOSAGE**

Overdose with verapamil may lead to pronounced hypotension, bradycardia, and conduction system abnormalities (e.g., junctional rhythm with AV dissociation and high degree AV block, including asystole). Other symptoms secondary to hypoperfusion (e.g., metabolic acidosis, hyperglycemia, hyperkalemia, renal dysfunction, and convulsions) may be evident.

Treat all verapamil overdoses as serious and maintain observation for at least 48 hours (especially verapamil hydrochloride extended-release tablets), preferably under continuous hospital care. Delayed pharmacodynamic consequences may occur with the sustained release formulation. Verapamil is known to decrease gastrointestinal transit time. In overdose, tablets of verapamil hydrochloride extended-release have occasionally been reported to form concretions within the stomach or intestines. These concretions have not been visible on plain radiographs of the abdomen, and no medical means of gastrointestinal emptying is of proven efficacy in removing them. Endoscopy might reasonably be considered in cases of massive overdose when symptoms are unusually prolonged.

Treatment of overdose should be supportive. Beta adrenergic stimulation or parenteral administration of calcium injections may increase calcium ion flux across the slow channel, and have been used effectively in treatment of deliberate overdose with verapamil. Continued treatment with large doses of calcium may produce a response. In a few reported cases overdose with calcium channel blockers that was initially refractory to atropine became more responsive to this treatment when the patients received large doses (close to 1 gram/hour for more than 24 hours) of calcium chloride. Verapamil cannot be removed by hemodialysis. Clinically significant hypotensive reactions or high degree AV block should be treated with vasopressor agents or cardiac pacing, respectively. Asystole should be handled by the usual measures including cardiopulmonary resuscitation.

**DOSAGE AND ADMINISTRATION**

**Essential Hypertension**

The dose of verapamil hydrochloride extended-release tablets should be individualized by titration and the drug should be administered with food. Initiate therapy with 180 mg of extended-release verapamil hydrochloride tablets, given in the morning. Lower, initial doses of 120 mg a day may be warranted in patients who may have an increased response to verapamil (e.g., the elderly or small people). Upward titration should be based on therapeutic efficacy and safety evaluated weekly and approximately 24 hours after the previous dose. The antihypertensive effects of verapamil hydrochloride extended-release tablets are evident within the first week of therapy.

If adequate response is not obtained with 180 mg of verapamil hydrochloride extended-release tablets, the dose may be titrated upward in the following manner:

- 240 mg each morning
- 180 mg each morning plus 180 mg each evening, or 240 mg each morning plus 120 mg each evening
- 240 mg every 12 hours

When switching from immediate-release verapamil hydrochloride tablets to verapamil hydrochloride extended-release tablets the total daily dose in milligrams may remain the same.

**HOW SUPPLIED**

Verapamil hydrochloride extended-release 240 mg tablets, USP are supplied as light tan colored, capsule-shaped, film-coated tablets containing 240 mg of verapamil hydrochloride. The tablet is bisected on one side and debossed "482" on the other side.

Verapamil hydrochloride extended-release 120 mg tablets are supplied as light pink colored, capsule-shaped, film-coated tablets containing 120 mg of verapamil hydrochloride. The tablets are debossed "480" on one side.

240 mg (light tan)-	Bottle of 100- NDC # 51285-482-02 Bottle of 500- NDC # 51285-482-04
120 mg (light pink)-	Bottle of 100- NDC # 51285-480-02 Bottle of 500- NDC # 51285-480-04

**STORAGE:** Store between 15° and 25°C (59° and 77°F).

Protect from light and moisture.  
Dispense in a light-resistant container as defined in the USP.

DURAMED PHARMACEUTICALS, INC.  
CINCINNATI, OH 45213 USA

100322B

REV. 04/89

# MESSAGE CONFIRMATION

05/13/98 08:16  
ID=OGD/CDER DOC RM1

DATE	S,R-TIME	DISTANT STATION ID	MODE	PAGES	RESULT
05/13	14'18"	513 731 5270	CALLING	12	OK 0000

05/13/98

08:00

OGD/CDER DOC RM1 → 915137315270

NO. 078

001

monary wedge pressure above 20 mmHg or ejection fraction less than 30%), or in patients taking beta-adrenergic blocking agents or other cardiodepressant drugs, deterioration of ventricular function may occur (see DRUG INTERACTIONS).

**Pulmonary Function:** ISOPTIN does not induce bronchoconstriction and hence, does not impair ventilatory function.

#### INDICATIONS AND USAGE

ISOPTIN SR (verapamil HCl) is indicated for the management of essential hypertension.

#### CONTRAINDICATIONS

Verapamil HCl is contraindicated in:

1. Severe left ventricular dysfunction (see WARNINGS)
2. Hypotension (systolic pressure less than 90 mmHg) or cardiogenic shock
3. Sick sinus syndrome (except in patients with a functioning artificial ventricular pacemaker)
4. Second- or third-degree AV block (except in patients with a functioning artificial ventricular pacemaker).
5. Patients with atrial flutter or atrial fibrillation and an accessory bypass tract (e.g., Wolff-Parkinson-White, Lown-Ganong-Levine syndromes). (see WARNINGS).
6. Patients with known hypersensitivity to verapamil hydrochloride.

#### WARNINGS

**Heart Failure:** Verapamil has a negative inotropic effect which, in most patients, is compensated by its afterload reduction (decreased systemic vascular resistance) properties without a net impairment of ventricular performance. In clinical experience with 4,954 patients, 87 (1.8%) developed congestive heart failure or pulmonary edema. Verapamil should be avoided in patients with severe left ventricular dysfunction (e.g., ejection fraction less than



DURAMED PHARMACEUTICALS, INC.  
CINCINNATI, OH 45213 USA  
LMB274A Rev. 06/08

**DURA med**  
NDC 51285-482-02  
**Verapamil HCl**  
Extended-Release  
Tablets, USP  
**240 mg** B only  
100 Tablets

Each tablet contains:  
Verapamil hydrochloride, USP ..... 240 mg  
Inert Ingredients: See package insert.  
Storage: Store between 15° and 25°C  
(59° and 77°F)  
Protect from light and moisture.  
Dispense in a light, light-resistant container as defined in the USP.

Lot No.:

Exp. Date:



DURAMED PHARMACEUTICALS, INC.  
CINCINNATI, OH 45213 USA  
LMB274A Rev. 06/08

**DURA med**  
NDC 51285-482-02  
**Verapamil HCl**  
Extended-Release  
Tablets, USP  
**240 mg** B only  
100 Tablets

Each tablet contains:  
Verapamil hydrochloride, USP ..... 240 mg  
Inert Ingredients: See package insert.  
Storage: Store between 15° and 25°C  
(59° and 77°F)  
Protect from light and moisture.  
Dispense in a light, light-resistant container as defined in the USP.

Lot No.:

Exp. Date:



DURAMED PHARMACEUTICALS, INC.  
CINCINNATI, OH 45213 USA  
LMB274A Rev. 06/08

**DURA med**  
NDC 51285-482-02  
**Verapamil HCl**  
Extended-Release  
Tablets, USP  
**240 mg** B only  
100 Tablets

Each tablet contains:  
Verapamil hydrochloride, USP ..... 240 mg  
Inert Ingredients: See package insert.  
Storage: Store between 15° and 25°C  
(59° and 77°F)  
Protect from light and moisture.  
Dispense in a light, light-resistant container as defined in the USP.

Lot No.:

Exp. Date:



DURAMED PHARMACEUTICALS, INC.  
CINCINNATI, OH 45213 USA  
LMB274A Rev. 06/08

**DURA med**  
NDC 51285-482-02  
**Verapamil HCl**  
Extended-Release  
Tablets, USP  
**240 mg** B only  
100 Tablets

Each tablet contains:  
Verapamil hydrochloride, USP ..... 240 mg  
Inert Ingredients: See package insert.  
Storage: Store between 15° and 25°C  
(59° and 77°F)  
Protect from light and moisture.  
Dispense in a light, light-resistant container as defined in the USP.

Lot No.:

Exp. Date:

mg



NDC 51285-482-04

**Verapamil HCl**  
Extended-Release  
Tablets, USP

**240 mg** Rx only

500 Tablets

Each tablet contains:  
Verapamil hydrochloride, USP ..... 240 mg  
Usual Dose: See package insert  
Storage: Store between 15° and 25°C (59° and 77°F)  
Protect from light and moisture.  
Dispense in a light, light-resistant container as defined in the USP.


Lot No.:

Exp. Date:

DURAMED PHARMACEUTICALS, INC.  
CINCINNATI, OH 45213 USA

L06575A

Rev. 05/98



0 28176 48204 8



NDC 51285-482-04

**Verapamil HCl**  
Extended-Release  
Tablets, USP

**240 mg** Rx only

500 Tablets

Each tablet contains:  
Verapamil hydrochloride, USP ..... 240 mg  
Usual Dose: See package insert  
Storage: Store between 15° and 25°C (59° and 77°F)  
Protect from light and moisture.  
Dispense in a light, light-resistant container as defined in the USP.

Lot No.:

Exp. Date:

DURAMED PHARMACEUTICALS, INC.  
CINCINNATI, OH 45213 USA

L06575A

Rev. 05/98



0 28176 48204 8



NDC 51285-482-04

**Verapamil HCl**  
Extended-Release  
Tablets, USP

**240 mg** Rx only

500 Tablets

Each tablet contains:  
Verapamil hydrochloride, USP ..... 240 mg  
Usual Dose: See package insert  
Storage: Store between 15° and 25°C (59° and 77°F)  
Protect from light and moisture.  
Dispense in a light, light-resistant container as defined in the USP.

Lot No.:

Exp. Date:

DURAMED PHARMACEUTICALS, INC.  
CINCINNATI, OH 45213 USA

L06575A

Rev. 05/98



0 28176 48204 8



NDC 51285-482-04

**Verapamil HCl**  
Extended-Release  
Tablets, USP

**240 mg** Rx only

500 Tablets

Each tablet contains:  
Verapamil hydrochloride, USP ..... 240 mg  
Usual Dose: See package insert  
Storage: Store between 15° and 25°C (59° and 77°F)  
Protect from light and moisture.  
Dispense in a light, light-resistant container as defined in the USP.

Lot No.:

Exp. Date:

DURAMED PHARMACEUTICALS, INC.  
CINCINNATI, OH 45213 USA

L06575A

Rev. 05/98



0 28176 48204 8