Revised: 12/2019

AUROBINDO

Efavirenz Capsules USP

P152310 1

Item Code With Serial Number in 2D format to be printed at printer's end
The position of product name and pharma code can be changed as per printer feasibility to print 2D data matrix

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use EFAVIRENZ CAPSULES safely and effectively. See full prescribing information for EFAVIRENZ CAPSULES.

EFAVIRENZ capsules for oral use
Initial U.S. Approval: 1998

----------RECENT MAJOR CHANGES----------

Warnings and Precautions, Nervous System Symptoms (5.6)	10/2019
Warnings and Precautions, Immune Reconstitution Syndrome (5.12)	10/2019

----------INDICATIONS AND USAGE----------

Efavirenz is a non-nucleoside reverse transcriptase inhibitor indicated in combination with other antiretroviral agents for the treatment of human immunodeficiency virus type 1 infection in adults and in pediatric patients at least 3 months old and weighing at least 3.5 kg. (1)

----------DOSAGE AND ADMINISTRATION----------

- Efavirenz capsules should be taken orally once daily on an empty stomach, preferably at bedtime. (2)
- Recommended adult dose: 600 mg. (2.2)
- With voriconazole, increase voriconazole maintenance dose to 400 mg every 12 hours and decrease efavirenz dose to 300 mg once daily using the capsule formulation. (2.2)
- With rifampin, increase efavirenz capsules dose to 800 mg once daily for patients weighing 50 kg or more. (2.2)
- Pediatric dosing is based on weight. (2.3)

----------DOSAGE FORMS AND STRENGTHS----------

- Capsules: 50 mg, 100 mg, and 200 mg. (3)

----------CONTRAINDICATIONS----------

- Patients with previously demonstrated hypersensitivity (e.g., Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to any of the components of this product. (4)
- Coadministration of efavirenz with elbasvir/grazoprevir.

----------WARNINGS AND PRECAUTIONS----------

- *QTc prolongation:* Consider alternatives to efavirenz in patients taking other medications with a known risk of Torsade de Pointes or in patients at higher risk of Torsade de Pointes. (5.2)
- *Do not use as a single agent* or add on as a sole agent to a failing regimen. Consider potential for cross-resistance when choosing other agents. (5.3)
- Not recommended with ATRIPLA, which contains efavirenz, emtricitabine, and tenofovir disoproxil fumarate, unless needed for dose adjustment when coadministered with rifampin. (5.4)
- *Serious psychiatric symptoms:* Immediate medical evaluation is recommended for serious psychiatric symptoms such as severe depression or suicidal ideation. (5.5, 17)
- *Nervous system symptoms (NSS):* NSS are frequent and usually begin 1 to 2 days after initiating therapy and resolve in 2 to 4 weeks. Dosing at bedtime may improve tolerability. NSS are not predictive of onset of psychiatric symptoms. (5.6, 6.1, 17)
- *Embryo-Fetal Toxicity:* Avoid administration in the first trimester of pregnancy as fetal harm may occur. (5.7, 8.1)
- *Hepatotoxicity:* Monitor liver function tests before and during treatment in patients with underlying hepatic disease, including hepatitis B or C coinfection, marked transaminase elevations, or who are taking medications associated with liver toxicity. Among reported cases of hepatic failure, a few occurred in patients with no pre-existing hepatic disease. (5.9, 6.1, 8.6)
- *Rash:* Rash usually begins within 1 to 2 weeks after initiating therapy and resolves within 4 weeks. Discontinue if severe rash develops. (5.8, 6.1, 17)
- *Convulsions:* Use caution in patients with a history of seizures. (5.10)
- *Lipids:* Total cholesterol and triglyceride elevations. Monitor before therapy and periodically thereafter. (5.11)
- *Immune reconstitution syndrome:* May necessitate further evaluation and treatment. (5.12)
- *Redistribution/accumulation of body fat:* Observed in patients receiving antiretroviral therapy. (5.13, 17)

----------ADVERSE REACTIONS----------

Most common adverse reactions (>5%, moderate-severe) are impaired concentration, abnormal dreams, rash, dizziness, nausea, headache, fatigue, insomnia, and vomiting. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Aurobindo Pharma USA, Inc. at 1-866-850-2876 or FDA at 1-800-FDA-1088 or *www.fda.gov/medwatch*.

----------DRUG INTERACTIONS----------

- Coadministration of efavirenz can alter the concentrations of other drugs and other drugs may alter the concentrations of efavirenz. The potential for drug-drug interactions should be considered before and during therapy. (7)

----------USE IN SPECIFIC POPULATIONS----------

- *Lactation:* Breastfeeding not recommended. (8.2)
- *Females and Males of Reproductive Potential:* Pregnancy testing and contraception are recommended. (8.3)
- *Hepatic impairment:* Efavirenz is not recommended for patients with moderate or severe hepatic impairment. Use caution in patients with mild hepatic impairment. (8.6)
- *Pediatric patients:* The incidence of rash was higher than in adults. (5.8, 6.2, 8.4)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 12/2019

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE
2 DOSAGE AND ADMINISTRATION
2.1 Hepatic Function
2.2 Adults
2.3 Pediatric Patients
2.4 Capsule Sprinkle Method of Administration
3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
5 WARNINGS AND PRECAUTIONS
5.1 Drug Interactions
5.2 QTc Prolongation
5.3 Resistance
5.4 Coadministration with Related Products
5.5 Psychiatric Symptoms
5.6 Nervous System Symptoms
5.7 Embryo-Fetal Toxicity
5.8 Rash
5.9 Hepatotoxicity
5.10 Convulsions
5.11 Lipid Elevations
5.12 Immune Reconstitution Syndrome
5.13 Fat Redistribution
6 ADVERSE REACTIONS
6.1 Clinical Trials Experience
6.2 Postmarketing Experience
7 DRUG INTERACTIONS
7.1 Potential for Efavirenz to Affect other Drugs
7.2 Potential for Other Drugs to Affect Efavirenz
7.3 QT Prolonging Drugs
7.4 Established and Other Potentially Significant Drug Interactions
7.5 Drugs Without Clinically Significant Interactions with Efavirenz
7.6 Cannabinoid Test Interaction
8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.2 Lactation
8.3 Females and Males of Reproductive Potential
8.4 Pediatric Use
8.5 Geriatric Use
8.6 Hepatic Impairment
10 OVERDOSAGE
11 DESCRIPTION
12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.2 Pharmacodynamics
12.3 Pharmacokinetics
12.4 Microbiology
13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
13.2 Animal Toxicology
14 CLINICAL STUDIES
14.1 Adults
14.2 Pediatric Patients
16 HOW SUPPLIED/STORAGE AND HANDLING
16.1 Capsules
16.3 Storage
17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Efavirenz capsules in combination with other antiretroviral agents are indicated for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in adults and in pediatric patients at least 3 months old and weighing at least 3.5 kg.

2 DOSAGE AND ADMINISTRATION

2.1 Hepatic Function

Monitor hepatic function prior to and during treatment with efavirenz capsules *[see Warnings and Precautions (5.9)]*.
Efavirenz capsules are not recommended in patients with moderate or severe hepatic impairment (Child Pugh B or C) *[see Warnings and Precautions (5.9) and Use in Specific Populations (8.6)]*.

2.2 Adults

The recommended dosage of efavirenz capsules is 600 mg orally, once daily, in combination with a protease inhibitor and/or nucleoside analogue reverse transcriptase inhibitors (NRTIs). It is recommended that efavirenz capsules be taken on an empty stomach, preferably at bedtime. The increased efavirenz concentrations observed following administration of efavirenz capsules with food may lead to an increase in frequency of adverse reactions *[see Clinical Pharmacology (12.3)]*. Dosing at bedtime may improve the tolerability of nervous system symptoms *[see Warnings and Precautions (5.6), Adverse Reactions (6.1),* and *Patient Counseling Information (17)]*. Efavirenz capsules should be swallowed intact with liquid. For patients who cannot swallow capsules, the capsule sprinkle method of administration is recommended *[see Dosage and Administration (2.4)]*.

Concomitant Antiretroviral Therapy

Efavirenz capsules must be given in combination with other antiretroviral medications *[see Indications and Usage (1), Warnings and Precautions (5.3), Drug Interactions (7.1),* and *Clinical Pharmacology (12.3)]*.

Dosage Adjustment

If efavirenz capsules are coadministered with voriconazole, the voriconazole maintenance dose should be increased to 400 mg every 12 hours and the efavirenz dose should be decreased to 300 mg once daily using the capsule formulation (one 200 mg and two 50 mg capsules or six 50 mg capsules).

If efavirenz capsules are coadministered with rifampin to patients weighing 50 kg or more, an increase in the dose of efavirenz capsules to 800 mg once daily is recommended *[see Drug Interactions (7.1, Table 5)* and *Clinical Pharmacology (12.3, Table 8)]*.

2.3 Pediatric Patients

It is recommended that efavirenz capsules be taken on an empty stomach, preferably at bedtime. Table 1 describes the recommended dose of efavirenz capsules for pediatric patients 3 months of age or older and weighing between 3.5 kg and 40 kg *[see Clinical Pharmacology (12.3)]*. The recommended dosage of efavirenz capsules for pediatric patients weighing 40 kg or greater is 600 mg once daily. For pediatric patients who cannot swallow capsules, the capsule contents can be administered with a small amount of food or infant formula using the capsule sprinkle method of administration *[see Dosage and Administration (2.4)]*.

Table 1: Efavirenz Capsules Dosing in Pediatric Patients

Patient Body Weight	Efavirenz Capsules Daily Dose	Number of Capsules[a] and Strength to Administer
3.5 kg to less than 5 kg	100 mg	two 50 mg capsules
5 kg to less than 7.5 kg	150 mg	three 50 mg capsules
7.5 kg to less than 15 kg	200 mg	one 200 mg capsule
15 kg to less than 20 kg	250 mg	one 200 mg + one 50 mg capsule
20 kg to less than 25 kg	300 mg	one 200 mg + two 50 mg capsules
25 kg to less than 32.5 kg	350 mg	one 200 mg + three 50 mg capsules
32.5 kg to less than 40 kg	400 mg	two 200 mg capsules
at least 40 kg	600 mg	three 200 mg capsules

[a] Capsules can be administered intact or as sprinkles *[see Dosage and Administration (2.4)]*.

2.4 Capsule Sprinkle Method of Administration

For pediatric patients at least 3 months old and weighing at least 3.5 kg and adults who cannot swallow capsules, the capsule contents may be administered with a small amount (1 to 2 teaspoons) of food. Use of infant formula for mixing should only be considered for those young infants who cannot reliably consume solid foods. Patients and caregivers should be instructed to open the capsule carefully to avoid spillage or dispersion of the capsule contents into the air. The capsule should be held horizontally over a small container and carefully twisted to open. For patients able to tolerate solid foods, the entire capsule contents should be gently mixed with an age-appropriate soft food, such as applesauce, grape jelly, or yogurt, in the small container. For young infants receiving the capsule sprinkle-infant formula mixture, the entire capsule contents should be gently mixed into 2 teaspoons of reconstituted room temperature infant formula in a small container by carefully stirring with a small spoon, and then drawing up the mixture into a 10 mL oral dosing syringe for administration. After administration of the efavirenz capsules-food or -formula mixture, an additional small amount (approximately 2 teaspoons) of food or formula must be added to the empty mixing container, stirred to disperse any remaining efavirenz residue, and administered to the patient. The efavirenz capsules-food or -formula mixture should be administered within 30 minutes of mixing. No additional food should be consumed for 2 hours after administration of efavirenz capsules.

Further patient instructions on the capsule sprinkle method of administration are provided in the FDA-approved patient labeling (see Patient Information and Instructions for Use).

3 DOSAGE FORMS AND STRENGTHS

Efavirenz capsules USP 50 mg are Yellow/White size '4' hard gelatin capsules imprinted with 'D' on yellow cap and '72' on white body with black edible ink filled with white to off-white colored powder.
Efavirenz capsules USP 100 mg are White/White size '2' hard gelatin capsules imprinted with 'D' on white cap and '71' on white body with black edible ink filled with white to off-white colored powder.
Efavirenz capsules USP 200 mg are Yellow/Yellow size '0EL' hard gelatin capsules imprinted with 'D' on yellow cap and '36' on yellow body with black edible ink filled with white to off-white colored powder.

4 CONTRAINDICATIONS

- Efavirenz capsules are contraindicated in patients with previously demonstrated clinically significant hypersensitivity (e.g., Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to any of the components of this product.
- Coadministration of efavirenz with elbasvir and grazoprevir is contraindicated *[see Warnings and Precautions (5.1)* and *Drug Interactions (7.1)]*.

5 WARNINGS AND PRECAUTIONS

5.1 Drug Interactions

Efavirenz plasma concentrations may be altered by substrates, inhibitors, or inducers of CYP3A. Likewise, efavirenz may alter plasma concentrations of drugs metabolized by CYP3A or CYP2B6. The most prominent effect of efavirenz at steady-state is induction of CYP3A and CYP2B6 *[see Dosage and Administration (2.2)* and *Drug Interactions (7.1)]*.

5.2 QTc Prolongation

QTc prolongation has been observed with the use of efavirenz *[see Drug Interactions (7.3, 7.4)* and *Clinical Pharmacology (12.2)]*. Consider alternatives to efavirenz when coadministered with a drug with a known risk of Torsade de Pointes or when administered to patients at higher risk of Torsade de Pointes.

5.3 Resistance

Efavirenz must not be used as a single agent to treat HIV-1 infection or added on as a sole agent to a failing regimen. Resistant virus emerges rapidly when efavirenz is administered as monotherapy. The choice of new antiretroviral agents to be used in combination with efavirenz should take into consideration the potential for viral cross-resistance.

5.4 Coadministration with Related Products

Coadministration of efavirenz with ATRIPLA (efavirenz 600 mg/emtricitabine 200 mg/tenofovir disoproxil fumarate 300 mg) is not recommended unless needed for dose adjustment (e.g., with rifampin), since efavirenz is one of its active ingredients.

5.5 Psychiatric Symptoms

Serious psychiatric adverse experiences have been reported in patients treated with efavirenz. In controlled trials of 1008 patients treated with regimens containing efavirenz for a mean of 2.1 years and 635 patients treated with control regimens for a mean of 1.5 years, the frequency (regardless of causality) of specific serious psychiatric events among patients who received efavirenz or control regimens, respectively, were severe depression (2.4%, 0.9%), suicidal ideation (0.7%, 0.3%), nonfatal suicide attempts (0.5%, 0), aggressive behavior (0.4%, 0.5%), paranoid reactions (0.4%, 0.3%), and manic reactions (0.2%, 0.3%). When psychiatric symptoms similar to those noted above were combined and evaluated as a group in a multifactorial analysis of data from Study 006, treatment with efavirenz was associated with an increase in the occurrence of these selected psychiatric symptoms. Other factors associated with an increase in the occurrence of these psychiatric symptoms were history of injection drug use, psychiatric history, and receipt of psychiatric medication at study entry; similar associations were observed in both the efavirenz and control treatment groups. In Study 006, onset of new serious psychiatric symptoms occurred throughout the study for both efavirenz-treated and control-treated patients. One percent of efavirenz-treated patients discontinued or interrupted treatment because of one or more of these selected psychiatric symptoms. There have also been occasional postmarketing reports of death by suicide, delusions, and psychosis-like behavior although a causal relationship to the use of efavirenz cannot be determined from these reports. Postmarketing cases of catatonia have also been reported and may be associated with increased efavirenz exposure. Patients with serious psychiatric adverse experiences should seek immediate medical evaluation to assess the possibility that the symptoms may be related to the use of efavirenz, and if so, to determine whether the risks of continued therapy outweigh the benefits *[see Adverse Reactions (6.1)]*.

5.6 Nervous System Symptoms

Fifty-three percent (531/1008) of patients receiving efavirenz in controlled trials reported central nervous system symptoms (any grade, regardless of causality) compared to 25% (156/635) of patients receiving control regimens *[see Adverse Reactions (6.1, Table 3)]*. These symptoms included, but were not limited to, dizziness (28.1% of the 1008 patients), insomnia (16.3%), impaired concentration (8.3%), somnolence (7.0%), abnormal dreams (6.2%), and hallucinations (1.2%). These symptoms were severe in 2.0% of patients; and 2.1% of patients discontinued therapy as a result. These symptoms usually begin during the first or second day of therapy and generally resolve after the first 2 to 4 weeks of therapy. After 4 weeks of therapy, the prevalence of nervous system symptoms of at least moderate severity ranged from 5% to 9% in patients treated with regimens containing efavirenz and from 3% to 5% in patients treated with a control regimen. Patients should be informed that these common symptoms were likely to improve with continued therapy and were not predictive of subsequent onset of the less frequent psychiatric symptoms *[see Warnings and Precautions (5.5)]*. Dosing at bedtime may improve the tolerability of these nervous system symptoms *[see Dosage and Administration (2)]*.

Analysis of long-term data from Study 006 (median follow-up 180 weeks, 102 weeks, and 76 weeks for patients treated with efavirenz + zidovudine + lamivudine, efavirenz + indinavir, and indinavir + zidovudine + lamivudine, respectively) showed that, beyond 24 weeks of therapy, the incidences of new-onset nervous system symptoms among efavirenz-treated patients were generally similar to those in the indinavir-containing control arm.

Late-onset neurotoxicity, including ataxia and encephalopathy (impaired consciousness, confusion, psychomotor slowing, psychosis, delirium), may occur months to years after beginning efavirenz therapy. Some events of late-onset neurotoxicity have occurred in patients with CYP2B6 genetic polymorphisms which are associated with increased efavirenz levels despite standard dosing of efavirenz. Patients presenting with signs and symptoms of serious neurologic adverse experiences should be evaluated promptly to assess the possibility that these events may be related to efavirenz use, and whether discontinuation of efavirenz is warranted.

Patients receiving efavirenz should be alerted to the potential for additive central nervous system effects when efavirenz is used concomitantly with alcohol or psychoactive drugs.

Patients who experience central nervous system symptoms such as dizziness, impaired concentration, and/or drowsiness should avoid potentially hazardous tasks such as driving or operating machinery.

5.7 Embryo-Fetal Toxicity

Efavirenz may cause fetal harm when administered during the first trimester to a pregnant woman. Advise females of reproductive potential who are receiving efavirenz to avoid pregnancy *[see Use in Specific Populations (8.1 and 8.3)]*.

5.8 Rash

In controlled clinical trials, 26% (266/1008) of adult patients treated with 600 mg efavirenz experienced new-onset skin rash compared with 17% (111/635) of those treated in control groups *[see Adverse Reactions (6.1)]*. Rash associated with blistering, moist desquamation, or ulceration occurred in 0.9% (9/1008) of patients treated with efavirenz. The incidence of Grade 4 rash (e.g., erythema multiforme, Stevens-Johnson syndrome) in adult patients treated with efavirenz in all studies and expanded access was 0.1%. Rashes are usually mild-to-moderate maculopapular skin eruptions that occur within the first 2 weeks of initiating therapy with efavirenz (median time to onset of rash in adults was 11 days) and, in most patients continuing therapy with efavirenz, rash resolves within 1 month (median duration, 16 days). The discontinuation rate for rash in adult clinical trials was 1.7% (17/1008).

Rash was reported in 59 of 182 pediatric patients (32%) treated with efavirenz *[see Adverse Reactions (6.2)]*. Two pediatric patients experienced Grade 3 rash (confluent rash with fever, generalized rash), and four patients had Grade 4 rash (erythema multiforme). The median time to onset of rash in pediatric patients was 28 days (range 3 to 1642 days). Prophylaxis with appropriate antihistamines before initiating therapy with efavirenz in pediatric patients should be considered.

Efavirenz can generally be reinitiated in patients interrupting therapy because of rash. Efavirenz should be discontinued in patients developing severe rash associated with blistering, desquamation, mucosal involvement, or fever. Appropriate antihistamines and/or corticosteroids may improve the tolerability and hasten the resolution of rash. For patients who have had a life-threatening cutaneous reaction (e.g., Stevens-Johnson syndrome), alternative therapy should be considered *[see Contraindications (4)]*.

5.9 Hepatotoxicity

Postmarketing cases of hepatitis, including fulminant hepatitis progressing to liver failure requiring transplantation or resulting in death, have been reported in patients treated with efavirenz. Reports have included patients with underlying hepatic disease, including coinfection with hepatitis B or C, and patients without pre-existing hepatic disease or other identifiable risk factors.

Efavirenz is not recommended for patients with moderate or severe hepatic impairment. Careful monitoring is recommended for patients with mild hepatic impairment receiving efavirenz. *[see Adverse Reactions (6.1)* and *Use in Specific Populations (8.6)]*.

Monitoring of liver enzymes before and during treatment is recommended for all patients *[see Dosage and Administration (2.1)]*. Consider discontinuing efavirenz in patients with persistent elevations of serum transaminases to greater than five times the upper limit of the normal range.

Discontinue efavirenz if elevation of serum transaminases is accompanied by clinical signs or symptoms of hepatitis or hepatic decompensation.

5.10 Convulsions

Convulsions have been observed in adult and pediatric patients receiving efavirenz, generally in the presence of known medical history of seizures *[see Nonclinical Toxicology (13.2)]*. Caution should be taken in any patient with a history of seizures. Patients who are receiving concomitant anticonvulsant medications primarily metabolized by the liver, such as phenytoin and phenobarbital, may require periodic monitoring of plasma levels *[see Drug Interactions (7.1)]*.

5.11 Lipid Elevations

Treatment with efavirenz has resulted in increases in the concentration of total cholesterol and triglycerides *[see Adverse Reactions (6.1)]*. Cholesterol and triglyceride testing should be performed before initiating efavirenz therapy and at periodic intervals during therapy.

5.12 Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in patients treated with combination antiretroviral therapy, including efavirenz. During the initial phase of combination antiretroviral treatment, patients whose immune system responds may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium* infection, cytomegalovirus, *Pneumocystis jiroveci* pneumonia [PCP], or tuberculosis), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis, Guillain-Barré syndrome, and autoimmune hepatitis) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable, and can occur many months after initiation of treatment.

5.13 Fat Redistribution

Redistribution/accumulation of body fat including central obesity, dorsocervical fat enlargement (buffalo hump), peripheral wasting, facial wasting, breast enlargement, and "cushingoid appearance" have been observed in patients receiving antiretroviral therapy. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been established.

6 ADVERSE REACTIONS

The most significant adverse reactions observed in patients treated with efavirenz are:

- psychiatric symptoms *[see Warnings and Precautions (5.5)]*,
- nervous system symptoms *[see Warnings and Precautions (5.6)]*,
- rash *[see Warnings and Precautions (5.8)]*,
- hepatotoxicity *[see Warnings and Precautions (5.9)]*

6.1 Clinical Trials Experience

Because clinical studies are conducted under widely varying conditions, the adverse reaction rates reported cannot be directly compared to rates in other clinical studies and may not reflect the rates observed in clinical practice.

Adverse Reactions in Adults

The most common (>5% in either efavirenz treatment group) adverse reactions of at least moderate severity among patients in Study 006 treated with efavirenz in combination with zidovudine/lamivudine or indinavir were rash, dizziness, nausea, headache, fatigue, insomnia, and vomiting.

Selected clinical adverse reactions of moderate or severe intensity observed in ≥2% of efavirenz-treated patients in two controlled clinical trials are presented in Table 2.

Table 2: Selected Treatment-Emergent[a] Adverse Reactions of Moderate or Severe Intensity Reported in ≥2% of Efavirenz-Treated Patients in Studies 006 and ACTG 364

	Study 006 LAM-, NNRTI-, and Protease Inhibitor-Naive Patients			Study ACTG 364 NRTI-experienced, NNRTI-, and Protease Inhibitor-Naive Patients		
Adverse Reactions	Efavirenz[b] + ZDV/LAM (n=412) 180 weeks[c]	Efavirenz[b] + Indinavir (n=415) 102 weeks[c]	Indinavir + ZDV/LAM (n=401) 76 weeks[c]	Efavirenz[b] + Nelfinavir + NRTIs (n=64) 71.1 weeks[c]	Efavirenz[b] + NRTIs (n=65) 70.9 weeks[c]	Nelfinavir + NRTIs (n=66) 62.7 weeks[c]
Body as a Whole						
Fatigue	8%	5%	9%	0	2%	3%
Pain	1%	2%	8%	13%	6%	17%
Central and Peripheral Nervous System						
Dizziness	9%	9%	2%	2%	6%	6%
Headache	8%	5%	3%	5%	2%	3%
Insomnia	7%	7%	2%	0	0	2%
Concentration impaired	5%	3%	<1%	0	0	0
Abnormal dreams	3%	1%	0	–	–	–
Somnolence	2%	2%	<1%	0	0	0
Anorexia	1%	<1%	<1%	0	2%	2%
Gastrointestinal						
Nausea	10%	6%	24%	3%	2%	2%
Vomiting	6%	3%	14%	–	–	–
Diarrhea	3%	5%	6%	14%	3%	9%
Dyspepsia	4%	4%	6%	0	0	2%
Abdominal pain	2%	2%	5%	3%	3%	3%
Psychiatric						
Anxiety	2%	4%	<1%	–	–	–
Depression	5%	4%	<1%	3%	0	5%
Nervousness	2%	2%	0	2%	0	2%
Skin & Appendages						
Rash[d]	11%	16%	5%	9%	5%	9%
Pruritus	<1%	1%	1%	9%	5%	9%

[a] Includes adverse events at least possibly related to study drug or of unknown relationship for Study 006. Includes all adverse events regardless of relationship to study drug for Study ACTG 364.
[b] Efavirenz provided as 600 mg once daily.
[c] Median duration of treatment.
[d] Includes erythema multiforme, rash, rash erythematous, rash follicular, rash maculopapular, rash petechial, rash pustular, and urticaria for Study 006 and macules, papules, rash, erythema, redness, inflammation, allergic rash, urticaria, welts, hives, itchy, and pruritus for ACTG 364.

— = Not Specified.
ZDV = zidovudine, LAM = lamivudine.

Pancreatitis has been reported, although a causal relationship with efavirenz has not been established. Asymptomatic increases in serum amylase levels were observed in a significantly higher number of patients treated with efavirenz 600 mg than in control patients (see *Laboratory Abnormalities*).

Nervous System Symptoms

For 1008 patients treated with regimens containing efavirenz and 635 patients treated with a control regimen in controlled trials, Table 3 lists the frequency of symptoms of different degrees of severity and gives the discontinuation rates for one or more of the following nervous system symptoms: dizziness, insomnia, impaired concentration, somnolence, abnormal dreaming, euphoria, confusion, agitation, amnesia, hallucinations, stupor, abnormal thinking, and depersonalization *[see Warnings and Precautions (5.6)]*. The frequencies of specific central and peripheral nervous system symptoms are provided in Table 2.

Table 3: Percent of Patients with One or More Selected Nervous System Symptoms[a,b]

Percent of Patients with:	Efavirenz 600 mg Once Daily (n=1008) %	Control Groups (n=635) %
Symptoms of any severity	52.7	24.6
Mild symptoms[c]	33.3	15.6
Moderate symptoms[d]	17.4	7.7
Severe symptoms[e]	2.0	1.3
Treatment discontinuation as a result of symptoms	2.1	1.1

[a] Includes events reported regardless of causality.
[b] Data from Study 006 and three Phase 2/3 studies.
[c] "Mild" = Symptoms which do not interfere with patient's daily activities.
[d] "Moderate" = Symptoms which may interfere with daily activities.
[e] "Severe" = Events which interrupt patient's usual daily activities.

Psychiatric Symptoms

Serious psychiatric adverse experiences have been reported in patients treated with efavirenz. In controlled trials, psychiatric symptoms observed at a frequency greater than 2% among patients treated with efavirenz or control regimens, respectively, were depression (19%, 16%), anxiety (13%, 9%), and nervousness (7%, 2%).

Rash

In controlled clinical trials, the frequency of rash (all grades, regardless of causality) was 26% for 1008 adults treated with regimens containing efavirenz and 17% for 635 adults treated with a control regimen. Most reports of rash were mild or moderate in severity. The frequency of Grade 3 rash was 0.8% for efavirenz-treated patients and 0.3% for control groups, and the frequency of Grade 4 rash was 0.1% for efavirenz and 0 for control groups. The discontinuation rates as a result of rash were 1.7% for efavirenz-treated patients and 0.3% for control groups *[see Warnings and Precautions (5.8)]*.

Experience with efavirenz in patients who discontinued other antiretroviral agents of the NNRTI class is limited. Nineteen patients who discontinued nevirapine because of rash have been treated with efavirenz. Nine of these patients developed mild-to-moderate rash while receiving therapy with efavirenz, and two of these patients discontinued because of rash.

Laboratory Abnormalities

Selected Grade 3 to 4 laboratory abnormalities reported in ≥2% of efavirenz-treated patients in two clinical trials are presented in Table 4.

Table 4: Selected Grade 3 to 4 Laboratory Abnormalities Reported in ≥2% of Efavirenz-Treated Patients in Studies 006 and ACTG 364

		Study 006 LAM-, NNRTI-, and Protease Inhibitor-Naive Patients			Study ACTG 364 NRTI-experienced, NNRTI-, and Protease Inhibitor-Naive Patients		
Variable	Limit	Efavirenz[a] + ZDV/LAM (n=412) 180 weeks[b]	Efavirenz[a] + Indinavir (n=415) 102 weeks[b]	Indinavir + ZDV/LAM (n=401) 76 weeks[b]	Efavirenz[a] + Nelfinavir + NRTIs (n=64) 71.1 weeks[b]	Efavirenz[a] + NRTIs (n=65) 70.9 weeks[b]	Nelfinavir + NRTIs (n=66) 62.7 weeks[b]
Chemistry							
ALT	>5 x ULN	5%	8%	5%	2%	6%	3%
AST	>5 x ULN	5%	6%	5%	6%	8%	8%
GGT[c]	>5 x ULN	8%	7%	3%	5%	0	5%
Amylase	>2 x ULN	4%	4%	1%	0	6%	2%
Glucose	>250 mg/dL	3%	3%	3%	5%	2%	3%
Triglycerides[d]	≥751 mg/dL	9%	6%	6%	11%	8%	17%
Hematology							
Neutrophils	<750/mm³	10%	3%	5%	2%	3%	2%

[a] Efavirenz provided as 600 mg once daily.
[b] Median duration of treatment.
[c] Isolated elevations of GGT in patients receiving efavirenz may reflect enzyme induction not associated with liver toxicity.
[d] Nonfasting.

ZDV = zidovudine, LAM = lamivudine, ULN = upper limit of normal, ALT = alanine aminotransferase, AST = aspartate aminotransferase, GGT = gamma-glutamyltransferase.

Patients Coinfected with Hepatitis B or C

Liver function tests should be monitored in patients with a history of hepatitis B and/or C. In the long-term data set from Study 006, 137 patients treated with efavirenz-containing regimens (median duration of therapy, 68 weeks) and 84 treated with a control regimen (median duration, 56 weeks) were seropositive at screening for hepatitis B (surface antigen positive) and/or C (hepatitis C antibody positive). Among these coinfected patients, elevations in AST to greater than five times ULN developed in 13% of patients in the efavirenz arms and 7% of those in the control arm, and elevations in ALT to greater than five times ULN developed in 20% of patients in the efavirenz arms and 7% of patients in the control arm. Among coinfected patients, 3% of those treated with efavirenz-containing regimens and 2% in the control arm discontinued from the study because of liver or biliary system disorders *[see Warnings and Precautions (5.9)]*.

Lipids

Increases from baseline in total cholesterol of 10 to 20% have been observed in some uninfected volunteers receiving efavirenz. In patients treated with efavirenz + zidovudine + lamivudine, increases from baseline in nonfasting total cholesterol and HDL of approximately 20% and 25%, respectively, were observed. In patients treated with efavirenz + indinavir, increases from baseline in nonfasting cholesterol and HDL of approximately 40% and 35%, respectively, were observed. Nonfasting total cholesterol levels ≥240 mg/dL and ≥300 mg/dL were reported in 34% and 9%, respectively, of patients treated with efavirenz + zidovudine + lamivudine; 54% and 20%, respectively, of patients treated with efavirenz + indinavir; and 28% and 4%, respectively, of patients treated with indinavir + zidovudine + lamivudine. The effects of efavirenz on triglycerides and LDL in this study were not well characterized since samples were taken from nonfasting patients. The clinical significance of these findings is unknown *[see Warnings and Precautions (5.11)]*.

Adverse Reactions in Pediatric Patients

Because clinical studies are conducted under widely varying conditions, the adverse reaction rates reported cannot be directly compared to rates in other clinical studies and may not reflect the rates observed in clinical practice.

Assessment of adverse reactions is based on three clinical trials in 182 HIV-1 infected pediatric patients (3 months to 21 years of age) who received efavirenz in combination with other antiretroviral agents for a median of 123 weeks. The adverse reactions observed in the three trials were similar to those observed in clinical trials in adults except that rash was more common in pediatric patients (32% for all grades regardless of causality) and more often of higher grade (i.e., more severe). Two (1.1%) pediatric patients experienced Grade 3 rash (confluent rash with fever, generalized rash), and four (2.2%) pediatric patients had Grade 4 rash (all erythema multiforme). Five pediatric patients (2.7%) discontinued from the study because of rash *[see Warnings and Precautions (5.8)]*.

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of efavirenz. Because these reactions are reported voluntarily from a population of unknown size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Body as a Whole: allergic reactions, asthenia, redistribution/accumulation of body fat *[see Warnings and Precautions (5.13)]*

Central and Peripheral Nervous System: abnormal coordination, ataxia, encephalopathy, cerebellar coordination and balance disturbances, convulsions, hypoesthesia, paresthesia, neuropathy, tremor, vertigo

Endocrine: gynecomastia

Gastrointestinal: constipation, malabsorption

Cardiovascular: flushing, palpitations

Liver and Biliary System: hepatic enzyme increase, hepatic failure, hepatitis.

Metabolic and Nutritional: hypercholesterolemia, hypertriglyceridemia

Musculoskeletal: arthralgia, myalgia, myopathy

Psychiatric: aggressive reactions, agitation, delusions, emotional lability, mania, neurosis, paranoia, psychosis, suicide, catatonia

Respiratory: dyspnea

Skin and Appendages: erythema multiforme, photoallergic dermatitis, Stevens-Johnson syndrome

Special Senses: abnormal vision, tinnitus

7 DRUG INTERACTIONS

7.1 Potential for Efavirenz to Affect other Drugs

Efavirenz has been shown *in vivo* to induce CYP3A and CYP2B6. Other compounds that are substrates of CYP3A or CYP2B6 may have decreased plasma concentrations when coadministered with efavirenz.

7.2 Potential for Other Drugs to Affect Efavirenz

Drugs that induce CYP3A activity (e.g., phenobarbital, rifampin, rifabutin) would be expected to increase the clearance of efavirenz resulting in lowered plasma concentrations *[see Dosage and Administration (2.2)]*.

7.3 QT Prolonging Drugs

There is limited information available on the potential for a pharmacodynamic interaction between efavirenz and drugs that prolong the QTc interval. QTc prolongation has been observed with the use of efavirenz *[see Clinical Pharmacology (12.2)]*. Consider alternatives to efavirenz when coadministered with a drug with a known risk of Torsade de Pointes.

7.4 Established and Other Potentially Significant Drug Interactions

Drug interactions with efavirenz are summarized in Table 5. For pharmacokinetics data, [*see Clinical Pharmacology (12.3)]* Tables 7 and 8. This table includes potentially significant interactions, but is not all inclusive.

**Table 5: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be

Coadministered Drug	Dose	Efavirenz Dose	Number of Subjects	Coadministered Drug (mean % change) C_{max} (90% CI)	AUC (90% CI)	C_{min} (90% CI)
Ethinyl estradiol/ Norgestimate	0.035 mg/0.25 mg x 14 days	600 mg qd x 14 days				
Ethinyl estradiol			21	↔	↔	↔
Norelgestromin			21	↓ 46% (39-52%)	↓ 64% (62-67%)	↓ 82% (79-85%)
Levonorgestrel			6	↓ 80% (77-83%)	↓ 83% (79-87%)	↓ 86% (80-90%)
Lorazepam	2 mg single dose	600 mg qd x 10 days	12	↑ 16% (2-32%)	↔	NA
Methadone	Stable maintenance 35 to 100 mg daily	600 mg qd x 14-21 days	11	↓ 45% (25-59%)	↓ 52% (33-66%)	NA
Bupropion	150 mg single dose (sustained-release)	600 mg qd x 14 days	13	↓34% (21-47%)	↓ 55% (48-62%)	NA
Hydroxy-bupropion				↑50% (20-80%)	↔	NA
Paroxetine	20 mg qd x 14 days	600 mg qd x 14 days	16	↔	↔	↔
Sertraline	50 mg qd x 14 days	600 mg qd x 14 days	13	↓ 29% (15-40%)	↓ 39% (27-50%)	↓ 46% (31-58%)

↑ Indicates increase ↓ Indicates decrease ↔ Indicates no change or a mean increase or decrease of <10%.
[a] Compared with atazanavir 400 mg qd alone.
[b] Comparator dose of indinavir was 800 mg q8h x 10 days.
[c] Parallel-group design; n for efavirenz + lopinavir/ritonavir, n for lopinavir/ritonavir alone.
[d] Values are for lopinavir; the pharmacokinetics of ritonavir in this study were unaffected by concurrent efavirenz.
[e] 95% CI.
[f] Soft Gelatin Capsule.
[g] Tenofovir disoproxil fumarate.
[h] 90% CI not available.
[i] Relative to steady-state administration of voriconazole (400 mg for 1 day, then 200 mg po q12h for 2 days).
[j] Not available because of insufficient data.
NA = not available.

Table 8: Effect of Coadministered Drug on Efavirenz Plasma C_{max}, AUC, and C_{min}

Coadministered Drug	Dose	Efavirenz Dose	Number of Subjects	Efavirenz (mean % change) C_{max} (90% CI)	AUC (90% CI)	C_{min} (90% CI)
Indinavir	800 mg q8h x 14 days	200 mg qd x 14 days	11	↔	↔	↔
Lopinavir/ ritonavir	400/100 mg q12h x 9 days	600 mg qd x 9 days	11,12[a]	↔	↓ 16% (↓ 38-↑ 15%)	↓ 16% (↓ 42-↑ 20%)
Nelfinavir	750 mg q8h x 7 days	600 mg qd x 7 days	10	↓ 12% (↓ 32-↑ 13%)[b]	↓ 12% (↓ 35-↑ 18%)[b]	↓ 21% (↓ 53-↑ 33%)
Ritonavir	500 mg q12h x 8 days	600 mg qd x 10 days	9	↑ 14% (4-26%)	↑ 21% (10-34%)	↑ 25% (7-46%)[b]
Saquinavir SGC[c]	1200 mg q8h x 10 days	600 mg qd x 10 days	13	↓ 13% (5-20%)	↓ 12% (4-19%)	↓ 14% (2-24%)[b]
Tenofovir[d]	300 mg qd	600 mg qd x 14 days	30	↔	↔	↔
Boceprevir	800 mg tid x 6 days	600 mg qd x 16 days	NA	↑11% (2-20%)	↑ 20% (15-26%)	NA
Simeprevir	150 mg qd x 14 days	600 mg qd x 14 days	23	↔	↓ 10% (5-15%)	↓ 13% (7-19%)
Azithromycin	600 mg single dose	400 mg qd x 7 days	14	↔	↔	↔
Clarithromycin	500 mg q12h x 7 days	400 mg qd x 7 days	12	↑ 11% (3-19%)	↔	↔
Fluconazole	200 mg x 7 days	400 mg qd x 7 days	10	↔	↑ 16% (6-26%)	↑ 22% (5-41%)
Itraconazole	200 mg q12h x 14 days	600 mg qd x 28 days	16	↔	↔	↔
Rifabutin	300 mg qd x 14 days	600 mg qd x 14 days	11	↔	↔	↓ 12% (↓ 24-↑ 1%)
Rifampin	600 mg x 7 days	600 mg qd x 7 days	12	↓ 20% (11-28%)	↓ 26% (15-36%)	↓ 32% (15-46%)
Voriconazole	400 mg po q12h x 1 day, then 200 mg po q 12 h x 8 days	400 mg qd x 9 days	NA	↑ 38%[g]	↑ 44%[g]	NA
	300 mg po q12h days 2-7	300 mg qd x 7 days	NA	↓ 14%[f] (7-21%)	↔[f]	NA
	400 mg po q12h days 2-7	300 mg qd x 7 days	NA	↔[f]	↑ 17%[f] (6-29%)	NA
Artemether/ Lumefantrine	Artemether 20 mg/lumefantrine 120 mg tablets (6 4-tablet doses over 3 days)	600 mg qd × 26 days	12	↔	↓ 17%	NA
Atorvastatin	10 mg qd x 4 days	600 mg qd x 15 days	14	↔	↔	↔
Pravastatin	40 mg qd x 4 days	600 mg qd x 15 days	11	↔	↔	↔
Simvastatin	40 mg qd x 4 days	600 mg qd x 15 days	14	↓ 12% (↓ 28-↑ 8%)	↔	↓ 12% (↓ 25-↑ 3%)
Aluminum hydroxide 400 mg, magnesium hydroxide 400 mg, plus simethicone 40 mg	30 mL single dose	400 mg single dose	17	↔	↔	NA
Carbamazepine	200 mg qd x 3 days, 200 mg bid x 3 days, then 400 mg qd x 15 days	600 mg qd x 35 days	14	↓ 21% (15-26%)	↓ 36% (32-40%)	↓ 47% (41-53%)
Cetirizine	10 mg single dose	600 mg qd x 10 days	11	↔	↔	↔
Diltiazem	240 mg x 14 days	600 mg qd x 28 days	12	↑ 16% (6-26%)	↑ 11% (5-18%)	↑ 13% (1-26%)
Famotidine	40 mg single dose	400 mg single dose	17	↔	↔	NA
Paroxetine	20 mg qd x 14 days	600 mg qd x 14 days	12	↔	↔	↔
Sertraline	50 mg qd x 14 days	600 mg qd x 14 days	13	↑ 11% (6-16%)	↔	↔

↑ Indicates increase ↓ Indicates decrease ↔ Indicates no change or a mean increase or decrease of <10%.
[a] Parallel-group design; n for efavirenz + lopinavir/ritonavir, n for efavirenz alone.
[b] 95% CI.
[c] Soft Gelatin Capsule.
[d] Tenofovir disoproxil fumarate.
[e] 90% CI not available.
[f] Relative to steady-state administration of efavirenz (600 mg once daily for 9 days).
NA = not available.

12.4 Microbiology

Mechanism of Action

Efavirenz is an NNRTI of HIV-1. Efavirenz activity is mediated predominantly by noncompetitive inhibition of HIV-1 reverse transcriptase. HIV-2 reverse transcriptase and human cellular DNA polymerases α, β, γ, and δ are not inhibited by efavirenz.

Antiviral Activity in Cell Culture

The concentration of efavirenz inhibiting replication of wild-type laboratory adapted strains and clinical isolates in cell culture by 90 to 95% (EC_{90-95}) ranged from 1.7 to 25 nM in lymphoblastoid cell lines, peripheral blood mononuclear cells (PBMCs), and macrophage/monocyte cultures. Efavirenz demonstrated antiviral activity against clade B and most non-clade B isolates (subtypes A, AE, AG, C, D, F, G, J, N), but had reduced antiviral activity against group O viruses. Efavirenz demonstrated additive antiviral activity without cytotoxicity against HIV-1 in cell culture when combined with the NNRTIs delavirdine and nevirapine, NRTIs (abacavir, didanosine, emtricitabine, lamivudine, stavudine, tenofovir, zalcitabine, zidovudine), PIs (amprenavir, indinavir, lopinavir, nelfinavir, ritonavir, saquinavir), and the fusion inhibitor enfuvirtide. Efavirenz demonstrated additive to antagonistic antiviral activity in cell culture with atazanavir. Efavirenz was not antagonistic with adefovir, used for the treatment of hepatitis B virus infection, or ribavirin, used in combination with interferon for the treatment of hepatitis C virus infection.

Resistance

In cell culture

In cell culture, HIV-1 isolates with reduced susceptibility to efavirenz (>380-fold increase in EC_{90} value) emerged rapidly in the presence of drug. Genotypic characterization of these viruses identified single amino acid substitutions L100I or V179D, double substitutions L100I/V108I, and triple substitutions L100I/V179D/Y181C in reverse transcriptase.

Clinical studies

Clinical isolates with reduced susceptibility in cell culture to efavirenz have been obtained. One or more substitutions at amino acid positions 98, 100, 101, 103, 106, 108, 188, 190, 225, and 227 in reverse transcriptase were observed in patients failing treatment with efavirenz in combination with indinavir, or with zidovudine plus lamivudine. The K103N substitution was the most frequently observed. Long-term resistance surveillance (average 52 weeks, range 4 to 106 weeks) analyzed 28 matching baseline and virologic failure isolates. Sixty-one percent (17/28) of these failure isolates had decreased efavirenz susceptibility in cell culture with a median 88-fold change in efavirenz susceptibility (EC_{90} value) from reference. The most frequent NNRTI substitution to develop in these patient isolates was K103N (54%). Other NNRTI substitutions that developed included L100I (7%), K101E/Q/R (14%), V108I (11%), G190S/T/A (7%), P225H (18%), and M230I/L (11%).

Cross-Resistance

Cross-resistance among NNRTIs has been observed. Clinical isolates previously characterized as efavirenz-resistant were also phenotypically resistant in cell culture to delavirdine and nevirapine compared to baseline. Delavirdine- and/or nevirapine-resistant clinical viral isolates with NNRTI resistance-associated substitutions (A98G, L100I, K101E/P, K103N/S, V106A, Y181X, Y188X, G190X, P225H, F227L, or M230L) showed reduced susceptibility to efavirenz in cell culture. Greater than 90% of NRTI-resistant clinical isolates tested in cell culture retained susceptibility to efavirenz.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term carcinogenicity studies in mice and rats were carried out with efavirenz. Mice were dosed with 0, 25, 75, 150, or 300 mg/kg/day for 2 years. Incidences of hepatocellular adenomas and carcinomas and pulmonary alveolar/bronchiolar adenomas were increased above background in females. No increases in tumor incidence above background were seen in males. There was no NOAEL in females established for this study because tumor findings occurred at all doses. AUC at the NOAEL (150 mg/kg) in the males was approximately 0.9 times that in humans at the recommended clinical dose. In the rat study, no increases in tumor incidence were observed at doses up to 100 mg/kg/day, for which AUCs were 0.1 (males) or 0.2 (females) times those in humans at the recommended clinical dose.

Mutagenesis

Efavirenz tested negative in a battery of *in vitro* and *in vivo* genotoxicity assays. These included bacterial mutation assays in *S. typhimurium* and *E. coli*, mammalian mutation assays in Chinese hamster ovary cells, chromosome aberration assays in human peripheral blood lymphocytes or Chinese hamster ovary cells, and an *in vivo* mouse bone marrow micronucleus assay.

Impairment of Fertility

Efavirenz did not impair mating or fertility of male or female rats, and did not affect sperm of treated male rats. The reproductive performance of offspring born to female rats given efavirenz was not affected. The AUCs at the NOAEL values in male (200 mg/kg) and female (100 mg/kg) rats were approximately ≤0.15 times that in humans at the recommended clinical dose.

13.2 Animal Toxicology

Nonsustained convulsions were observed in 6 of 20 monkeys receiving efavirenz at doses yielding plasma AUC values 4- to 13-fold greater than those in humans given the recommended dose *[see Warnings and Precautions (5.10)]*.

14 CLINICAL STUDIES

14.1 Adults

Study 006, a randomized, open-label trial, compared efavirenz (600 mg once daily) + zidovudine (ZDV, 300 mg q12h) + lamivudine (LAM, 150 mg q12h) or efavirenz (600 mg once daily) + indinavir (IDV, 1000 mg q8h) with indinavir (800 mg q8h) + zidovudine (300 mg q12h) + lamivudine (150 mg q12h). Twelve hundred sixty-six patients (mean age 36.5 years [range 18 to 81], 60% Caucasian, 83% male) were enrolled. All patients were efavirenz-, lamivudine-, NNRTI-, and PI-naive at study entry. The median baseline CD4+ cell count was 320 cells/mm^3 and the median baseline HIV-1 RNA level was 4.8 log_{10} copies/mL. Treatment outcomes with standard assay (assay limit 400 copies/mL) through 48 and 168 weeks are shown in Table 9. Plasma HIV RNA levels were quantified with standard (assay limit 400 copies/mL) and ultrasensitive (assay limit 50 copies/mL) versions of the AMPLICOR HIV-1 MONITOR assay. During the study, version 1.5 of the assay was introduced in Europe to enhance detection of non-clade B virus.

Table 9: Outcomes of Randomized Treatment Through 48 and 168 Weeks, Study 006

Outcome	Efavirenz + ZDV + LAM (n=422)		Efavirenz + IDV (n=429)		IDV + ZDV + LAM (n=415)	
	Week 48	Week 168	Week 48	Week 168	Week 48	Week 168
Responder[a]	69%	48%	57%	40%	50%	29%
Virologic failure[b]	6%	12%	15%	20%	13%	19%
Discontinued for adverse events	7%	8%	6%	8%	16%	20%
Discontinued for other reasons[c]	17%	31%	22%	32%	21%	32%
CD4+ cell count (cells/mm^3)						
Observed subjects (n)	(279)	(205)	(256)	(158)	(228)	(129)
Mean change from baseline	190	329	191	319	180	329

[a] Patients achieved and maintained confirmed HIV-1 RNA <400 copies/mL through Week 48 or Week 168.
[b] Includes patients who rebounded, patients who were on study at Week 48 and failed to achieve confirmed HIV-1 RNA <400 copies/mL at time of discontinuation, and patients who discontinued due to lack of efficacy.
[c] Includes consent withdrawn, lost to follow-up, noncompliance, never treated, missing data, protocol violation, death, and other reasons. Patients with HIV-1 RNA levels <400 copies/mL who chose not to continue in the voluntary extension phases of the study were censored at date of last dose of study medication.

For patients treated with efavirenz + zidovudine + lamivudine, efavirenz + indinavir, or indinavir + zidovudine + lamivudine, the percentage of responders with HIV-1 RNA <50 copies/mL was 65%, 50%, and 45%, respectively, through 48 weeks, and 43%, 31%, and 23%, respectively, through 168 weeks. A Kaplan-Meier analysis of time to loss of virologic response (HIV RNA <400 copies/mL) suggests that both the trends of virologic response and differences in response continue through 4 years.

ACTG 364 is a randomized, double-blind, placebo-controlled, 48-week study in NRTI-experienced patients who had completed two prior ACTG studies. One-hundred ninety-six patients (mean age 41 years [range 18 to 76], 74% Caucasian, 88% male) received NRTIs in combination with efavirenz (600 mg once daily), or nelfinavir (NFV, 750 mg three times daily), or efavirenz (600 mg once daily) + nelfinavir in a randomized, double-blinded manner. The mean baseline CD4+ cell count was 389 cells/mm^3 and mean baseline HIV-1 RNA level was 8130 copies/mL. Upon entry into the study, all patients were assigned a new open-label NRTI regimen, which was dependent on their previous NRTI treatment experience. There was no significant difference in the mean CD4+ cell count among treatment groups; the overall mean increase was approximately 100 cells at 48 weeks among patients who continued on study regimens. Treatment outcomes are shown in Table 10. Plasma HIV RNA levels were quantified with the AMPLICOR HIV-1 MONITOR assay using a lower limit of quantification of 500 copies/mL.

Table 10: Outcomes of Randomized Treatment Through 48 Weeks, Study ACTG 364*

Outcome	Efavirenz + NFV + NRTIs (n=65)	Efavirenz + NRTIs (n=65)	NFV + NRTIs (n=66)
HIV-1 RNA <500 copies/mL[a]	71%	63%	41%
HIV-1 RNA ≥500 copies/mL[b]	17%	34%	54%
CDC Category C Event	2%	0%	0%
Discontinuations for adverse events[c]	3%	3%	5%
Discontinuations for other reasons[d]	8%	0%	0%

* For some patients, Week 56 data were used to confirm the status at Week 48.
[a] Subjects achieved virologic response (two consecutive viral loads <500 copies/mL) and maintained it through Week 48.
[b] Includes viral rebound and failure to achieve confirmed <500 copies/mL by Week 48.
[c] See *Adverse Reactions (6.1)* for a safety profile of these regimens.
[d] Includes loss to follow-up, consent withdrawn, noncompliance.

A Kaplan-Meier analysis of time to treatment failure through 72 weeks demonstrates a longer duration of virologic suppression (HIV RNA <500 copies/mL) in the efavirenz-containing treatment arms.

14.2 Pediatric Patients

Study AI266922 is an open-label study to evaluate the pharmacokinetics, safety, tolerability, and antiviral activity of efavirenz in combination with didanosine and emtricitabine in antiretroviral-naive and -experienced pediatric patients. Thirty-seven patients 3 months to 6 years of age (median 0.7 years) were treated with efavirenz. At baseline, median plasma HIV-1 RNA was 5.88 log_{10} copies/mL, median CD4+ cell count was 1144 cells/mm^3, and median CD4+ percentage was 25%. The median time on study therapy was 60 weeks; 27% of patients discontinued before Week 48. Using an ITT analysis, the overall proportions of patients with HIV RNA <400 copies/mL and <50 copies/mL at Week 48 were 57% (21/37) and 46% (17/37), respectively. The median increase from baseline in CD4+ count at 48 weeks was 196 cells/mm^3 and the median increase in CD4+ percentage was 6%.

Study PACTG 1021 was an open-label study to evaluate the pharmacokinetics, safety, tolerability, and antiviral activity of efavirenz in combination with didanosine and emtricitabine in pediatric patients who were antiretroviral therapy naive. Forty-three patients 3 months to 21 years of age (median 9.6 years) were dosed with efavirenz. At baseline, median plasma HIV-1 RNA was 4.8 log_{10} copies/mL, median CD4+ cell count was 367 cells/mm^3, and median CD4+ percentage was 18%. The median time on study therapy was 181 weeks; 16% of patients discontinued before Week 48. Using an ITT analysis, the overall proportions of patients with HIV RNA <400 copies/mL and <50 copies/mL at Week 48 were 77% (33/43) and 70% (30/43), respectively. The median increase from baseline in CD4+ count at 48 weeks of therapy was 238 cells/mm^3 and the median increase in CD4+ percentage was 13%.

Study PACTG 382 was an open-label study to evaluate the pharmacokinetics, safety, tolerability, and antiviral activity of efavirenz in combination with nelfinavir and an NRTI in antiretroviral-naive and NRTI-experienced pediatric patients. One hundred two patients 3 months to 16 years of age (median 5.7 years) were treated with efavirenz. Eighty-seven percent of patients had received prior antiretroviral therapy. At baseline, median plasma HIV-1 RNA was 4.57 log_{10} copies/mL, median CD4+ cell count was 755 cells/mm^3, and median CD4+ percentage was 30%. The median time on study therapy was 118 weeks; 25% of patients discontinued before Week 48. Using an ITT analysis, the overall proportion of patients with HIV RNA <400 copies/mL and <50 copies/mL at Week 48 were 57% (58/102) and 43% (44/102), respectively. The median increase from baseline in CD4+ count at 48 weeks of therapy was 128 cells/mm^3 and the median increase in CD4+ percentage was 5%.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 Capsules

Efavirenz capsules USP are available as follows:

Efavirenz capsules USP 50 mg are Yellow/White size '4' hard gelatin capsules imprinted with 'D' on yellow cap and '72' on white body with black edible ink filled with white to off-white colored powder.

Bottles of 30	NDC 65862-104-30
3 x 10 Unit-dose Capsules	NDC 65862-104-10

Efavirenz capsules USP 100 mg are White/White size '2' hard gelatin capsules imprinted with 'D' on white cap and '71' on white body with black edible ink filled with white to off-white colored powder.

Bottles of 30	NDC 65862-105-30
3 x 10 Unit-dose Capsules	NDC 65862-105-10

Efavirenz capsules USP 200 mg are Yellow/Yellow size 'OEL' hard gelatin capsules imprinted with 'D' on yellow cap and '36' on yellow body with black edible ink filled with white to off-white colored powder.

Bottles of 30	NDC 65862-106-30
Bottles of 90	NDC 65862-106-90
3 x 10 Unit-dose Capsules	NDC 65862-106-10
9 x 10 Unit-dose Capsules	NDC 65862-106-09

16.3 Storage

Efavirenz capsules should be stored at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information and Instructions for Use).

Drug Interactions

A statement to patients and healthcare providers is included on the product's bottle labels: *ALERT: Find out about medicines that should NOT be taken with Efavirenz Capsules.*

Efavirenz may interact with some drugs; therefore, advise patients to report to their doctor the use of any other prescription or nonprescription medication.

General Information for Patients

Inform patients that efavirenz is not a cure for HIV-1 infection and patients may continue to experience illnesses associated with HIV-1 infection, including opportunistic infections. Patients should remain under the care of a physician while taking efavirenz.

Advise patients to avoid doing things that can spread HIV-1 infection to others.

- **Do not share or reuse needles or other injection equipment.**
- **Do not share personal items that can have blood or body fluids on them, like toothbrushes and razor blades.**
- **Do not have any kind of sex without protection.** Always practice safer sex by using a latex or polyurethane condom to lower the chance of sexual contact with semen, vaginal secretions, or blood.
- **Do not breastfeed.** Mothers with HIV-1 should not breastfeed because HIV-1 can be passed to the baby in breast milk.

Dosing Instructions

Advise patients to take efavirenz every day as prescribed. If a patient forgets to take efavirenz, tell the patient to take the missed dose right away, unless it is almost time for the next dose. Advise the patient not to take 2 doses at one time and to take the next dose at the regularly scheduled time. Advise the patient to ask a healthcare provider if he/she needs help in planning the best times to take his/her medicine.

Efavirenz must always be used in combination with other antiretroviral drugs. Advise patients to take efavirenz on an empty stomach, preferably at bedtime. Taking efavirenz with food increases efavirenz concentrations and may increase the frequency of adverse reactions. Dosing at bedtime may improve the tolerability of nervous system symptoms *[see Dosage and Administration (2)* and *Adverse Reactions (6.1)]*. Healthcare providers should assist parents or caregivers in determining the best efavirenz dosing schedule for infants and young children.

For adult and pediatric patients who cannot swallow capsules, patients or their caregivers should be advised to read and carefully follow the instructions for administering the capsule contents in a small amount of food or infant formula *[see Dosage and Administration (2.3)* and *FDA-approved patient labeling (Patient Information* and *Instructions for Use)]*. Patients should call their healthcare provider or pharmacist if they have any questions.

Nervous System Symptoms

Inform patients that central nervous system symptoms (NSS) including dizziness, insomnia, impaired concentration, drowsiness, and abnormal dreams are commonly reported during the first weeks of therapy with efavirenz *[see Warnings and Precautions (5.6)]*. Dosing at bedtime may improve the tolerability of these symptoms, which are likely to improve with continued therapy. Alert patients to the potential for additive effects when efavirenz is used concomitantly with alcohol or psychoactive drugs. Instruct patients that if they experience NSS they should avoid potentially hazardous tasks such as driving or operating machinery.

Inform patients that there is a risk of developing late-onset neurotoxicity, including ataxia and encephalopathy which may occur months to years after beginning efavirenz therapy *[see Warnings and Precautions (5.6)]*.

Psychiatric Symptoms

Inform patients that serious psychiatric symptoms including severe depression, suicide attempts, aggressive behavior, delusions, paranoia, psychosis-like symptoms and catatonia have been reported in patients receiving efavirenz *[see Warnings and Precautions (5.5)]*. If they experience severe psychiatric adverse experiences they should seek immediate medical evaluation. Advise patients to inform their physician of any history of mental illness or substance abuse.

Rash

Inform patients that a common side effect is rash *[see Warnings and Precautions (5.8)]*. Rashes usually go away without any change in treatment. However, since rash may be serious, advise patients to contact their physician promptly if rash occurs.

Hepatotoxicity

Inform patients to watch for early warning signs of liver inflammation or failure, such as fatigue, weakness, lack of appetite, nausea and vomiting, as well as later signs such as jaundice, confusion, abdominal swelling, and discolored feces, and to consult their health care professional without delay if such symptoms occur *[see Warnings and Precautions (5.9) and Adverse Reactions (6.1)]*.

Females of Reproductive Potential

Advise females of reproductive potential to use effective contraception as well as a barrier method during treatment with efavirenz and for 12 weeks after discontinuing efavirenz. Advise patients to contact their healthcare provider if they plan to become pregnant, become pregnant, or if pregnancy is suspected during treatment with efavirenz *[see Warnings and Precautions (5.7)* and *Use in Specific Populations (8.1, 8.3)]*.

Pregnancy Exposure Registry

Advise patients that there is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to efavirenz during pregnancy *[see Use in Specific Populations (8.1)]*.

Fat Redistribution

Inform patients that redistribution or accumulation of body fat may occur in patients receiving antiretroviral therapy and that the cause and long-term health effects of these conditions are not known *[see Warnings and Precautions (5.13)]*.

Dispense with Patient Information available at: www.aurobindousa.com/product-medication-guides

Patient Information
Efavirenz Capsules USP
(eh-FAV-ir-enz)

Important: Ask your doctor or pharmacist about medicines that should not be taken with efavirenz capsules. For more information, see the section **"What should I tell my doctor before taking efavirenz capsules?"**

Read this Patient Information before you start taking efavirenz capsules and each time you get a refill. There may be new information. This information does not take the place of talking with your doctor about your medical condition or treatment.

What are efavirenz capsules?

Efavirenz capsules are a prescription HIV-1 (Human Immunodeficiency Virus type 1) medicine used with other antiretroviral medicines to treat HIV-1 infection in adults and in children who are at least 3 months old and who weigh at least 7 pounds 12 ounces (3.5 kg). HIV is the virus that causes AIDS (Acquired Immune Deficiency Syndrome).

It is not known if efavirenz capsules are safe and effective in children younger than 3 months of age or who weigh less than 7 pounds 12 ounces (3.5 kg).

When used with other antiretroviral medicines to treat HIV-1 infection, efavirenz capsules may help:

- reduce the amount of HIV-1 in your blood. This is called viral load.
- increase the number of CD4+ (T) cells in your blood that help fight off other infections.

Reducing the amount of HIV-1 and increasing the CD4+ (T) cells in your blood may help improve your immune system. This may reduce your risk of death or getting infections that can happen when your immune system is weak (opportunistic infections).

Efavirenz capsules do not cure HIV-1 infection or AIDS. You should keep taking HIV-1 medicines to control HIV-1 infection and decrease HIV-related illnesses.

Avoid doing things that can spread HIV-1 infection to others:

- Do not share or reuse needles or other injection equipment.
- Do not share personal items that can have blood or body fluids on them, like toothbrushes and razor blades.
- Do not have any kind of sex without protection. Always practice safer sex by using a latex or polyurethane condom to lower the chance of sexual contact with any body fluids such as semen, vaginal secretions, or blood.

Ask your doctor if you have any questions about how to prevent passing HIV to other people.

Who should not take efavirenz capsules?

Do not take efavirenz capsules if you are allergic to efavirenz or any of the ingredients in efavirenz capsules. See the end of this leaflet for a complete list of ingredients in efavirenz capsules.

Do not take efavirenz capsules if you are currently taking elbasvir and grazoprevir (ZEPATIER®).

What should I tell my doctor before taking efavirenz capsules?

Before taking efavirenz capsules, tell your doctor if you have any medical conditions and in particular, if you:

- have a heart condition
- have ever had a mental health problem
- have ever used street drugs or large amounts of alcohol
- have liver problems, including hepatitis B or C virus infection
- have a history of seizures
- are pregnant or plan to become pregnant. Efavirenz capsules may harm your unborn baby. If you are able to become pregnant your healthcare provider should do a pregnancy test before you start efavirenz capsules. You should not become pregnant while taking efavirenz capsules and for 12 weeks after stopping treatment with efavirenz capsules.

  Females who are able to become pregnant should use 2 effective forms of birth control during treatment and for 12 weeks after stopping treatment with efavirenz capsules. A barrier form of birth control should always be used along with another type of birth control.
  - Barrier forms of birth control may include latex or polyurethane condom, contraceptive sponge, diaphragm with spermicide, and cervical cap.
  - **Hormonal forms of birth control, such as birth control pills, injections, vaginal rings, or implants may not work during treatment with efavirenz capsules.**
  - Talk to your doctor about forms of birth control that may be used during treatment with efavirenz capsules.
  - **Pregnancy Registry.** There is a pregnancy registry for women who take antiretroviral medicines during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk to your doctor about how you can take part in this registry.
  - **Do not breastfeed if you take efavirenz capsules.**
  - You should not breastfeed if you have HIV because of the risk of passing HIV to your baby.

Tell your doctor and pharmacist about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Efavirenz capsules may affect the way other medicines work, and other medicines may affect how efavirenz capsules work, and may cause serious side effects. If you take certain medicines with efavirenz capsules, the amount of efavirenz in your body may be too low and it may not work to help control your HIV infection. The HIV virus in your body may become resistant to efavirenz or other HIV medicines that are like it.

You should not take efavirenz capsules if you take ATRIPLA (efavirenz, emtricitabine, tenofovir disoproxil fumarate) unless your doctor tells you to.

Tell your doctor and pharmacist about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Some medicines interact with efavirenz capsules.

Keep a list of your medicines to show your doctor and pharmacist.

- You can ask your doctor or pharmacist for a list of medicines that interact with efavirenz capsules.
- **Do not start taking a new medicine without telling your doctor.** Your doctor can tell you if it is safe to take efavirenz capsules with other medicines.

How should I take efavirenz capsules?

- Take efavirenz capsules exactly as your doctor tells you to.
- Do not change your dose or stop taking efavirenz capsules unless your doctor tells you to.
- Stay under the care of your doctor during treatment with efavirenz capsules.
- Efavirenz capsules must be used with other antiretroviral medicines.
- Take efavirenz capsules 1 time each day.
- Efavirenz comes as capsules.
- Swallow efavirenz capsules whole with liquid.

How and when to take efavirenz capsules.

- You should take efavirenz capsules on an empty stomach at bedtime. Taking efavirenz capsules with food increases the amount of medicine in your body. Some side effects may bother you less if you take efavirenz capsules on an empty stomach and at bedtime.
- Your child's doctor will prescribe the right dose of efavirenz capsules based on your child's weight.
- If you have difficulty swallowing capsules, tell your doctor. Your doctor may recommend opening the efavirenz capsule and mixing the contents with food or infant formula. **See the detailed "Instructions for Use" at the end of this Patient Information to learn the right way to take efavirenz capsules using the capsule sprinkle method.**
- Adults and children who take efavirenz capsules using the capsule sprinkle method should not eat for 2 hours after taking a dose of efavirenz capsules.
- Babies should not be given infant formula for 2 hours after taking a dose of efavirenz capsules using the capsule sprinkle method.
- Do not miss a dose of efavirenz capsules. If you forget to take efavirenz capsules, take the missed dose right away, unless it is almost time for your next dose. Do not take 2 doses at one time. Just take your next dose at your regularly scheduled time. If you need help in planning the best times to take your medicine, ask your doctor or pharmacist.
- If you take too much efavirenz, call your doctor or go to the nearest hospital emergency room right away.
- When your efavirenz capsules supply starts to run low, get more from your doctor or pharmacy. It is important not to run out of efavirenz capsules. The amount of HIV-1 in your blood may increase if the medicine is stopped for even a short time. The virus may become resistant to efavirenz and harder to treat.

What are the possible side effects of efavirenz capsules?

Efavirenz capsules may cause serious side effects, including:

- **Serious mental health problems** can happen in people who take efavirenz capsules. Tell your doctor right away if you have any of the following symptoms:
  - feel sad or hopeless
  - feel anxious or restless
  - have thoughts of hurting yourself (suicide) or have tried to hurt yourself or others
  - are not able to tell the difference between what is true or real and what is false or unreal
  - do not trust other people
  - hear or see things that are not real
  - are not able to move or speak normally
- **Nervous system symptoms** are common in people who take efavirenz capsules and can be severe. These symptoms usually begin during the first or second day of treatment with efavirenz capsules and usually go away after 2 to 4 weeks of treatment. Some symptoms may occur months to years after beginning efavirenz capsules therapy. These symptoms may become worse if you drink alcohol, take a medicine for mental health problems, or use certain street drugs during treatment with efavirenz capsules. Symptoms may include:
  - dizziness
  - trouble sleeping
  - trouble concentrating
  - drowsiness
  - unusual dreams
  - lack of coordination or difficulty with balance

  If you have dizziness, trouble concentrating or drowsiness, do not drive a car, use machinery, or do anything that needs you to be alert.

  Some nervous system symptoms (e.g., confusion, slow thoughts and physical movement, and delusions [false beliefs] or hallucinations [seeing or hearing things that others do not see or hear]) may occur months to years after beginning efavirenz capsules therapy. Promptly contact your health care provider should any of these symptoms occur.
- **Skin rash** is common with efavirenz capsules but can sometimes be severe. Skin rash usually goes away without any change in treatment. If you develop a rash with any of the following symptoms, tell your doctor right away:
  - skin rash, with or without itching
  - fever
  - swelling of your face
  - blisters or skin lesions
  - peeling skin
  - mouth sores
  - red or inflamed eyes, like "pink eye" (conjunctivitis)
- **Liver problems, including liver failure and death** can happen in people who take efavirenz capsules. Liver problems can happen in people without a history of liver problems. Your doctor will do blood tests to check your liver before you start efavirenz capsules and during treatment. Tell your doctor right away if you get any of the following symptoms:
  - your skin or the white part of your eyes turns yellow (jaundice)
  - your urine turns dark
  - your bowel movements (stools) turn light in color
  - you don't feel like eating food for several days or longer
  - you feel sick to your stomach (nausea)
  - you have lower stomach area (abdominal) pain
- **Seizures** can happen in people who take efavirenz capsules. Seizures are more likely to happen if you have had seizures in the past. Tell your doctor if you have had a seizure or if you take a medicine to help prevent seizures.
- **Changes in your immune system (Immune Reconstitution Syndrome)** can happen when you start taking HIV-1 medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your doctor if you start having new symptoms after starting your HIV-1 medicine.
- **Changes in body fat** can happen in people who take HIV-1 medicine. These changes may include increased amount of fat in the upper back and neck ("buffalo hump"), breast, and around the main part of your body (trunk). Loss of fat from the legs, arms, and face may also happen. The cause and long-term health effects of these conditions are not known.

The most common side effects of efavirenz capsules include:

- rash
- dizziness
- nausea
- headache
- difficulty concentrating
- abnormal dreams
- tiredness
- trouble sleeping
- vomiting

Some patients taking efavirenz capsules have experienced increased levels of lipids (cholesterol and triglycerides) in the blood. Tell your doctor if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of efavirenz capsules. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store efavirenz capsules?

- Store efavirenz capsules at room temperature between 68° to 77°F (20° to 25°C).

Keep efavirenz capsules and all medicines out of the reach of children.

General information about efavirenz capsules

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use efavirenz capsules for a cond