504-2018-11

EFAVIRENZ tablets
600 mg
Rx Only

2660624

Space for
2D Code

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use EFAVIRENZ TABLETS safely and effectively. See full prescribing information for EFAVIRENZ TABLETS.

EFAVIRENZ tablets for oral use

Initial U.S. Approval: 1998

----------RECENT MAJOR CHANGES----------

Dosage and Administration, Hepatic Function (2.1)	10/2017
Contraindications, Antiviral Agents (4)	10/2017
Warnings and Precautions, Psychiatric Symptoms (5.5)	01/2017
Warnings and Precautions, Hepatotoxicity (5.9)	10/2017

----------INDICATIONS AND USAGE----------

Efavirenz is a non-nucleoside reverse transcriptase inhibitor indicated in combination with other antiretroviral agents for the treatment of human immunodeficiency virus type 1 infection in adults and in pediatric patients atleast 3 months old and weighing at least 3.5 kg. (1)

----------DOSAGE AND ADMINISTRATION----------

- Efavirenz tablets should be taken orally once daily on an empty stomach, preferably at bedtime. (2)
- Recommended adult dose: 600 mg. (2.2)
- Pediatric dosing is based on weight. (2.3)

----------DOSAGE FORMS AND STRENGTHS----------

- Tablets: 600 mg (3)

----------CONTRAINDICATIONS----------

- Patients with previously demonstrated hypersensitivity (eg, Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to any of the components of this product. (4)
- Coadministration of efavirenz with elbasvir/grazoprevir.

----------WARNINGS AND PRECAUTIONS----------

- *QTc prolongation:* Consider alternatives to efavirenz in patients taking other medications with a known risk of Torsade de Pointes or in patients at higher risk of Torsade de Pointes. (5.2)
- *Do not use as a single agent or add on as a sole agent to a failing regimen.* Consider potential for cross-resistance when choosing other agents. (5.3)
- Not recommended with ATRIPLA, which contains efavirenz, emtricitabine, and tenofovir disoproxil fumarate, unless needed for dose adjustment when coadministered with rifampin. (5.4)
- *Serious psychiatric symptoms:* Immediate medical evaluation is recommended for serious psychiatric symptoms such as severe depression or suicidal ideation. (5.5, 17)
- *Nervous system symptoms* (NSS): NSS are frequent and usually begin 1 to 2 days after initiating therapy and resolve in 2 to 4 weeks. Dosing at bedtime may improve tolerability. NSS are not predictive of onset of psychiatric symptoms. (5.6, 6.1, 17)
- *Embryo-Fetal Toxicity:* Avoid administration in the first trimester of pregnancy as fetal harm may occur. (5.7, 8.1)
- *Hepatotoxicity:* Monitor liver function tests before and during treatment in patients with underlying hepatic disease, including hepatitis B or C infection, marked transaminase elevations, or who are taking medications associated with serious liver toxicity. Among reported cases of hepatic failure, a few occurred in patients with no pre-existing hepatic disease. (5.9, 6.1, 8.6)
- *Rash:* Rash usually begins within 1 to 2 weeks after initiating therapy and resolves within 4 weeks. Discontinue if severe rash develops. (5.8, 6.1, 17)
- *Convulsions:* Use caution in patients with a history of seizures. (5.10)
- *Lipids:* Total cholesterol and triglyceride elevations. Monitor before therapy and periodically thereafter. (5.11)
- *Immune reconstitution syndrome:* May necessitate further evaluation and treatment. (5.12)
- *Redistribution/accumulation of body fat:* Observed in patients receiving antiretroviral therapy. (5.13, 17)

----------ADVERSE REACTIONS----------

Most common adverse reactions (>5%, moderate-severe) are impaired concentration, abnormal dreams, rash, dizziness, nausea, headache, fatigue, insomnia, and vomiting. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Hetero Labs Limited at 1-866-495-1995 or FDA at 1-800-FDA-1088 or *www.fda.gov/medwatch*.

----------DRUG INTERACTIONS----------

- *Coadministration of efavirenz can alter the concentrations of other drugs and other drugs may alter the* concentrations of efavirenz. The potential for drug-drug interactions should be considered before and during therapy. (7)

----------USE IN SPECIFIC POPULATIONS----------

- *Lactation:* Breastfeeding not recommended. (8.2)
- *Females and Males of Reproductive Potential:* Pregnancy testing and contraception are recommended. (8.3)
- *Hepatic impairment:* Efavirenz tablets are not recommended for patients with moderate or severe hepatic impairment. Use caution in patients with mild hepatic impairment. (8.6)
- *Pediatric patients:* The incidence of rash was higher than in adults. (5.8, 6.2, 8.4)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 11/2018

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE
2 DOSAGE AND ADMINISTRATION
2.1 Hepatic Function
2.2 Adults
2.3 Pediatric Patients
3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
5 WARNINGS AND PRECAUTIONS
5.1 Drug Interactions
5.2 QTc Prolongation
5.3 Resistance
5.4 Coadministration with Related Products
5.5 Psychiatric Symptoms
5.6 Nervous System Symptoms
5.7 Embryo-Fetal Toxicity
5.8 Rash
5.9 Hepatotoxicity
5.10 Convulsions
5.11 Lipid Elevations
5.12 Immune Reconstitution Syndrome
5.13 Fat Redistribution
6 ADVERSE REACTIONS
6.1 Clinical Trials Experience
6.2 Postmarketing Experience
7 DRUG INTERACTIONS
7.1 Potential for Efavirenz to Affect other Drugs
7.2 Potential for Other Drugs to Affect Efavirenz
7.3 QT Prolonging Drugs
7.4 Established and Other Potentially Significant Drug Interactions
7.5 Drugs Without Clinically Significant Interactions with Efavirenz
7.6 Cannabinoid Test Interaction
8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.2 Lactation
8.3 Females and Males of Reproductive Potential
8.4 Pediatric Use
8.5 Geriatric Use
8.6 Hepatic Impairment
10 OVERDOSAGE
11 DESCRIPTION
12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.2 Pharmacodynamics
12.3 Pharmacokinetics
12.4 Microbiology
13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
13.2 Animal Toxicology
14 CLINICAL STUDIES
14.1 Adults
14.2 Pediatric Patients
16 HOW SUPPLIED/STORAGE AND HANDLING
16.2 Tablets
16.3 Storage
17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Efavirenz tablets in combination with other antiretroviral agents is indicated for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in adults and in pediatric patients at least 3 months old and weighing at least 3.5 kg.

2 DOSAGE AND ADMINISTRATION

2.1 Hepatic Function

Monitor hepatic function prior to and during treatment with efavirenz tablets *[see Warnings and Precautions (5.9)]*.

Efavirenz tablets are not recommended in patients with moderate or severe hepatic impairment (Child Pugh B or C) *[see Warnings and Precautions (5.9) and Use in Specific Populations (8.6)]*.

2.2 Adults

The recommended dosage of efavirenz tablets are 600 mg orally, once daily, in combination with a protease inhibitor and/or nucleoside analogue reverse transcriptase inhibitors (NRTIs). It is recommended that efavirenz tablets be taken on an empty stomach, preferably at bedtime. The increased efavirenz concentrations observed following administration of efavirenz tablets with food may lead to an unusual manifestation or the use of efavirenz *[see Clinical Pharmacology (12.3)]*. Dosing at bedtime may improve the tolerability of nervous system symptoms *[see Warnings and Precautions (5.6), Adverse Reactions (6.1), and Patient Counseling Information (17)]*. Efavirenz tablets should be swallowed intact with liquid.

Concomitant Antiretroviral Therapy

Efavirenz tablets must be given in combination with other antiretroviral medications *[see Indications and Usage (1), Warnings and Precautions (5.3), Drug Interactions (7.1), and Clinical Pharmacology (12.3)]*.

2.3 Pediatric Patients

It is recommended that efavirenz tablets be taken on an empty stomach, preferably at bedtime. Table 1 describes the recommended dose of efavirenz tablets for pediatric patients 3 months of age or older and weighing between 3.5 kg and 40 kg *[see Clinical Pharmacology (12.3)]*. The recommended dosage of efavirenz tablets for pediatric patients weighing 40 kg or greater is 600 mg once daily.

Table 1: Efavirenz Tablets Dosing in Pediatric Patients

Patient Body Weight	Efavirenz Tablets Daily Dose	Number of Tablets* and Strength to Administer
at least 40 kg	600 mg	one 600 mg tablet

* Tablets must not be crushed.

3 DOSAGE FORMS AND STRENGTHS

- Tablets

600 mg tablets are yellow, capsular-shaped, film-coated tablets, with 'H' on one side and '4' on the other side.

4 CONTRAINDICATIONS

- Efavirenz tablets are contraindicated in patients with previously demonstrated clinically significant hypersensitivity (eg, Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to any of the components of this product.
- Coadministration of efavirenz with elbasvir and grazoprevir is contraindicated *[see Warnings and Precautions (5.1) and Drug Interactions (7.1)]*.

5 WARNINGS AND PRECAUTIONS

5.1 Drug Interactions

Clinically significant drug interactions may be altered by substrates, inhibitors, or inducers of CYP3A. Likewise, efavirenz may alter plasma concentrations of drugs metabolized by CYP3A or CYP2B6. The most prominent effect of efavirenz at steady-state is induction of CYP3A and CYP2B6. *[See Dosage and Administration (2.2) and Drug Interactions (7.1)]*.

5.2 QTc Prolongation

QTc prolongation has been observed with the use of efavirenz *[see Drug Interactions (7.3, 7.4) and Clinical Pharmacology (12.2)]*. Consider alternatives to efavirenz tablets when coadministered with a drug with a known risk of Torsade de Pointes or when administered to patients at higher risk of Torsade de Pointes.

5.3 Resistance

Efavirenz tablets must not be used as a single agent to treat HIV-1 infection or added on as a sole agent to a failing regimen. Resistant virus emerges rapidly when efavirenz is administered as monotherapy. The choice of new antiretroviral agents to be used in combination with efavirenz should take into consideration the potential for viral cross-resistance.

5.4 Coadministration with Related Products

Coadministration of efavirenz tablets with ATRIPLA (efavirenz 600 mg/emtricitabine 200 mg/ tenofovir disoproxil fumarate 300 mg) is not recommended unless needed for dose adjustment (eg, with rifampin), since efavirenz is one of its active ingredients.

5.5 Psychiatric Symptoms

Serious psychiatric adverse experiences have been reported in patients treated with efavirenz tablets. In controlled trials of 1,008 patients who received treatment with efavirenz tablets for a median duration of 2 years and 635 patients treated with control regimens for a mean of 1.5 years, the frequency (regardless of causality) of specific serious psychiatric events among patients who received efavirenz tablets or control regimens, respectively, were severe depression (2.4%, 0.9%), suicidal ideation (0.7%, 0.3%), nonfatal suicide attempts (0.5%, 0), aggressive behavior (0.4%, 0.3%), paranoid reactions (0.4%, 0.3%), and manic reactions (0.2%, 0.3%). When psychiatric symptoms similar to those noted above were combined and evaluated as a group in a multifactorial analysis of data from these 1,008 patients, female gender, history of psychiatric illness, and longer duration of efavirenz tablets exposure were significant predictors of experiencing one of these symptoms. Other factors associated with an increase in the occurrence of these psychiatric symptoms were history of injection drug use, psychiatric history, and receipt of psychiatric medication at study entry; similar associations were observed in both the efavirenz tablets and control groups. Discontinuations of efavirenz tablets due to these psychiatric symptoms occurred throughout the study for both efavirenz tablets-treated and control-treated patients. Nine percent of efavirenz tablets-treated patients discontinued or interrupted treatment because of one or more of these selected psychiatric symptoms. There have also been occasional postmarketing reports of death by suicide, delusions, psychosis-like behavior and catatonia, although a causal relationship to the use of efavirenz tablets cannot be determined from these reports. Postmarketing cases of catatonia have also been reported and were associated with multiple concomitant medications. Patients should be informed that if they experience psychosis, they should seek immediate medical evaluation to assess the possibility that the symptoms may be related to the use of efavirenz tablets, and if so, to determine whether the risks of continued therapy outweigh the benefits. *[See Adverse Reactions (6.1)]*.

5.6 Nervous System Symptoms

Fifty-three percent (531/1,008) of patients receiving efavirenz tablets in controlled trials reported central nervous system symptoms (any grade, regardless of causality) compared to 25% (156/635) of patients receiving control regimens *[see Adverse Reactions (6.1)]*. In these symptoms included, but were not limited to, dizziness (8.8%, 4.7%), insomnia (4.1%, 2.0%), somnolence (2.2%, 0.8%), abnormal dreams (6.2%, 1.6%), hallucinations (1.2%, 0.3%), confusion (1.3%, 0.6%), and impaired concentration (3.2%, 0.9%). Overall, 2.1% of patients discontinued therapy as a result. These symptoms usually begin during the first or second day of therapy and generally resolve after the first 2 to 4 weeks of therapy. After 4 weeks of therapy, the prevalence of nervous system symptoms of at least moderate severity ranged from 5% to 9% in patients treated with regimens containing efavirenz tablets and from 3% to 5% in patients treated with a control regimen. Patients should be informed that these common symptoms were likely to improve with continued therapy and were not predictive of subsequent onset of the less frequent psychotic symptoms *[see Warnings and Precautions (5.5)]*. Dosing at bedtime may improve the tolerability of these nervous system symptoms *[see Dosage and Administration (2)]*.

Analysis of long-term data from Study 006 (median follow-up 180 weeks, 102 weeks, and 76 weeks for patients treated with efavirenz tablets + zidovudine + lamivudine, efavirenz tablets + indinavir, and indinavir + zidovudine + lamivudine, respectively) showed that, beyond 24 weeks of therapy, the incidences of new-onset nervous system symptoms among efavirenz tablets-treated patients were generally similar to those in the indinavir-containing control arm.

Patients receiving efavirenz tablets should be alerted to the potential for additive central nervous system effects when efavirenz tablets are used concomitantly with alcohol or psychoactive drugs.

Patients who experience central nervous system symptoms such as dizziness, impaired concentration, and/or drowsiness should avoid potentially hazardous tasks such as driving or operating machinery.

5.7 Embryo-Fetal Toxicity

Efavirenz may cause fetal harm when administered during the first trimester to a pregnant woman. Advise females of reproductive potential who are receiving efavirenz tablets to avoid pregnancy. *[See Use in Specific Populations (8.1 and 8.3)]*.

5.8 Rash

In controlled clinical trials, 26% (266/1,008) of adult patients treated with 600 mg efavirenz tablets experienced new-onset skin rash compared with 17% (111/635) of those treated in control groups *[see Adverse Reactions (6.1)]*. Rash associated with blistering, moist desquamation, or ulceration occurred in 0.9% (9/1,008) of patients in efavirenz tablets-containing regimens. The incidence of Stevens-Johnson syndrome and erythema multiforme in adult patients treated with efavirenz tablets in all studies was 0.1%. Rashes are usually mild-to-moderate maculopapular skin eruptions that occur within the first 2 weeks of initiating therapy with efavirenz (median time to onset of rash in adults was 11 days) and, in most patients continuing therapy with efavirenz, rash resolves within 1 month (median duration, 16 days). The discontinuation rate for rash in adult clinical trials was 1.7% (17/1,008).

Rash was reported in 59 of 182 pediatric patients (32%) treated with efavirenz tablets *[see Adverse Reactions (6.2)]*. Two pediatric patients experienced Grade 3 rash (confluent rash with fever, generalized rash), and four patients had Grade 4 rash (erythema multiforme). The median time to onset of rash in pediatric patients was 28 days (range 3 to 1,642 days). Prophylaxis with appropriate antihistamines before initiating therapy with efavirenz tablets in pediatric patients should be considered.

Efavirenz tablets can generally be reinitiated in patients interrupting therapy because of rash. Efavirenz tablets should be discontinued in patients developing severe rash associated with blistering, desquamation, mucosal involvement, or fever. Appropriate antihistamines and/or corticosteroids may improve the tolerability and hasten the resolution of rash. For patients who have had a life-threatening cutaneous reaction (eg, Stevens-Johnson syndrome), alternative therapy should be considered *[see also Contraindications (4)]*.

5.9 Hepatotoxicity

Postmarketing cases of hepatitis, including fulminant hepatitis progressing to liver failure requiring transplantation or resulting in death, have been reported in patients treated with efavirenz. Reports have included patients with underlying hepatic disease, including coinfection with hepatitis B or C, and patients without pre-existing hepatic disease or other identifiable risk factors.

Efavirenz is not recommended for patients with moderate or severe hepatic impairment. Careful monitoring is recommended for patients with mild hepatic impairment receiving efavirenz. *[see Adverse Reactions (6.1) and Use in Specific Populations (8.6)]*.

Monitoring of liver enzymes before and during treatment is recommended for all patients *[see Dosage and Administration (2.1)]*. Consider discontinuing efavirenz in patients with persistent elevations of serum transaminases to greater than five times the upper limit of the normal range.

Discontinue efavirenz if elevation of serum transaminases is accompanied by clinical signs or symptoms of hepatitis or hepatic decompensation.

5.10 Convulsions

Convulsions have been observed in adult and pediatric patients receiving efavirenz, generally in the presence of known medical history of seizures *[see Nonclinical Toxicology (13.2)]*. Caution should be taken in any patient with a history of seizures. Patients who are receiving concomitant anticonvulsant medications primarily metabolized by the liver, such as phenytoin and phenobarbital, may require periodic monitoring of plasma levels *[see Drug Interactions (7.1)]*.

5.11 Lipid Elevations

Treatment with efavirenz tablets has resulted in increases in the concentration of total cholesterol and triglycerides *[see Adverse Reactions (6.1)]*. Cholesterol and triglyceride testing should be performed before initiating efavirenz tablets therapy and at periodic intervals during therapy.

5.12 Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in patients treated with combination antiretroviral therapy, including efavirenz tablets. During the initial phase of combination antiretroviral treatment, patients whose immune system responds may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium* infection, cytomegalovirus, *Pneumocystis jiroveci* pneumonia (PCP), or tuberculosis), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis, and Guillain-Barré syndrome) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable, and can occur many months after initiation of treatment.

5.13 Fat Redistribution

Redistribution/accumulation of body fat including central obesity, dorsocervical fat enlargement (buffalo hump), peripheral wasting, facial wasting, breast enlargement, and "cushingoid appearance" have been observed in patients receiving antiretroviral therapy. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been established.

6 ADVERSE REACTIONS

The most significant adverse reactions observed in patients treated with efavirenz tablets are:

- psychiatric symptoms *[see Warnings and Precautions (5.5)]*,
- nervous system symptoms *[see Warnings and Precautions (5.6)]*,
- rash *[see Warnings and Precautions (5.8)]*,
- hepatotoxicity *[see Warnings and Precautions (5.9)]*

6.1 Clinical Trials Experience

Because clinical studies are conducted under widely varying conditions, the adverse reaction rates reported cannot be directly compared to rates in other clinical studies and may not reflect the rates observed in clinical practice.

Adverse Reactions in Adults

The most common (>5% in either efavirenz treatment group) adverse reactions of at least moderate severity in patients in Study 006 treated with efavirenz tablets + zidovudine/lamivudine or indinavir were rash, dizziness, nausea, headache, fatigue, insomnia, and vomiting.

Selected clinical adverse experiences and laboratory abnormalities observed in ≥2% of efavirenz tablets-treated patients in two controlled clinical trials are presented in Table 2.

Table 2: Selected Treatment-Emergent* Adverse Reactions of Moderate or Severe Intensity Reported in ≥2% of Efavirenz-Treated Patients in Studies 006 and ACTG 364

	Study 006 LAM-, NRTI-, and Protease Inhibitor-Naive Patients			Study ACTG 364 NRTI-experienced, NRTI-, and Protease Inhibitor-Naive Patients		
	Efavirenz tablets† + ZDV/LAM (n=412)	Efavirenz tablets† + Indinavir (n=415)	Indinavir + ZDV/LAM (n=401)	Efavirenz tablets† + NRTIs (n=64)	Efavirenz tablets† + NRTIs (n=65)	Nelfinavir + NRTIs (n=66)
Adverse Reactions	180 weeks‡	102 weeks‡	76 weeks‡	71.1 weeks‡	70.9 weeks‡	62.7 weeks‡
Body as a Whole						
Fatigue	8%	5%	9%	0	2%	3%
Pain	1%	2%	8%	13%	6%	17%
Central and Peripheral Nervous System						
Dizziness	9%	9%	2%	2%	6%	6%
Headache	8%	5%	3%	5%	2%	3%
Insomnia	7%	7%	2%	0	0	2%
Concentration impaired	5%	3%	<1%	0	0	0
Abnormal dreams	3%	1%	0	—	—	—
Somnolence	2%	2%	<1%	0	0	0
Amnesia	1%	<1%	<1%	0	2%	2%
Gastrointestinal						
Nausea	10%	6%	24%	3%	2%	2%
Vomiting	6%	3%	14%	—	—	—
Diarrhea	3%	5%	14%	3%	9%	9%
Dyspepsia	4%	4%	6%	0	0	2%
Abdominal pain	2%	2%	5%	3%	3%	3%
Psychiatric						
Anxiety	2%	4%	<1%	—	—	—
Depression	5%	4%	<1%	3%	0	5%
Nervousness	2%	2%	0	2%	0	2%
Skin & Appendages						
Rash§	11%	18%	5%	9%	5%	9%
Pruritus	<1%	1%	1%	9%	5%	9%

* Includes adverse events at least possibly related to study drug or of unknown relationship for Study 006. Includes all adverse events regardless of relationship to study drug for Study ACTG 364.

† Efavirenz tablets provided as 600 mg once daily.

‡ Median duration of treatment.

§ Includes pruritus, maculopapular rash, rash erythematous, rash follicular, rash maculopapular, rash petechial, rash pustular, and urticaria for Study 006 and macules, papules, rash, erythema, redness, inflammation, allergic rash, urticaria, welts, hives, itchy, and pruritus for ACTG 364.

— = Not Specified.

ZDV = zidovudine, LAM=lamivudine.

Pancreatitis has been reported, although a causal relationship with efavirenz has not been established. Asymptomatic increases in serum amylase levels were observed in a significantly higher number of patients treated with efavirenz 600 mg than in control patients *[see Laboratory Abnormalities]*.

Nervous System Symptoms

For 1,008 patients treated with regimens containing efavirenz tablets and 635 patients treated with a control regimen in controlled trials, Table 3 lists the frequency of symptoms of different degrees of severity and gives the discontinuation rates for one or more of the following nervous system symptoms: dizziness, insomnia, impaired concentration, somnolence, abnormal dreaming, euphoria, confusion, agitation, amnesia, hallucinations, stupor, abnormal thinking, and depersonalization that were considered drug-related (5.6). The frequencies of specific central and peripheral nervous system symptoms are provided in Table 3.

Table 3: Percent of Patients with One or More Selected Nervous System Symptoms[a,b]

Percent of Patients with:	Efavirenz Tablets 600 mg Once Daily (n=1,008)	Control Groups (n=635)
	%	%
Symptoms of any severity	52.7	24.6
Mild symptoms[c]	33.3	15.6
Moderate symptoms[d]	17.4	7.7
Severe symptoms[e]	2	1.3
Treatment discontinuation as a result of symptoms	2.1	1.1

[a] Includes events reported regardless of causality.

[b] Data from Study 006 and three Phase 2/3 studies.

[c] "Mild" = Symptoms which do not interfere with patient's daily activities.

[d] "Moderate" = Symptoms which may interfere with daily activities.

[e] "Severe" = Events which interrupt patient's usual daily activities.

Psychiatric Symptoms

Serious psychiatric adverse experiences have been reported in patients treated with efavirenz tablets. In controlled trials, psychiatric symptoms observed at a frequency greater than 2% among patients treated with efavirenz tablets or control regimens, respectively, were depression (19%, 16%), anxiety (13%, 9%), and nervousness (7%, 2%).

Rash

In controlled clinical trials, the frequency of rash (all grades, regardless of causality) was 26% for 1,008 adults treated with 600 mg efavirenz tablets and 17% for 635 adults treated with a control regimen. Most reports of rash were mild or moderate in severity. The frequency of Grade 3 rash was 6.8% for efavirenz tablets-treated patients and 3.0% for control patients. Grade 4 rash was 0.4% for each group. The discontinuation rate as a result of rash was 1.7% for efavirenz tablets-treated patients and 0.3% for control groups *[see Warnings and Precautions (5.8)]*.

Experience with efavirenz tablets in patients who discontinued other antiretroviral agents of the NNRTI class is limited. Of 49 patients who discontinued nevirapine because of rash and were then treated with efavirenz tablets, Nine of these patients developed mild-to-moderate rash while receiving therapy with efavirenz tablets, and two of these patients discontinued because of rash.

Laboratory Abnormalities

Selected Grade 3 to 4 laboratory abnormalities reported in ≥2% of efavirenz tablets-treated patients in two clinical trials are presented in Table 4.

Table 4: Selected Grade 3 to 4 Laboratory Abnormalities Reported in ≥2% of Efavirenz-Treated Patients in Studies 006 and ACTG 364

		Study 006 LAM-, NRTI-, and Protease Inhibitor-Naive Patients			Study ACTG 364 NRTI-experienced, NRTI-, and Protease Inhibitor-Naive Patients		
		Efavirenz tablets* +ZDV/LAM (n=412)	Efavirenz tablets* +Indinavir (n=415)	Indinavir +ZDV/LAM (n=401)	Efavirenz tablets* + Nelfinavir + NRTIs (n=64)	Efavirenz tablets* + NRTIs (n=65)	Nelfinavir + NRTIs (n=66)
Variable	Limit	180 weeks†	102 weeks†	76 weeks†	71.1 weeks†	70.9 weeks†	62.7 weeks†
Chemistry							
ALT	>5 x ULN	5%	8%	5%	2%	6%	3%
AST	>5 x ULN	5%	6%	5%	6%	8%	8%
GGT[‡]	>5 x ULN	8%	7%	3%	5%	0	5%
Amylase	>2 x ULN	4%	4%	1%	0	6%	2%
Glucose	>250 mg/dL	3%	3%	1%	2%	2%	3%
Triglycerides[§]	>751 mg/dL	9%	6%	6%	11%	8%	17%
Hematology							
Neutrophils	<750/mm³	10%	3%	5%	2%	3%	2%

* Efavirenz tablets provided as 600 mg once daily.

† Median duration of treatment.

‡ Isolated elevations of GGT in patients receiving efavirenz tablets may reflect enzyme induction not associated with liver toxicity.

§ Fasting.

ZDV = zidovudine, LAM = lamivudine, ULN = upper limit of normal, ALT = alanine aminotransferase, AST = aspartate aminotransferase, GGT = gamma glutamyltransferase.

Patients Coinfected with Hepatitis B or C

Liver function tests should be monitored in patients with a history of hepatitis B and/or C. In the long-term data from Study 006 (median follow-up 180 weeks), patients coinfected with hepatitis or chronic duration of therapy (48 weeks) and 84 treated with a control regimen (24 weeks) were comparable in the incidence of grade 3 or 4 hepatitis B surface antigen (HBsAg) positivity and/or hepatitis C antibody positivity. Among these coinfected patients, the incidence in AST is greater than five times ULN developed in 13% of patients in the efavirenz tablets arms and 7% of those in the control arm, and elevations in ALT to greater than five times ULN developed in 20% of patients in the efavirenz tablets arms and 7% of patients in the control arm. Among coinfected patients, 3% of those treated with efavirenz tablets-containing regimens and 2% in the control arm discontinued from the study because of liver or biliary system disorders *[see Warnings and Precautions (5.9)]*.

Lipids

Increases from baseline in total cholesterol of 10 to 20% have been observed in some uninfected volunteers receiving efavirenz. In patients treated with efavirenz tablets + zidovudine + lamivudine, increases from baseline in fasting total cholesterol and HDL of approximately 20% and 5%, respectively, were observed. In patients treated with efavirenz tablets + indinavir, increases from baseline in nonfasting cholesterol and HDL of approximately 40% and 35%, respectively, were observed. Nonfasting total triglycerides ≥240 mg/dL and ≥300 mg/dL were reported in 34% and 9%, respectively, of patients treated with efavirenz tablets + zidovudine + lamivudine; 54% and 20%, respectively, of patients treated with efavirenz tablets + indinavir; and 28% and 4%, respectively, of patients treated with indinavir + zidovudine + lamivudine. The effects of efavirenz tablets on triglycerides and LDL in this study were not well characterized since samples were taken from nonfasting patients. The clinical significance of these findings is unknown *[see Warnings and Precautions (5.11)]*.

Adverse Reactions in Pediatric Patients

Because clinical studies are conducted under widely varying conditions, the adverse reaction rates reported cannot be directly compared to rates in other clinical studies and may not reflect the rates observed in clinical practice.

Assessment of adverse reactions is based on three clinical trials in 182 HIV-1 infected pediatric patients (3 months to 21 years of age) who received efavirenz tablets in combination with other antiretroviral agents for a median of 120 weeks. The adverse reaction profile observed in the three trials was similar to those observed in clinical trials in adults except that rash was more common in pediatric patients (32% for all grades regardless of causality) and was severe in 4% of patients. Discontinuations for rash were 2.2%. Two pediatric patients had Grade 3 rash (confluent rash with fever, generalized rash), and two patients had Grade 4 rash (erythema multiforme). Five pediatric patients (2.7%) discontinued from the study because of rash *[see Warnings and Precautions (5.8)]*.

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of efavirenz tablets. Because these reactions are reported voluntarily from a population of unknown size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Body as a Whole: allergic reactions, asthenia, redistribution/accumulation of body fat *[see Warnings and Precautions (5.13)]*.

Central and Peripheral Nervous System: abnormal coordination, ataxia, cerebellar coordination and balance disturbances, convulsions, hypoesthesia, paresthesia, neuropathy, tremor, vertigo

Endocrine: gynecomastia

Gastrointestinal: constipation, malabsorption

Cardiovascular: palpitations, vasculitis

Liver and Biliary System: hepatic enzyme increase, hepatic failure, hepatitis.

Metabolic and Nutritional: hypercholesterolemia, hypertriglyceridemia

Musculoskeletal: arthralgia, myalgia, myopathy

Psychiatric: aggressive reactions, agitation, delusions, emotional lability, mania, neurosis, paranoia, psychosis, suicide, catatonia

Respiratory: dyspnea

Skin and Appendages: erythema multiforme, photoallergic dermatitis, Stevens-Johnson syndrome

Special Senses: abnormal vision, tinnitus

7 DRUG INTERACTIONS

7.1 Potential for Efavirenz to Affect Other Drugs

Efavirenz has been shown *in vivo* to induce CYP3A and CYP2B6. Other compounds that are substrates of CYP3A or CYP2B6 may have decreased plasma concentrations when coadministered with efavirenz tablets.

7.2 Potential for Other Drugs to Affect Efavirenz

Drugs that induce CYP3A activity (eg, phenobarbital, rifampin, rifabutin) would be expected to increase the clearance of efavirenz resulting in lowered plasma concentrations *[see Dosage and Administration (2.2)]*.

7.3 QT Prolonging Drugs

There is limited information available on the potential for a pharmacodynamic interaction between efavirenz and drugs that prolong the QTc interval. QTc prolongation has been observed with the use of efavirenz *[see Clinical Pharmacology (12.2)]*. Consider alternatives to efavirenz when coadministered with a drug with a known risk of Torsade de Pointes.

7.4 Established and Other Potentially Significant Drug Interactions

Drug interactions with efavirenz tablets are summarized in Tables 5. for pharmacokinetics data, *[see Clinical Pharmacology (12.3)]* Tables 7 and 8. This table is not all-inclusive.

Table 5: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies or Predicted Interaction

Concomitant Drug Class: Drug Name	Effect	Clinical Comment
HIV antiviral agents		
Protease inhibitor: Fosamprenavir calcium	↓ amprenavir	Fosamprenavir (unboosted): Appropriate doses of the combinations with respect to safety and efficacy have not been established. Fosamprenavir/ritonavir: An additional 100 mg/day (300 mg total) of ritonavir is recommended when efavirenz tablets are administered with fosamprenavir/ritonavir 700 mg/100 mg twice daily. Additional ritonavir is not required when efavirenz tablets are administered with fosamprenavir plus ritonavir twice daily.
Protease inhibitor: Atazanavir	↓ atazanavir*	*Treatment-naive patients:* When coadministered with efavirenz tablets, the dose of atazanavir should be increased to 400 mg with ritonavir 100 mg (together once daily with food) and efavirenz tablets 600 mg (once daily on an empty stomach, preferably at bedtime). *Treatment-experienced patients:* Coadministration of efavirenz tablets and atazanavir is not recommended.
Protease inhibitor: Indinavir	↓ indinavir*	The optimal dose of indinavir, when given in combination with efavirenz tablets, is not known. Increasing the indinavir dose to 1,000 mg every 8 hours does not compensate for the increased indinavir metabolism due to efavirenz tablets.
Protease inhibitor: Lopinavir/ritonavir	↓ lopinavir*	Lopinavir/ritonavir once daily dosing is not recommended when coadministered with efavirenz tablets. The dose of lopinavir/ritonavir must be increased when coadministered with efavirenz tablets. See the lopinavir/ritonavir prescribing information for dose recommendations. Dose increase may not adequately maintain plasma concentrations of both drugs in all patients.
Protease inhibitor: Ritonavir	↑ ritonavir* ↑ efavirenz*	Monitor for elevation of liver enzymes and for adverse clinical experiences (eg, dizziness, nausea, dyspepsia) when efavirenz tablets is coadministered with ritonavir.

Table 5: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies or Predicted Interaction

Concomitant Drug Class: Drug Name	Effect	Clinical Comment
Protease inhibitor: Saquinavir	↓ saquinavir*	Appropriate doses of the combination of efavirenz tablets and saquinavir/ritonavir with respect to safety and efficacy have not been established.
NNRTI: Other NNRTIs	↑ or ↓ efavirenz and/or NNRTI	Combining two NNRTIs has not been shown to be beneficial. Efavirenz tablets should not be coadministered with other NNRTIs.
CCR5 co-receptor antagonist: Maraviroc	↓ maraviroc*	Refer to the full prescribing information for maraviroc for guidance on coadministration with efavirenz.
Hepatitis C antiviral agents		
Boceprevir	↓ boceprevir*	Concomitant administration of boceprevir with efavirenz tablet is not recommended because it may result in loss of therapeutic effect of boceprevir.
Elbasvir/Grazoprevir	↓ elbasvir ↓ grazoprevir	Coadministration of clopidogrel with elbasvir/grazoprevir is not recommended *[see Contraindications (4)]* because it may lead to loss of virologic response to elbasvir/ grazoprevir.
Pibrentasvir/Glecaprevir	↓ pibrentasvir ↓ glecaprevir	Coadministration of clopidogrel is not recommended because it may lead to reduced therapeutic effect of pibrentasvir/glecaprevir.
Simeprevir	↓ simeprevir* ↔efavirenz*	Concomitant administration of simeprevir with efavirenz tablets is not recommended because it may result in loss of therapeutic effect of sime

8.4 Pediatric Use

The safety, pharmacokinetic profile, and virologic and immunologic responses of efavirenz tablets were evaluated in antiretroviral-naive and -experienced HIV-1 infected pediatric patients 3 months to 21 years of age in three open-label clinical trials [*see Adverse Reactions (6.2), Clinical Pharmacology (12.3), and Clinical Studies (14.2)*]. The type and frequency of adverse reactions in these trials were generally similar to those of adult patients with the exception of a higher frequency of rash, including a higher frequency of Grade 3 or 4 rash, in pediatric patients compared to adults [*see Warnings and Precautions (5.8) and Adverse Reactions (6.2)*].

Use of efavirenz tablets in patients younger than 3 months of age OR less than 3.5 kg body weight is not recommended because the safety, pharmacokinetics, and antiviral activity of efavirenz tablets have not been evaluated in this age group and there is a risk of developing HIV resistance if efavirenz tablets are underdosed. See *Dosage and Administration (2.2)* for dosing recommendations for pediatric patients.

8.5 Geriatric Use

Clinical studies of efavirenz tablets did not include sufficient numbers of subjects aged 65 years and over to determine whether they respond differently from younger subjects. In general, dose selection for an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other therapy.

8.6 Hepatic Impairment

Efavirenz tablets are not recommended for patients with moderate or severe hepatic impairment because there are insufficient data to determine whether dose adjustment is necessary. Patients with mild hepatic impairment may be treated with efavirenz without any adjustment in dose. Because of the extensive cytochrome P450-mediated metabolism of efavirenz and limited clinical experience in patients with hepatic impairment, caution should be exercised in administering efavirenz tablets to these patients [*see Warnings and Precautions (5.9) and Clinical Pharmacology (12.3)*].

10 OVERDOSAGE

Some patients accidentally taking 600 mg twice daily have reported increased nervous system symptoms. One patient experienced involuntary muscle contractions.

Treatment of overdose with efavirenz tablets should consist of general supportive measures, including monitoring of vital signs and observation of the patient's clinical status. Administration of activated charcoal may be used to aid removal of unabsorbed drug. There is no specific antidote for overdose with efavirenz tablets. Since efavirenz is highly protein bound, dialysis is unlikely to significantly remove the drug from blood.

11 DESCRIPTION

Efavirenz is an HIV-1 specific, non-nucleoside, reverse transcriptase inhibitor (NNRTI). Efavirenz USP is chemically described as (*S*)-6-chloro-4-(cyclopropylethynyl)-1,4-dihydro-4-(trifluoromethyl)-2*H*-3,1-benzoxazin-2-one. Its empirical formula is $C_{14}H_9ClF_3NO_2$ and its structural formula is:



Efavirenz USP is a white to slightly pink crystalline powder with a molecular mass of 315.68. It is practically insoluble in water (<10 microgram/mL).

Tablets: Efavirenz are available as film-coated tablets for oral administration containing 600 mg of efavirenz USP and the following inactive ingredients: microcrystalline cellulose, sodium lauryl sulfate, croscarmellose sodium, hydroxypropyl cellulose, lactose monohydrate, magnesium stearate. The film coating contains Opadry® Yellow (hypromellose, titanium dioxide, iron oxide yellow and polyethylene glycol)

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Efavirenz is an antiviral drug [*see Microbiology (12.4)*].

12.2 Pharmacodynamics

Cardiac Electrophysiology

The effect of efavirenz on the QTc interval was evaluated in an open-label, positive and placebo controlled, fixed single sequence 3-period, 3-treatment crossover QT study in 39 healthy subjects enrolled for CYP2B6 polymorphism. The mean C_{max} of efavirenz in subjects with CYP2B6 *6/*6 genotype following the administration of 600 mg daily dose for 14 days was 2.25-fold the mean C_{max} observed in subjects with CYP2B6 *1/*1 genotype. A positive relationship between efavirenz concentration and QTc prolongation was observed. Based on the concentration-QTc relationship, the mean QTc prolongation and its upper bound 90% confidence interval are 8.7 ms and 11.3 ms in subjects with CYP2B6*6/*6 genotype following the administration of 600 mg daily dose for 14 days *[see Warnings and Precautions (5.2)]*.

12.3 Pharmacokinetics

Absorption

Peak efavirenz plasma concentrations of 1.6 to 9.1 µM were attained by 5 hours following single oral doses of 100 mg to 1,600 mg administered to uninfected volunteers. Dose-related increases in C_{max} and AUC were seen for doses up to 1,600 mg; the increases were less than proportional suggesting diminished absorption at higher doses.

In HIV-1-infected patients at steady state, mean C_{max}, mean C_{min}, and mean AUC were dose proportional following 200 mg, 400 mg, and 600 mg daily doses. Time-to-peak plasma concentrations were approximately 3 to 5 hours and steady-state plasma concentrations were reached in 6 to 10 days. In 35 patients receiving efavirenz tablets 600 mg once daily, steady-state C_{max} was 12.9 ± 3.7 µM (mean ± SD), steady-state C_{min} was 5.6 ± 3.2 µM, and AUC was 184 ± 73 µM•h.

Effect of Food on Oral Absorption:

Tablets: Administration of a single 600 mg efavirenz tablet with a high-fat/high-calorie meal (approximately 1,000 kcal, 500 to 600 kcal from fat) was associated with a 28% increase in mean AUC, of efavirenz and a 79% increase in mean C_{max} of efavirenz relative to the exposures achieved under fasted conditions. [*See Dosage and Administration (2) and Patient Counseling Information (17)*.]

Distribution

Efavirenz is highly bound (approximately 99.5 to 99.75%) to human plasma proteins, predominantly albumin. In HIV-1 infected patients (n=9) who received efavirenz tablets 200 to 600 mg once daily for at least one month, cerebrospinal fluid concentrations ranged from 0.26 to 1.19% (mean 0.69%) of the corresponding plasma concentration. This proportion is approximately 3-fold higher than the non-protein-bound (free) fraction of efavirenz in plasma.

Metabolism

Studies in humans and *in vitro* studies using human liver microsomes have demonstrated that efavirenz is principally metabolized by the cytochrome P450 system to hydroxylated metabolites with subsequent glucuronidation of these hydroxylated metabolites. These metabolites are essentially inactive against HIV-1. The *in vitro* studies suggest that CYP3A and CYP2B6 are the major isozymes responsible for efavirenz metabolism.

Efavirenz has been shown to induce CYP enzymes, resulting in the induction of its own metabolism. Multiple doses of 200 to 400 mg per day for 10 days resulted in a lower than predicted extent of accumulation (22 to 42% lower) and a shorter terminal half-life of 40 to 55 hours (single dose half-life 52 to 76 hours).

Elimination

Efavirenz has a terminal half-life of 52 to 76 hours after single doses and 40 to 55 hours after multiple doses. A one-month mass balance/excretion study was conducted using 400 mg per day with a ^{14}C-labeled dose administered on Day 8. Approximately 14 to 34% of the radiolabel was recovered in the urine and 16 to 61% was recovered in the feces. Nearly all of the urinary excretion of the radiolabel was attributable to the form of metabolites. Efavirenz accounted for the majority of the total radioactivity measured in feces.

Special Populations

Pediatrics: The pharmacokinetic parameters for efavirenz at steady state in pediatric patients were predicted by a population pharmacokinetic model.

Gender and race: The pharmacokinetics of efavirenz in patients appear to be similar between men and women and among the racial groups studied.

Renal impairment: The pharmacokinetics of efavirenz have not been studied in patients with renal insufficiency; however, less than 1% of efavirenz is excreted unchanged in the urine, so the impact of renal impairment on efavirenz elimination should be minimal.

Hepatic impairment: A multiple-dose study showed no significant effect on efavirenz pharmacokinetics in patients with mild hepatic impairment (Child-Pugh Class A) compared with controls. There were insufficient data to determine whether moderate or severe hepatic impairment (Child-Pugh Class B or C) affects efavirenz pharmacokinetics.

Drug Interaction Studies

Efavirenz has been shown *in vivo* to cause hepatic enzyme induction, thus increasing the biotransformation of some drugs metabolized by CYP3A and CYP2B6. *In vitro* studies have shown that efavirenz inhibited CYP isozymes 2C9 and 2C19 with K_i values (8.5 to 17 µM) in the range of observed efavirenz plasma concentrations. In *in vitro* studies, efavirenz did not inhibit CYP2E1 and inhibited CYP3D6 and CYP1A2 (K_i values 82 to 160 µM) only at concentrations well above those achieved clinically. Coadministration of efavirenz with drugs primarily metabolized by CYP2C9, CYP2C19, CYP2C8 or CYP2B6 isozymes may result in altered plasma concentrations of the coadministered drug. Drugs which induce CYP3A and CYP2B6 activity would be expected to increase the clearance of efavirenz resulting in lowered plasma concentrations.

Drug interaction studies were performed with efavirenz and other drugs likely to be coadministered or drugs commonly used as probes for pharmacokinetic interaction. The effects of coadministration of efavirenz on the C_{max}, AUC, and C_{min} are summarized in Table 7 (effect of efavirenz on other drugs) and Table 8 (effect of other drugs on efavirenz). For information regarding clinical recommendations see *Drug Interactions (7.1)*.

Table 7: Effect of Efavirenz on Coadministered Drug Plasma C_{max}, AUC, and C_{min}

Coadministered Drug	Dose	Efavirenz Dose	Number of Subjects	Coadministered Drug (mean % change)		
				C_{max} (90% CI)	AUC (90% CI)	C_{min} (90% CI)
Atazanavir	400 mg qd with a light meal ‡ 1-20	600 mg qd with a light meal ‡ 21-28	27	↓ 59% (49-67%)	↓ 74% (68-78%)	↓ 85% (90-95%)
	400 mg qd ‡ 1-6, then 300 mg qd ‡ 7-20 with ritonavir 100 mg qd with a light meal ‡	600 mg qd ‡ 21 after atazanavir and ritonavir at 7-20	13	↑ 14%* (-17 - 1 5%)	↑ 99%* (2- 88%)	↑ 48%* (24-76%)
	300 mg qd/ritonavir 100 mg qd ‡ 1-10 with a light meal, then 400 mg qd/ritonavir 100 mg qd ‡ 11-24 (pm) (simultaneous with efavirenz)	600 mg qd with a light snack ‡ 11-24 (pm)	14	↑ 17% (8-27%)	**	↓ 42% (31-51%)
Indinavir	1,000 mg q8h x 10 days	600 mg qd x 10 days	20			
	After morning dose		**	↓ 33%* (26-39%)	↓ 39%* (24-51%)
	After afternoon dose		**	↓ 37% ‡ (26-46%)	↓ 52% ‡ (47-57%)
	After evening dose		**	↓ 29% ‡ (11-42%)	↓ 46% ‡ (37-54%)	↓ 57% ‡ (50-63%)
Lopinavir/ ritonavir	400/100 mg capsules q12h x 9 days	600 mg qd x 9 days	11_7*	**	↓ 19% ‡ (4 36-13%)	↓ 39% ‡ (3-62%)
	500/125 mg tablet q12h x 10 days with efavirenz compared to 400/100 mg q12h dose	600 mg qd x 9 days	19	↑ 12% ‡ (2-23%)	** ‡	↓ 10% ‡ (↓ 22- 1 4%)
	600/150 mg tablet q12h x 10 days with efavirenz compared to 400/100 mg q12 h dose	600 mg qd x 9 days	23	↑ 36% ‡ (28-44%)	↑ 56% ‡ (28-44%)	↑ 32% ‡ (21-44%)
Nelfinavir	750 mg q8h x 7 days	600 mg qd x 7 days	10	↑ 21% (10-33%)	↑ 20% (8-34%)	**
Metabolite AG-1402				↓ 40% (30-48%)	↓ 37% (25-48%)	↓ 43% (21-59%)
Ritonavir	500 mg q12h x 8 days	600 mg qd x 10 days	11			
	After AM dose			↑ 24% (12-38%)	↑ 18% (6-33%)	↑ 42% (9-86%)
	After PM dose			**	**	↑ 24% (3-50%) ¶

Table 7: Effect of Efavirenz on Coadministered Drug Plasma C_{max}, AUC, and C_{min}

Coadministered Drug	Dose	Efavirenz Dose	Number of Subjects	Coadministered Drug (mean % change)		
				C_{max} (90% CI)	AUC (90% CI)	C_{min} (90% CI)
Saquinavir SGC	1,200 mg q8h x 10 days	600 mg qd x 10 days	12	↓ 50% (39-68%)	↓ 62% (42-74%)	↓ 95% (97-77%)#
Lamivudine	150 mg q12h x 14 days	600 mg qd x 14 days	9	**	**	↑ 265% (37-672%)
Tenofovir§	300 mg qd	600 mg qd x 14 days	29	**	**	**
Zidovudine	300 mg q12h x 14 days	600 mg qd x 14 days	9	**	**	↑ 22% (40-64%)
Maraviroc	100 mg bid	600 mg qd	12	↓ 51% (37-62%)	↓ 45% (38-51%)	↓ 45% (28-57%)
Raltegravir	400 mg single dose	600 mg qd x 9 days	9	↓ 36% (2-59%)	↓ 36% (20-48%)	↓ 21% (↓ 51- 7 38%)
Bocoprevir	800 mg tid x 6 days	600 mg qd x 16 days	NA	↓ 8% (↓ 22- 7 8%)	↓ 19% (11-25%)	↓ 44% (26-58%)
Simeprevir	150 mg qd x 14 days	600 mg qd x 14 days	23	↓ 51% (↓ 46-↓28%)	↓ 71% (↓ 67-↓74%)	↓ 91% (↓ 89-↓92%)
Azithromycin	600 mg single dose	400 mg qd x 14 days	14	↑ 22% (4-42%)	**	NA
Clarithromycin	500 mg q12h x 7 days	400, mg qd x 7 days	11	↓ 26% (↓ 53-5%)	↓ 39% (↓ 49-↓26%)	↓ 53% (↓ 62-↓43%)
14-OH metabolite				↑ 49% (32-69%)	↑ 34% (↓ 18-53%)	↑ 26% (9-45%)
Fluconazole	200 mg x 7 days	400 mg qd x 7 days	10	**	**	**
Itraconazole	200 mg q12h x 28 days	200 mg qd x 14 days	18	↓ 37% (20-51%)	↓ 39% (21-53%)	↓ 44% (27-58%)
Hydroxy-itraconazole				↓ 35% (12-52%)	↓ 37% (14-55%)	↓ 43% (19-60%)
Posaconazole	400 mg (oral suspension) bid x 10 and 20 days	400 mg qd x 10 and 20 days	11	↓ 45% (34-55%)	↓ 50% (40-57%)	NA
Rifabutin	300 mg qd x 14 days	600 mg qd x 14 days	9	↓ 32% (15-46%)	↓ 38% (28-47%)	↓ 45% (31-58%)
Voriconazole	400 mg q12h x 1 day, then 200 mg q12h x 8 days	400 mg qd x 9 days	NA	↓ 61%‡	↓ 77%‡	NA
	300 mg po q12h days 2-7	300 mg qd x 7 days	NA	↓ 36%‡ (21-49%)	↓ 55%‡ (45-62%)	NA
	400 mg po q12h days 2-7	300 mg qd x 7 days	NA	↓ 23% (↓ 1 - 7 43%)	↓ 7% (↓ 23- 1 5%)	NA
Artemether/ lumefantrine	Artemether 20 mg/lumefantrine 120 mg tablets (6/4 tablet doses over 3 days)	600 mg qd x 26 days	12			
Artemether				↓ 21%	↓ 51%	NA
dihydroartemisinin				↓ 38%	↓ 46%	NA
lumefantrine				**	↓ 21%	NA
Atorvastatin	10 mg qd x 4 days	600 mg qd x 14 days	14	↓ 14% (↓ 26-3%)	↓ 43% (34-50%)	↓ 69% (49-81%)
Total active (including metabolites)				↓ 15% (2-26%)	↓ 32% (21-41%)	↓ 48% (23-64%)
Pravastatin	40 mg qd x 4 days	600 mg qd x 15 days	13	↓ 32% (↓ 49-↓12%)	↓ 44% (26-57%)	↓ 19% (0-35%)
Simvastatin	40 mg qd x 4 days	600 mg qd x 14 days	14	↓ 72% (63-79 %)	↓ 68% (62- 7 3 %)	↓ 45% (30-62%)
Total active (including metabolites)				↓ 68 % (55-78 %)	↓ 60 % (52-68 %)	NA‡
Carbamazepine	200 mg qd x 3 days, 200 mg bid x 3 days, then 400 mg qd x 29 days	600 mg qd x 14 days	12	↓ 20% (↓ 27-↓ 2%)	↓ 27% (20-33%)	↓ 35% (24-44%)
Epoxide metabolite				**	**	↓ 13% (↓ 30-↑7%)
Cetirizine	10 mg single dose	600 mg qd x 10 days	11	↓ 24% (↓ 6-↑ 2%)	**	NA
Diltiazem	240 mg x 21 days	600 mg qd x 14 days	13	↓ 69% (58-69%)	↓ 69% (63-74%)	↓ 63% (44-75%)
Desacetyl diltiazem				↓ 64% (57-69%)	↓ 75% (59-84%)	↓ 62% (44-75%)
N-monodes-methyl diltiazem				↓ 38% (7-44%)	↓ 37% (17-52%)	↓ 52% (17-52%)
Ethinyl estradiol/ Norgestimate	0.035 mg/0.25 mg x 14 days	600 mg qd x 14 days				
Ethinyl estradiol			21	**	**	**
Norelgestromin			21	↓ 46% (59-32%)	↓ 64% (69-57%)	↓ 82% (79-85%)
Levonorgestrel			6	↓ 80% (77-83%)	↓ 83% (79-87%)	↓ 86% (80-90%)
Lorazepam	2 mg single dose	600 mg qd x 10 days	12	↑ 16% (2-32%)	↓ 12%	NA
Methadone	Stable maintenance 35-100 mg daily	600 mg qd x 14-21 days	11	↓ 45% (25-59%)	↓ 52% (33-66%)	NA
Bupropion	150 mg single dose (sustained-release)	600 mg qd x 14 days	13	↓ 34% (21-47%)	↓ 55% (48-62%)	NA
Hydroxy-bupropion				↑ 50% (20-80%)	**	NA
Paroxetine	20 mg qd x 14 days	600 mg qd x 14 days	16	**	**	**
Sertraline	50 mg qd x 14 days	600 mg qd x 14 days	13	↓ 29% (15-40%)	↓ 39% (27-50%)	↓ 46% (31-58%)

‡Indicates increase ↓ Indicates decrease ** Indicates no change or a mean increase or decrease of <10%.

* Compared with atazanavir 400 mg qd alone.

† Comparator dose of indinavir 800 mg q8h x 10 days.

‡ Parallel-group design; *n* for efavirenz + lopinavir/ritonavir, *n* for lopinavir/ritonavir alone.

§ Values are for lopinavir, the pharmacokinetics of ritonavir in this study were unaffected by concurrent efavirenz.

¶ 95% CI.

Soft Gelatin Capsule.

Ω Tenofovir disoproxil fumarate.

Ω 90% CI not available.

‡ Relative to steady-state administration of voriconazole (400 mg for 1 day, then 200 mg po q12h for 2 days).

¶ Not available because of insufficient data.

NA = not available.

Table 8: Effect of Coadministered Drug on Efavirenz Plasma C_{max}, AUC, and C_{min}

Coadministered Drug	Dose	Efavirenz Dose	Number of Subjects	Efavirenz (mean % change)		
				C_{max} (90% CI)	AUC (90% CI)	C_{min} (90% CI)
Indinavir	800 mg q8h x 14 days	200 mg qd x 14 days	11	**	**	**
	800 mg/ritonavir 100 mg bid x 9 days	600 mg qd x 9 days	11_12*	**	↓ 16% (↓ 33-↑15%)	↓16% (↓42-↑20%)
Nelfinavir	750 mg q8h x 7 days	600 mg qd x 7 days	10	↑ 12% (↓ 2 2-↑ 3 3%)‡	↑ 13% (↓ 2 5-↑ 5 3%)‡	↑ 21% (↓ 4 3-↑ 5 3%)‡
Ritonavir	500 mg q12h x 8 days	600 mg qd x 10 days	9	↑ 14% (4-26%)	↑ 21% (10-34%)	↑ 25% (↑ 4-↑ 51%)
Saquinavir SGC	1,200 mg q8h x 10 days	600 mg qd x 10 days	13	↓ 33% (5-58%)	↓ 12% (4-19%)	↓ 14% (2-24%)‡
Tenofovir§	300 mg qd	600 mg qd x 14 days	30	**	**	**
Bocoprevir	800 mg tid x 6 days	600 mg qd x 16 days	NA	↑ 11% (2-20%)	↑ 20% (15-26%)	NA
simeprevir	150 mg qd x 14 days	600 mg qd x 14 days	23	**	↑ 10% (5-15%)	↑13% (↑ 1-19%)
Azithromycin	600 mg single dose	400 mg qd x 7 days	14	**	**	↑ 22% (5-41%)
Clarithromycin	500 mg q12h x 7 days	400 mg qd x 7 days	12	↑ 11% (3-19%)‡	**	**
Fluconazole	200 mg x 7 days	400 mg qd x 7 days	10	**	↑ 16% (6-26%)	↑ 22% (5-41%)
Itraconazole	200 mg q12h x 14 days	600 mg qd x 28 days	16	**	**	**
Rifabutin	300 mg qd x 14 days	600 mg qd x 14 days	11	**	**	↓ 12% (↓ 24-↑11%)
Rifampin	600 mg x 7 days	600 mg qd x 7 days	12	↓ 26% (11-39%)	↓ 26% (15-36%)	↓ 32% (15-46%)
Voriconazole	400 mg po q12h x 1 day, then 200 mg po q12h x 8 days	400 mg qd x 9 days	NA	↑ 38%‡	↑44%‡	NA
	300 mg po q12h days 2-7	300 mg qd x 7 days	NA	↑ 14%‡ (↑ 2-↑ 2 7%)	↑ 17%‡ (↑ 2-↑ 3 3%)	NA
	400 mg po q12h days 2-7	300 mg qd x 7 days	NA	↑ 38%‡ (↑ 1 7-↑ 6 3%)	↑ 44%‡ (↑ 2 3-↑ 7 0%)	NA
Artemether/ Lumefantrine	Artemether 20 mg/ lumefantrine 120 mg tablets (6/4 tablet doses over 3 days)	600 mg qd x 26 days	12	**	↑ 17%	NA
Atorvastatin	10 mg qd x 4 days	600 mg qd x 14 days	14	**	**	NA
Pravastatin	40 mg qd x 4 days	600 mg qd x 15 days	11	**	**	**

Table 8: Effect of Coadministered Drug on Efavirenz Plasma C_{max}, AUC, and C_{min}

Coadministered Drug	Dose	Efavirenz Dose	Number of Subjects	Efavirenz (mean % change)		
				C_{max} (90% CI)	AUC (90% CI)	C_{min} (90% CI)
Simvastatin	40 mg qd x 4 days	600 mg qd x 15 days	14	↓ 12% (↓ 28-↑ 8%)	**	↓ 12% (↓ 25-↑ 3%)
Aluminum hydroxide 400 mg, magnesium hydroxide 400 mg, plus simethicone 40 mg	30 mL single dose	400 mg single dose	17	**	**	NA
Carbamazepine	200 mg qd x 3 days, 200 mg bid x 3 days, then 400 mg qd x 15 days	600 mg qd x 35 days	14	↓ 21% (15-26%)	↓ 36% (32-40%)	↓ 47% (41-53%)
Cetirizine	10 mg single dose	600 mg qd x 10 days	11	**	**	**
Diltiazem	240 mg x 14 days	600 mg qd x 28 days	12	↑ 16% (6-26%)	↑ 11% (↓ 5-28%)	↑ 13% (↓ 7-34%)
Famotidine	40 mg single dose	400 mg single dose	17	**	**	NA
Paroxetine	20 mg qd x 14 days	600 mg qd x 14 days	12	**	**	**
Sertraline	50 mg qd x 14 days	600 mg qd x 14 days	13	↑ 11% (6-16%)	**	**

‡ Indicates increase ↓ Indicates decrease ** Indicates no change or a mean increase or decrease of <10%.

* Parallel-group design; *n* for efavirenz + lopinavir/ritonavir, *n* for efavirenz alone.

§ 95% CI.

† Soft Gelatin Capsule.

‡ Tenofovir disoproxil fumarate.

¶ 90% CI not available.

Relative to steady-state administration of efavirenz (600 mg once daily for 9 days).

NA = not available.

12.4 Microbiology

Mechanism of Action

Efavirenz is an NNRTI of HIV-1. Efavirenz activity is mediated predominantly by noncompetitive inhibition of HIV-1 reverse transcriptase. HIV-2 reverse transcriptase and human cellular DNA polymerases α, β, γ, and δ are not inhibited by efavirenz.

Antiviral Activity in Cell Culture

The concentration of efavirenz inhibiting replication of wild-type laboratory adapted strains and clinical isolates in cell culture by 90 to 95% (EC$_{90}$ to EC$_{95}$) ranged from 1.7 to 25 nM in lymphoblastoid cell lines, peripheral blood mononuclear cells (PBMCs), and macrophage/ monocyte cultures. Efavirenz demonstrated antiviral activity against clade B and most non-clade B isolates (subtypes A, AE, AG, C, D, F, G, J, N), but had reduced antiviral activity against group O viruses. Efavirenz demonstrated additive antiviral activity without cytotoxicity against HIV-1 in cell culture when combined with the nucleoside reverse transcriptase inhibitors (NRTIs) (abacavir, didanosine, lamivudine, stavudine, tenofovir disoproxil fumarate, zalcitabine, zidovudine), PIs (amprenavir, indinavir, nelfinavir, ritonavir, saquinavir), and the fusion inhibitor enfuvirtide. Efavirenz demonstrated additive to antagonistic antiviral activity in cell culture with atazanavir. Efavirenz was not antagonistic with adefovir, used for the treatment of chronic hepatitis B virus infection, or ribavirin, used in combination with interferon for the treatment of hepatitis C virus infection.

Resistance

In cell culture

In cell culture, HIV-1 isolates with reduced susceptibility to efavirenz (≥380-fold increase in EC$_{90}$ value) emerged rapidly in the presence of drug. Genotypic characterization of three viruses identified single amino acid substitutions L100I or V179D, double substitutions L100I/V108I, and triple substitutions L100I/V179D/Y181C in reverse transcriptase.

Clinical studies

Clinical isolates with reduced susceptibility in cell culture to efavirenz have been obtained. One or more substitutions at amino acid positions 98, 100, 101, 103, 106, 108, 138, 179, 181, 188, 190, 225, and 227 in reverse transcriptase were observed in patients failing treatment with efavirenz in combination with indinavir, or with zidovudine plus lamivudine. The K103N substitution was the most frequently observed. Long-term resistance surveillance (average 52 weeks, range 4 to 106 weeks) analyzed 28 matching baseline and virologic failure isolates. Sixty-one percent (17/28) of these failure isolates had decreased efavirenz susceptibility in cell culture with a median 88-fold change in efavirenz susceptibility (EC$_{90}$ value) from reference. The most frequent NNRTI substitution to develop in these patient isolates was K103N (64%). Other NNRTI substitutions that developed included L100I (7%), K101E/Q/R (14%), V106I (11%), G190S/T/A (7%), P225H (18%), and M230L (11%).

Cross-Resistance

Cross-resistance among NNRTIs has been observed. Clinical isolates previously characterized as efavirenz-resistant were also phenotypically resistant in cell culture to delavirdine and nevirapine compared to baseline. Delavirdine- and/or nevirapine-resistant clinical viral isolates with NNRTI resistance-associated substitutions (A98G, L100I, K101E/P, K103N/S, V106A/M, Y181C, Y188L/C/H, G190A, P225H, or M230L) showed reduced susceptibility to efavirenz in cell culture. Greater than 90% of NNRTI-resistant clinical isolates tested in cell culture retained susceptibility to efavirenz.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term carcinogenicity studies in mice and rats were carried out with efavirenz. Mice were dosed with 0, 25, 75, 150, or 300 mg/kg/day for 2 years. Incidences of hepatocellular adenomas and carcinomas and pulmonary alveolar/bronchiolar adenomas were increased above background in females. No increases in tumor incidence above background were seen in males. There was no NOAEL in mice established for this study because tumor findings occurred at all doses. AUC at the NOAEL (150 mg/kg) in the males was approximately 0.9 times that in humans at the recommended clinical dose. In the female mice, no increase in tumor incidence were observed at doses up to 100 mg/kg/day, for which AUCs were 0.1 (males) or 0.2 (females) times those in humans at the recommended clinical dose.

Mutagenesis

Efavirenz tested negative in a battery of *in vitro* and *in vivo* genotoxicity assays. These included bacterial mutation assays in *S. typhimurium* and *E. coli*, mammalian mutation assays in Chinese hamster ovary cells, chromosome aberration assays in human peripheral blood lymphocytes or Chinese hamster ovary cells, and an *in vivo* mouse bone marrow micronucleus assay.

Impairment of Fertility

Efavirenz did not impair mating or fertility of male or female rats, and did not affect sperm of treated male rats. The reproductive performance of offspring born to female rats given efavirenz was not affected. The AUCs at the NOAEL values in male (200 mg/kg) and female (100 mg/kg) rats were approximately ≤0.15 times that in humans at the recommended clinical dose.

13.2 Animal Toxicology

Microscopic observations were observed in 6 of 20 monkeys receiving efavirenz at doses yielding plasma AUC values 4- to 10-fold greater than those in humans given the recommended dose [*see Warnings and Precautions (5.10)*].

14 CLINICAL STUDIES

14.1 Adults

Study 006, a randomized, open-label trial, compared efavirenz tablets (600 mg once daily) + zidovudine (ZDV, 300 mg q12h) + lamivudine (LAM, 150 mg q12h) or efavirenz tablets (600 mg once daily) + indinavir (IDV, 1,000 mg q8h) without ritonavir (800 mg q8h) + zidovudine (300 mg q12h) + lamivudine (150 mg q12h). Twelve hundred sixty-six patients (mean age 36.5 years (range 18 to 81), 60% Caucasian, 83% male) were enrolled. All patients were efavirenz-, lamivudine-, NNRTI-, and PI-naive at study entry. The median time-to-AIDS or death was 2,500 cells/mm³ and the median baseline HIV-1 RNA level was 4.8 log$_{10}$ copies/mL. Treatment outcomes with standard assay (lower limit of detection 400 copies/mL) and ultrasensitive (assay limit 50 copies/mL) versions of the HIV-1 RNA assay through 48 weeks, and 3-drug regimen CD4 cell count changes through 48 weeks are summarized in Table 9.

Table 9: Outcomes of Randomized Treatment Through 48 and 168 Weeks, Study 006

Outcome	Efavirenz tablets + ZDV + LAM (n=422)		Efavirenz tablets + IDV (n=428)		IDV + ZDV + LAM (n=413)	
	Week 48	Week 168	Week 48	Week 168	Week 48	Week 168
Responder[a]	69%	48%	57%	40%	50%	29%
Virologic failure[b]	6%	12%	15%	26%	13%	19%
Discontinued for adverse events	7%	8%	6%	8%	16%	20%
Discontinued for other reasons[c]	17%	31%	22%	32%	21%	32%
CD4+ cell count cells/mm³)						
Observed subjects (n)	(279)	(205)	(256)	(158)	(228)	(129)
Mean change from baseline	190	329	191	310	180	329

[a] Patients achieved and maintained confirmed HIV-1 RNA <400 copies/mL through Week 48 or Week 168.

[b] Includes patients who rebounded, patients who were on study at Week 48 and failed to achieve confirmed HIV-1 RNA <400 copies/mL at time of discontinuation, and patients who discontinued due to lack of efficacy.

[c] Includes consent withdrawn, lost to follow-up, noncompliance, never treated, missing data, protocol violation, death, and other reasons. Patients with HIV-1 RNA levels <400 copies/mL who chose not to continue in the voluntary extension phases of the study were censored at date of last dose of study medication.

For patients treated with efavirenz tablets + zidovudine + lamivudine, efavirenz tablets + indinavir, or indinavir + zidovudine + lamivudine, the percentage of responders with HIV-1 RNA <50 copies/mL was 65%, 50%, and 45%, respectively, through 48 weeks, and 43%, 31%, and 23%, respectively, through 168 weeks. A Kaplan-Meier analysis of time to loss of virologic response (HIV RNA <400 copies/mL) suggests that both the trends of virologic response and differences in response continue through 4 years.

ACTG 364 is a randomized, double-blind, placebo-controlled, 48-week study in NRTI-experienced patients who had completed two prior ACTG studies. One-hundred-ninety-six patients (mean age 40 years (range 18 to 76), 74% Caucasian, 88% male) received NRTIs in combination with efavirenz tablets (600 mg once daily), or nelfinavir (750 mg three times daily), or efavirenz tablets (600 mg once daily) + nelfinavir in a randomized, double-blinded manner. The mean baseline CD4+ cell count was 389 cells/mm³ and mean baseline HIV-1 RNA level was 4.139 copies/mL. Upon entry into the study, all patients were assigned a genotypic sensitivity NRTI regimen that was augmented as their previous NRTI treatment experience dictated. There was no significant difference in the CD4+ cell count among the 3 treatment groups. The antiviral response was comparable in approximately 100 cells at 48 weeks among patients who continued on study regimens. Treatment outcomes are shown in Table 10. Plasma HIV RNA levels were quantified with the AMPLICOR HIV-1 MONITOR assay using a lower limit of quantification of 500 copies/mL.

Table 10: Outcomes of Randomized Treatment Through 48 Weeks, Study ACTG 364*

Outcome	Efavirenz tablets + NRTI + NRTIs (n=65)	Efavirenz tablets + NRTIs (n=65)	NFV +NRTIs (n=66)
HIV-1 RNA <500 copies/mL†	71%	63%	41%
HIV-1 RNA ≥500 copies/mL†	17%	34%	54%
CDC category C Event	2%	0%	5%
Discontinuations for adverse events‡	3%	3%	5%
Discontinuations for other reasons§	8%	0%	0%

* For some patients, Week 56 data were used to confirm the status at Week 48.

† Subjects achieved virologic response (two consecutive viral loads <500 copies/mL) and maintained it through Week 48.

‡ Includes viral rebound and failure to achieve confirmed <500 copies/mL by Week 48.

§ See *Adverse Reactions (6.1)* for a safety profile of these regimens.

¶ Includes loss to follow-up, consent withdrawn, noncompliance.

A Kaplan-Meier analysis of time to treatment failure through 72 weeks demonstrates a longer duration of virologic suppression (HIV RNA <500 copies/mL) in the efavirenz tablets-containing treatment arms.

14.2 Pediatric Patients

Study AI266022 is an open-label study to evaluate the pharmacokinetics, safety, tolerability, and antiviral activity of efavirenz in combination with didanosine and emtricitabine in antiretroviral-naive and -experienced pediatric patients. Thirty-seven patients 3 months to 6 years of age (median 0.7 years) were treated with efavirenz. At baseline, median plasma HIV-1 RNA was 5.88 log$_{10}$ copies/mL, median CD4+ cell count was 1,144 cells/mm³, and median CD4+ percentage was 25%. The median time on study therapy was 60 weeks; 27% of patients discontinued before Week 48. Using an ITT analysis, the overall proportions of patients with HIV RNA <400 copies/mL and <50 copies/mL at Week 48 were 57% (21/37) and 46% (17/37), respectively. The median increase from baseline in CD4+ count at 48 weeks was 196 cells/mm³ and the median increase in CD4+ percentage was 6%.

Study PACTG 1,021 was an open