HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all the information needed to use EFAVIRENZ, LAMIVUDINE AND TENOFOVIR DISOPROXIL FUMARATE TABLETS safely and effectively. See full prescribing information for EFAVIRENZ, LAMIVUDINE AND TENOFOVIR DISOPROXIL FUMARATE TABLETS.

EFAVIRENZ, LAMIVUDINE and TENOFOVIR DISOPROXIL FUMARATE tablets, for oral use — Initial U.S. Approval: 2006

WARNING: LACTIC ACIDOSIS/SEVERE HEPATOMEGALY WITH STEATOSIS and POST TREATMENT ACUTE EXACERBATIONS OF HEPATITIS B
See full prescribing information for complete boxed warning.
- **Lactic acidosis and hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogues including lamivudine and tenofovir disoproxil fumarate. (5.1)**
- **Severe acute exacerbations of hepatitis B have been reported in patients who are co-infected with HIV-1 and hepatitis B virus and have discontinued lamivudine, which is a component of these products. Monitor hepatic function closely in these patients and, if appropriate, initiate anti-hepatitis B treatment. (5.2)**

INDICATIONS AND USAGE
Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, a combination of one non-nucleoside reverse transcriptase inhibitor (efavirenz) and two nucleoside reverse transcriptase inhibitors (lamivudine and tenofovir disoproxil fumarate), are indicated for use alone as a complete regimen for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in adults and pediatric patients weighing at least 40 kg. (1)

DOSAGE AND ADMINISTRATION
- Recommended dose: One tablet (containing 600 mg of efavirenz, 300 mg of lamivudine and 300 mg of tenofovir disoproxil fumarate) taken once daily orally on an empty stomach, preferably at bedtime. (2.1)

DOSAGE FORMS AND STRENGTHS
Tablets: 600 mg of efavirenz, 300 mg lamivudine and 300 mg tenofovir disoproxil fumarate. (3)

CONTRAINDICATIONS
- Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are contraindicated in patients with previous hypersensitivity (e.g., Stevens-Johnson syndrome, erythema multiforme, or toxic skin epidermal necrolysis) to any of the components of this product. (4.1)

WARNINGS AND PRECAUTIONS
- Coadministration with Other Products: Do not use with other efavirenz, lamivudine or tenofovir-containing products or with lamivudine-containing products. Do not administer in combination with adefovir dipivoxil. (5.3)
- Hepatic decompensation, some fatal, has occurred in HIV-1/HCV co-infected patients receiving combination antiretroviral therapy. Monitor patients with pre-existing liver dysfunction, including chronic active hepatitis, for exacerbations of hepatic symptoms and laboratory abnormalities. Interrupt or discontinue therapy as clinically appropriate and consider dose reduction or discontinuation of interferon alfa, ribavirin, or both. (5.4)
- Nervous System Symptoms: Dizziness, poor concentration, and abnormal dreams are common efavirenz side effects. Discontinue efavirenz if these symptoms occur and evaluate patients to identify whether other potential risk factors for psychiatric disorders are present, and if so, consider whether the risk of continued therapy is clinically appropriate. (5.6)
- New-Onset or Worsening Renal Impairment: Can include acute renal failure and Fanconi syndrome. Assess estimated creatinine clearance before initiating treatment with tenofovir disoproxil fumarate. A component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, tenofovir disoproxil fumarate can cause renal toxicity. Avoid concomitant use of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets with nephrotoxic agents, including the chronic use of high-dose or multiple NSAIDs. (5.7)
- Serious Psychiatric Symptoms: Immediate medical evaluation is recommended for serious psychiatric symptoms such as severe depression or suicidal ideation. (5.7, 17)
- Nervous System Symptoms (NSS): NSS are frequent, usually begin 1 to 2 days after initiating therapy and resolve in 2 to 4 weeks. Dosing of sedating medications may improve tolerability. NSS are not predictive of more serious psychiatric symptoms. (5.6, 17)
- Embryo-Fetal Toxicity: Avoid administration in the first trimester of pregnancy as fetal harm may occur. (5.8, 8.1)
- Neurocognitive: Monitor liver function tests before and during treatment in patients with underlying hepatic disease, including those co-infected with hepatitis B and/or hepatitis C virus, or who are taking medications associated with liver toxicity. Among reported cases of hepatic failure, a few occurred in patients with no pre-existing hepatic disease. (5.11, 6.1)
- Rash: Rash usually begins within 1 to 2 weeks after initiating therapy and resolves within 4 weeks. Discontinue if severe rash develops. (5.16, 17)
- Convulsions: Use caution in patients with a history of seizures. (5.12)
- Lipids: Total cholesterol and triglyceride elevations. Monitor before therapy and periodically thereafter. (5.13)
- Decreases in Bone Mineral Density (BMD): Observed in HIV-infected patients. Consider assessment of BMD in patients with a history of pathologic fracture or other risk factors for osteoporosis or bone loss. (5.14)
- Immune Reconstitution Syndrome: Observed in HIV-infected patients. May necessitate further evaluation and treatment. (5.15)
- Redistribution/Accumulation of Body Fat: Observed in HIV-infected patients receiving antiretroviral therapy. (5.16)

ADVERSE REACTIONS
- Most common adverse reactions are impaired concentration, abnormal dreams, headache, nausea, malaise and fatigue, visual signs and symptoms, diarrhea, rash, dizziness, insomnia, pain, depression, asthenia, and cough. (6)

DRUG INTERACTIONS
Coadministration of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets can alter the concentrations of other drugs and other drugs may alter the concentrations of efavirenz, lamivudine and tenofovir disoproxil fumarate. The possibility of clinically significant drug interactions should be considered before and during therapy. (7)

USE IN SPECIFIC POPULATIONS
- Pregnancy: Women should avoid becoming pregnant while receiving efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, and for 12 weeks after discontinuation. (8.6)
- Lactation: Breastfeeding not recommended. (8.3)
- Females and Males of Reproductive Potential: Pregnancy testing and contraception are recommended. (8.3)
- Hepatic Impairment: Severe or pre-existing hepatic disease in patients with moderate or severe hepatic impairment. Use caution in patients with mild hepatic impairment. (8.7)
- Pediatrics: Safety in patients below 40 kg body weight has not been established. (8.4)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 09/2007

Efavirenz, Lamivudine and Tenofovir Disoproxil Fumarate Tablets 200405/04

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: LACTIC ACIDOSIS/SEVERE HEPATOMEGALY WITH STEATOSIS and POST TREATMENT ACUTE EXACERBATIONS OF HEPATITIS B

1 INDICATIONS AND USAGE
2 DOSAGE AND ADMINISTRATION
2.1 Recommended Dosage for Adult Patients and Pediatric Patients (40 kg or More)
2.2 Dose Adjustment for Renal Impairment
3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
4.1 Hypersensitivity
5 WARNINGS AND PRECAUTIONS
5.1 Lactic Acidosis/Severe Hepatomegaly With Steatosis
5.2 Patients Coinfected with HIV-1 and HBV
5.3 Coadministration with Other Products
5.4 Use with Interferon- and Ribavirin-Regimens
5.5 Pancreatitis
5.6 New Onset or Worsening Renal Impairment
5.7 Psychiatric Symptoms
5.8 Nervous System Symptoms
5.9 Embryo-Fetal Toxicity
5.10 Rash
5.11 Hepatotoxicity
5.12 Convulsions
5.13 Lipid Elevations
5.14 Bone Effects
5.15 Immune Reconstitution Syndrome
5.16 Redistribution of Body Fat
5.17 Effects on Ability to Drive and Use Machines
6 ADVERSE REACTIONS
6.1 Clinical Trials Experience
6.2 Postmarketing Experience

FULL PRESCRIBING INFORMATION

WARNING: LACTIC ACIDOSIS/SEVERE HEPATOMEGALY WITH STEATOSIS and POST TREATMENT ACUTE EXACERBATIONS OF HEPATITIS B

Lactic acidosis and hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogues including lamivudine and tenofovir disoproxil fumarate. Suspend treatment if clinical or laboratory findings suggestive of lactic acidosis or pronounced hepatotoxicity occur (see Warnings and Precautions (5.1)).

Severe acute exacerbations of hepatitis B have been reported in patients who are co-infected with hepatitis B virus (HBV) and HIV-1 and have discontinued lamivudine, which is a component of these products. Monitor hepatic function closely in these patients and, if appropriate, initiate anti-hepatitis B treatment (see Warnings and Precautions (5.2)).

1 INDICATIONS AND USAGE

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are indicated for use alone as a complete regimen for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in adults and in pediatric patients weighing at least 40 kg.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage for Adult and Pediatric Patients weighing at least 40 kg

The recommended dosage of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets is 600 mg/300 mg/300 mg orally, once daily. It is recommended that efavirenz, lamivudine and tenofovir disoproxil fumarate tablets be taken on an empty stomach, preferably at bedtime. Administration with food may increase the exposure of efavirenz (see Clinical Pharmacology (12.3) in the full prescribing information), which may increase the frequency of adverse reactions (see Adverse Reactions (6.1) and Patient Counseling Information (17)). Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets should be swallowed intact with liquid.

2.2 Dose Adjustment for Renal Impairment

Because efavirenz, lamivudine and tenofovir disoproxil fumarate is a fixed-dose combination tablet, it is not recommended for use in patients with creatinine clearance less than 50 mL/min, or in patients who end-stage renal disease (ESRD) requiring hemodialysis.

3 DOSAGE FORMS AND STRENGTHS

Efavirenz, Lamivudine and Tenofovir Disoproxil Fumarate Tablets are available containing 600 mg of efavirenz, 300 mg of lamivudine, USP and 300 mg tenofovir disoproxil fumarate equivalent to 300 mg of tenofovir disoproxil.

- The 600 mg/300 mg/300 mg tablets are yellow colored, capsule-shaped, beveled edged biconvex film coated tablets debossed with "1" on one side and "227" on the other side.

4 CONTRAINDICATIONS

4.1 Hypersensitivity

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are contraindicated in patients with a previous hypersensitivity reaction to any of the components contained in the formulations.

5 WARNINGS AND PRECAUTIONS

5.1 Lactic Acidosis/Severe Hepatomegaly With Steatosis

Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogues including lamivudine and tenofovir disoproxil fumarate. A class effect of these drugs is prolonged nucleoside exposure may be risk factor. Caution should be exercised when administering any nucleoside analogs to any patient with known risk factors for liver disease; however, cases have also been reported in patients with no known risk factors. Treatment should be suspended in any patient who develops clinical or laboratory findings suggestive of lactic acidosis or pronounced hepatotoxicity (which may include hepatomegaly and steatosis even in the absence of marked transaminase elevations).

5.2 Patients Coinfected with HIV-1 and HBV

It is not known whether efavirenz is active for the treatment of chronic hepatitis B virus (HBV) before initiating antiretroviral therapy. Discontinuation of anti-HBV therapy, including lamivudine and tenofovir disoproxil fumarate, may be associated with severe acute exacerbations of hepatitis. Patients infected with HBV who discontinue efavirenz, lamivudine and tenofovir disoproxil fumarate tablets should be closely monitored with both clinical and laboratory follow-up for at least several months after stopping treatment. If appropriate, resumption of anti-hepatitis B therapy may be warranted.

Important Differences Among Lamivudine-Containing Products: Lamivudine tablets contain a higher dose of the same active ingredient (lamivudine) than EPIVIR-HBV tablets. EPIVIR-HBV was developed for patients with chronic hepatitis B. The formulation and dosage of lamivudine in EPIVIR-HBV are not appropriate for co-infected patients with HIV-1 and HBV. Safety and efficacy of lamivudine have not been established for treatment of chronic hepatitis B in patients who are co-infected with HIV-1.

If treatment with EPIVIR-HBV or tenofovir disoproxil fumarate-containing product such as VIREAD is indicated for chronic hepatitis B for a patient with unrecognized or untreated HIV-1 infection, rapid emergence of HIV-1 resistance is likely to result because of the subtherapeutic dose and the inappropriateness of monotherapy in HIV-1 treatment.

5.3 Coadministration with Other Products

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are a fixed-dose combination product of efavirenz, lamivudine and tenofovir disoproxil fumarate. Do not administer with another efavirenz-containing product, lamivudine-containing product, lamivudine-containing, or emtricitabine-containing drugs. Avoid administration with adefovir dipivoxil. Didanosine (dideoxyinosine), EPIVIR® or EPIVIR-HBV® (lamivudine), EMTRIVA® (emtricitabine), TRUVADA® (abacavir sulfate/lamivudine/zidovudine), ABETREXA® (emtricitabine), EMTRIVA® (emtricitabine/tenofovir disoproxil fumarate), VIREAD® (tenofovir disoproxil fumarate), ATRIPLA® (emtricitabine/efavirenz/tenofovir disoproxil fumarate), COMBIVIR® (lamivudine/zidovudine), or EPIVIR-HBV® (lamivudine) are not recommended for use with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets.

5.4 Use With Interferon- and Ribavirin-Based Regimens

In vitro studies have shown ribavirin can reduce the phosphorylation of pyrimidine nucleoside analogues such as lamivudine, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Although no evidence of an interaction between ribavirin and lamivudine has been observed in HIV-1 infected patients, caution should be exercised when ribavirin was coadministered with lamivudine or with HIV-1/HCV co-infected patients (see *Clinical Pharmacology (12.3)*). Hepatic decompensation (some fatal) has occurred in HIV-1/HCV co-infected patients receiving combination antiretroviral therapy. Monitor patients with pre-existing liver dysfunction, including chronic active hepatitis, or without ribavirin and lamivudine should be closely monitored for treatment-associated function, especially hepatic decompensation. Discontinuation of lamivudine should be considered as medically appropriate. Dose reduction or discontinuation of interferon alfa, ribavirin, or both should also be considered if worsening clinical toxicities are observed, including hepatic decompensation (e.g., Child-Pugh-c). See the full prescribing information for ribavirin.

5.5 Pancreatitis

In pediatric patients with a history of prior antiretroviral nucleoside exposure, a history of pancreatitis, or other significant risk factors for the development of pancreatitis, lamivudine should be used with caution. Treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets should be stopped immediately if clinical signs, symptoms, or laboratory abnormalities suggestive of pancreatitis occur.

5.6 New Onset or Worsening Renal Impairment

Tenofovir, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets is principally eliminated by the kidney. Renal impairment, including cases of acute renal failure and Fanconi syndrome (renal tubular injury with severe hypophosphatemia), has been reported with the use of tenofovir disoproxil fumarate (see *Clinical Pharmacology (12.3)*).

It is recommended that estimated creatinine clearance be assessed in all patients prior to initiating therapy and as clinically appropriate during therapy with tenofovir disoproxil fumarate. In patients at risk of renal dysfunction, including patients who have previously experienced renal events while receiving ADEFOVIR dipivoxil, alternative therapies should be considered. It is recommended that the dosing interval be modified for patients with estimated creatinine clearance between 30 and 49 mL/min and phosphate in patients with estimated creatinine clearance between 10 and 49 mL/min. Renal function should be monitored closely during tenofovir disoproxil fumarate therapy.

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets should be avoided with concurrent or recent use of a nephrotoxic agent (e.g., high-dose or multiple non-steroidal anti-inflammatory drugs (NSAIDs)) (see *Drug Interactions (7.6)*). Cases of acute renal failure have been reported following the initiation of high-dose or multiple NSAIDs in HIV-infected patients with risk factors for renal dysfunction who appeared stable on tenofovir disoproxil fumarate. Some patients required hospitalization and renal replacement therapy. Alternatives to NSAIDs should be considered, if needed, in patients at risk for renal dysfunction.

Persistent or worsening bone pain, pain in extremities, fractures and/or muscular pain or weakness may be manifestations of proximal renal tubulopathy and should prompt an evaluation of renal function in at-risk patients.

5.7 Psychiatric Symptoms

Serious psychiatric adverse experiences have been reported in pregnant women. Advise a prospective or pregnant patient to discuss the risks and benefits of efavirenz therapy with her doctor. In controlled trials of 1008 patients who had not previously received efavirenz, 19% (194/1008) of patients treated with control regimens for a mean of 1.1 years, the frequency (regardless of causality) of specific serious psychiatric events among patients exposed efavirenz or control regimens, respectively, were severe depression (2.4%, 0.9%), suicidal ideation (0.7%, 0.4%), nonfatal suicide attempts (0.5%, 0.0%), aggressive behavior (1.2%, 0.4%), paranoia (1.2%, 0.4%), and manic reactions (0.3%, 0.2%). When psychiatric symptoms similar to those noted above were combined and evaluated as a group in a multifactorial analysis of data from Study 006, treatment with efavirenz was associated with a 2.4-fold increase (95% CI: 1.3, 4.4) in the occurrence of these events. The contribution of other risk factors to the occurrence of these psychiatric symptoms were history of injection drug use, psychiatric history, and receipt of psychiatric medication at study entry; similar associations were observed in both the efavirenz and control treatment groups. In 96 of 969, short of nine serious psychiatric symptoms occurred throughout the study for both efavirenz-treated and control-treated patients. One percent of efavirenz-treated patients discontinued therapy.

Interrupted treatment because of one or more of these selected psychiatric symptoms.

There have also been occasional postmarketing reports of death by suicide, delusions, psychosis-like behavior, and catatonia, although a causal relationship to the use of efavirenz cannot be determined from these reports. Patients with serious psychiatric adverse experiences should seek immediate medical evaluation to assess the possibility of suicide, the likelihood that symptoms are a consequence of the drug, and the potential risks and benefits of continued therapy schedule the baseline.

5.8 Nervous System Symptoms

Fifty-three percent (531/1008) of patients receiving efavirenz, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, in controlled trials reported central nervous system symptoms (any grade, regardless of causality) compared to 35% (180/509) of patients receiving control regimens. These symptoms usually began within the first or second day of therapy and generally resolve after the first 2 to 4 weeks of therapy. After 4 weeks of therapy, 2% (19/1008) of patients receiving efavirenz and 1% (5/509) of patients receiving control drugs had these symptoms. Symptoms of any duration led to discontinuation of efavirenz in 2% (19/1008) of patients treated with efavirenz and from 2% to 5% in patients treated with control regimens. Patients should be informed that these common symptoms were likely to improve with continued therapy and were not predictive of subsequent onset of the less frequent psychiatric symptoms (see *Warnings and Precautions (5.7)*). Dosing of bedtime may improve the tolerability of these nervous system symptoms (see *Dosage and Administration (2.1)*).

Analysis of long-term data from Study 006 (median follow-up 180 weeks, 100 weeks, and 76 weeks for patients treated with efavirenz + lamivudine + stavudine, efavirenz + indinavir, and indinavir + zidovudine + lamivudine, respectively) showed that, beyond the 4 weeks of therapy, the incidence of new nervous system symptoms among efavirenz-treated patients were generally similar to those in the indinavir-containing control arm.

Efavirenz can cause dizziness in patients who receive it. Advise patients to avoid potentially hazardous tasks such as driving or operating machinery until they are familiar with the potential effects of efavirenz on mental alertness and coordination.

Patients who experience central nervous system symptoms such as dizziness, impaired concentration, and/or drowsiness should avoid potentially hazardous tasks such as driving or operating machinery.

5.9 Embryo-Fetal Toxicity

Efavirenz, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, may cause fetal harm when administered during the first trimester to a pregnant woman. Advise females of reproductive potential who are receiving efavirenz to avoid pregnancy. (*See Use in Specific Populations (8.1)*.)

5.10 Rash

In controlled clinical trials, 26% (266/1008) of patients treated with 600 mg efavirenz experienced new-onset skin rash compared with 17% (117/682) of patients treated in control groups. Rash associated with blistering, moist desquamation, or ulceration occurred in 0.9% (9/1008) of patients treated with efavirenz. The incidence of rash (e.g., erythema multiforme, Stevens-Johnson syndrome) was highest during the first 1 to 2 weeks of therapy and expanded scores was 9.1%. Rashes are usually mild-to-moderate maculopapular skin eruptions that occur within the first 2 weeks of initiation of therapy with efavirenz and resolve spontaneously within 4 weeks. In clinical trials, in most patients continuing therapy with efavirenz, rash resolves within 1 month (median duration, 16 days). The discontinuation rate for rash in clinical trials was 1.7% (17/1008).

Efavirenz can generally be reinitiated in patients interrupting therapy because of rash. Efavirenz should be discontinued in patients developing severe rash associated with blistering, desquamation, mucosal involvement, or fever. Appropriate antihistamines and/or corticosteroids may improve the tolerability and hasten the resolution of rash. For patients who have had a life-threatening cutaneous reaction (e.g., Stevens-Johnson syndrome), alternative therapy should be considered. (see *Contraindications (4.1)*)

Rash was reported in 5% of 3 (3/57) pediatric patients (40%) treated with efavirenz capsules, patient experienced Grade 3 rash (confluent rash with fever), and two patients had Grade 4 rash (erythema multiforme). The median time to onset of rash in pediatric patients was 8 days. Prophylaxis with appropriate antihistamines before initiating therapy with efavirenz in pediatric patients should be considered.

5.11 Hepatotoxicity

Monitoring of liver enzymes before and during treatment is recommended for patients with underlying hepatic disease, including those co-infected with hepatitis B and/or hepatitis C virus, or who are taking medications associated with liver toxicity (see *Use in Specific Populations (8.7)*). A few of the postmarketing reports of hepatic failure occurred in patients with no pre-existing hepatic disease or other identifiable risk factors. Consideration should be given to interrupting therapy in patients with persistent elevations of serum transaminases to greater than five times the upper limit of the normal range, the benefit of continued therapy with efavirenz needs to be carefully reassessed in patients with significant liver toxicity.

5.12 Convulsions

Convulsions have been observed in patients receiving efavirenz, generally in the presence of known medical history of seizures (see *Nonclinical Toxicology (13.2)*). Caution should be taken in any patient with a history of seizures. These patients should be monitored during coadministration with medications that may affect the clearance of efavirenz and phenobarbital, may require periodic monitoring of plasma levels (see *Drug Interactions (7.2)*).

5.13 Lipid Elevations

Treatment with efavirenz has resulted in increases in the concentration of total cholesterol and triglycerides. Cholesterol and triglyceride testing should be performed before initiating efavirenz therapy and at periodic intervals during therapy.

5.14 Bone Effects

Bone Mineral Density In clinical trials in HIV-1 infected adults, tenofovir disoproxil fumarate was associated with slightly greater decreases in bone mineral density (BMD) and increases in biochemical markers of bone metabolism, suggesting increased bone turnover relative to comparators. These changes were generally not progressive over 144 weeks in clinical trials. Clinical relevance of these findings has not been established.

Clinical trials evaluating tenofovir disoproxil fumarate in pediatric and adolescent subjects were conducted. Under normal circumstances, BMD increases rapidly in pediatric patients. In HIV-1 infected subjects aged 2 years to less than 18 years, bone effects were similar to those observed in adult subjects and suggest increased bone turnover. Serum parathyroid hormone levels and 1,25 vitamin D levels were higher in tenofovir disoproxil fumarate treated subjects compared to the control groups. Similar trends were observed in chronic hepatitis B infected adolescent subjects aged 12 years to less than 18 years. In all pediatric trials, skeletal growth (height, BMI) was observed (see *Adverse Reactions (6.1)*).

Tenofovir Disoproxil Fumarate

Immune System Disorders: allergic reaction, including angioedema.

Respiratory, Thoracic, and Mediastinal Disorders: dyspnea.

Gastrointestinal Disorders: pancreatitis, increased amylase, abdominal pain.

Renal and Urinary Disorders: renal insufficiency, Acute renal failure, renal failure, acute tubular necrosis, Fanconi syndrome (renal tubular injury with severe hypophosphatemia), proximal renal tubulopathy, interstitial nephritis, renal insufficiency, increased creatinine, proteinuria, polyuria (see *Warnings and Precautions (5.6)*)

Hepatobiliary Disorders: hepatic steatosis, hepatitis, increased liver enzymes (most commonly AST, ALT, gamma GT).

Skin and Subcutaneous Tissue Disorders: rash.

Musculoskeletal and Connective Tissue Disorders: rhabdomyolysis, osteomalacia (manifested as bone pain and which may contribute to fractures), muscular weakness, myopathy.

General Disorders and Administration Site Conditions: asthenia.

The following adverse reactions, listed under the body system headings above, may occur as a consequence of Fanconi syndrome: osteomalacia, hypophosphatemic osteomalacia, hypokalemia, muscular weakness, myopathy.

7 DRUG INTERACTIONS

7.1 Not Recommended with Other Antiretroviral Medications

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are a complete regimen for the treatment of HIV-1 infection; therefore, they should not be administered with other antiretroviral medications for treatment of HIV-1 infection.

7.2 UP Protegrating Drugs

There is limited information available on the potential for a pharmacokinetic interaction between efavirenz and drugs that prolong the QTc interval. QTc prolongation has been observed with the use of efavirenz (see *Clinical Pharmacology (12.2)*). Consider alternatives to efavirenz when coadministered with a drug with a known risk of torsades de pointes.

7.3 Drugs Affecting Renal Function

Since tenofovir is primarily eliminated by the kidney (see *Clinical Pharmacology (12.3)*), coadministration of efavirenz, lamivudine and tenofovir disoproxil fumarate with drugs that reduce renal function or compete for active tubular secretion may increase serum concentrations of tenofovir and/or increase the concentrations of other renally eliminated drugs. Coadministration of tenofovir disoproxil fumarate with other drugs that are eliminated by active tubular secretion (e.g., gentamicin), and high-dose or multiple NSAIDs (see *Warnings and Precautions (5.6)*).

7.4 Cannabinoid Test Interactions

Efavirenz does not bind to cannabinoid receptors. False-positive urine cannabinoid test results have been observed in uninfected volunteers who received efavirenz when using some assay methods during efavirenz. Confirmation of positive screening tests for cannabinoids by a more specific method is recommended.

7.5 Drug-Drug Interactions

Efavirenz has been shown *in vivo* to induce CYP3A4 and CYP2B6. Other compounds that are substrates of CYP3A4 or CYP2B6 may have decreased plasma concentrations when coadministered with efavirenz.

Drugs that induce CYP3A4 activity (e.g., phenobarbital, rifampin, rifabutin) would be expected to increase the clearance of efavirenz, thereby decreasing efavirenz plasma concentrations.

No drug interactions studies have been conducted using efavirenz, lamivudine and tenofovir disoproxil fumarate. However, drug interaction studies have been conducted with the individual components efavirenz, lamivudine and tenofovir disoproxil fumarate tablets (see *Clinical Pharmacology (12.3)*).

Drug interactions with efavirenz are summarized in Table 3 (for pharmacokinetic data see *Clinical Pharmacology (12.3, Tables 4 and 5)*). The recommendations in this table are based on single-dose studies, but is not all inclusive.

Table 3. Established and Other Potentially Significant Drug Interactions: Alterations in Dose or Regimen May Be Recommended Based on Drug Interaction Studies or Predicted Interaction

Concomitant Drug Class: Drug Name	Effect	Clinical Comment
Anticonvulsant: Carbamazepine	↑ or ↓ or ↔efavirin	Monitor INR and adjust warfarin dosage if necessary.
Antimycobacterial: Carbamazepine	↓ carbamazepine* ↓ efavirenz*	Data are insufficient data to make a dose recommendation for efavirenz. Alternative antimycobacterial agents should be considered.
Rifampin Rifabutin	↓ anticonvulsant ↓ efavirenz	Monitor anticonvulsant plasma levels particularly for those anticonvulsants with a narrow therapeutic index and administer dose adjustments.
Antidepressants: Bupropion Sertraline Fluoxetine Paroxetine	↓ bupropion* ↓ sertraline* ↓ fluoxetine* ↓ paroxetine*	Increases in bupropion dosage should be guided by clinical response, but the maximum recommended dose. Increases in sertraline dosage should be guided by clinical response. Avoid concomitant use unless the benefit outweighs the risk.
Anti-infective: Clarithromycin	↑ clarithromycin*	Consider alternatives to macrolide antibiotics because of risk of QTc prolongation.
Antifungal: Voriconazole	↓ voriconazole*	Coadministration is not recommended.
Antimycobacterial: Rifabutin	↓ rifabutin*	Concomitant administration is not recommended.
Antiretrovirals: Atazanavir/ritonavir Lopinavir/ritonavir Ritonavir Atovaquone/ritonavir Nelfinavir	↓ atazanavir* ↓ efavirenz* ↓ lopinavir* ↓ efavirenz* ↓ ritonavir* ↓ efavirenz* ↓ nelfinavir* ↓ efavirenz*	Concomitant administration is not recommended.
Cancer chemoattenuators: Cyclophosphamide Ifosfamide Vinca alkaloids	↓ efavirenz* ↓ ifosfamide* ↓ vinca alkaloids* (e.g., vinblastine, vincristine)	Clinical dose adjustments should be guided by clinical response (refer to the full prescribing information for efavirenz).
Calcium channel blockers: Diltiazem Verapamil Nifedipine Nicardipine	↓ calcium channel blocker	Where coadministered with efavirenz, dosage adjustment of calcium channel blocker may be needed and should be guided by clinical response (refer to the full prescribing information for calcium channel blockers).
Others (e.g., felodipine, nisoldipine, verapamil)	↓ calcium channel blocker*	Plasma concentrations of atovaquone, proguanil and simvastatin decreased. Consult the full prescribing information for atovaquone/proguanil for guidance on administering the dose.
GABA-Cob reduction inhibitor: Divalproex Finasteride Valproic acid	↓ phenobarbital* ↓ phenytoin* ↓ valproic acid*	Plasma concentrations of atovaquone, proguanil and simvastatin decreased. Consult the full prescribing information for atovaquone/proguanil for guidance on administering the dose.
Hepatitis C antiviral agents: Boceprevir Telaprevir	↓ boceprevir* ↓ telaprevir*	Concomitant administration of efavirenz is not recommended.
Herbal products: St. John's wort (Hypericum perforatum)	↓ efavirenz*	Concomitant administration of efavirenz is not recommended.
Hepatitis B antiviral agents: Adefovir dipivoxil	↑ lamivudine/tenofovir disoproxil fumarate	Coadministration of adefovir dipivoxil is not recommended.
Hepatitis B antiviral agents: Telbivudine	↓ telbivudine*	Concomitant administration of telbivudine is not recommended.
Immunosuppressants: Cyclosporine, tacrolimus, sirolimus, everolimus, metabolized by CYP3A	↓ immunosuppressant	Dose adjustments of the immunosuppressant may be required. Close monitoring of immunosuppressant concentrations for at least 2 weeks (until stable concentrations) after initiation, dose adjustment, or discontinuation of efavirenz is warranted.
Narcotic analgesic: Methadone	↓ methadone*	Monitor for signs of withdrawal, adjust dose and increase methadone dose if required to alleviate withdrawal symptoms.

↑ = increase, ↓ = decrease, ↔ = no change, * = see text

- The interaction between efavirenz and the drug was evaluated in a clinical study. All other drug interactions observed are clinically mild.

7.6 Drugs without Clinically Significant Interactions

No clinically significant drug interactions have been observed when lamivudine, and tenofovir disoproxil fumarate tablets are administered with the following: atazanavir/lopinavir tipranavir/ritonavir, atorvastatin, azithromycin, cetirizine, famotidine, fluconazole.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Efavirenz:

Pregnancy Exposure Registry: There are retrospective case reports of neural tube defects in infants whose mothers were exposed to efavirenz-containing regimens in the first trimester of pregnancy. Prospective pregnancy data from the Antiretroviral Pregnancy Registry (APR) on 766 pregnancies with first trimester exposure to efavirenz-containing regimens (600 mg daily dose) did not show an increased risk of overall birth defects. The APR has received reports of 19 infants with first trimester exposure to efavirenz and birth defects of 2.7% in the U.S. reference population of the Metropolitan Atlanta Congenital Defects Program (MACDP). Although the sample size is not sufficient to exclude small differences in risk, the data from the APR do not suggest that the overall risk of birth defects in humans exposed to efavirenz in the first trimester is greater than the background birth defect rate of 2.7% in the U.S. reference population of the Metropolitan Atlanta Congenital Defects Program. As of the January 2007 report, there were 14 prospectively reported cases of neural tube defects among these 766 exposures (1.8% [95% CI: 0.9%, 3.0%]). Two cases of neural tube defects were reported among 104 first trimester exposures to efavirenz-containing regimens (including one fetal loss) in the Ibadan cohort but one of these cases was not confirmed by examination. There were no malformed fetuses or infants from placebo-treated mothers. The malformations that occurred in these three cases included anencephaly and cephalocele (exencephaly) in one fetus, micrognathia in a second, and cleft palate in the third. There was no MSNG, the observable adverse effect level for efavirenz in animals (see *Nonclinical Toxicology (13.1)*). Although the mechanism of action is unknown, efavirenz may interfere with folate metabolism. Periconceptional use of folic acid supplements has been shown to decrease the risk of neural tube defects in the general population. Women should be informed that these common symptoms were likely to improve with continued therapy and were not predictive of subsequent onset of the less frequent psychiatric symptoms (see *Warnings and Precautions (5.7)*). Dosing of bedtime may improve the tolerability of these nervous system symptoms (see *Dosage and Administration (2.1)*).

Analysis of long-term data from Study 006 (median follow-up 180 weeks, 100 weeks, and 76 weeks for patients treated with efavirenz + lamivudine + stavudine, efavirenz + indinavir, and indinavir + zidovudine + lamivudine, respectively) showed that, beyond the 4 weeks of therapy, the incidence of new nervous system symptoms among efavirenz-treated patients were generally similar to those in the indinavir-containing control arm.

Efavirenz can cause dizziness in patients who receive it. Advise patients to avoid potentially hazardous tasks such as driving or operating machinery until they are familiar with the potential effects of efavirenz on mental alertness and coordination.

Patients who experience central nervous system symptoms such as dizziness, impaired concentration, and/or drowsiness should avoid potentially hazardous tasks such as driving or operating machinery.

5.9 Embryo-Fetal Toxicity

Efavirenz, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, may cause fetal harm when administered during the first trimester to a pregnant woman. Advise females of reproductive potential who are receiving efavirenz to avoid pregnancy. (*See Use in Specific Populations (8.1)*.)

5.10 Rash

In controlled clinical trials, 26% (266/1008) of patients treated with 600 mg efavirenz experienced new-onset skin rash compared with 17% (117/682) of patients treated in control groups. Rash associated with blistering, moist desquamation, or ulceration occurred in 0.9% (9/1008) of patients treated with efavirenz. The incidence of rash (e.g., erythema multiforme, Stevens-Johnson syndrome) was highest during the first 1 to 2 weeks of therapy and expanded scores was 9.1%. Rashes are usually mild-to-moderate maculopapular skin eruptions that occur within the first 2 weeks of initiation of therapy with efavirenz and resolve spontaneously within 4 weeks. In clinical trials, in most patients continuing therapy with efavirenz, rash resolves within 1 month (median duration, 16 days). The discontinuation rate for rash in clinical trials was 1.7% (17/1008).

Efavirenz can generally be reinitiated in patients interrupting therapy because of rash. Efavirenz should be discontinued in patients developing severe rash associated with blistering, desquamation, mucosal involvement, or fever. Appropriate antihistamines and/or corticosteroids may improve the tolerability and hasten the resolution of rash. For patients who have had a life-threatening cutaneous reaction (e.g., Stevens-Johnson syndrome), alternative therapy should be considered. (see *Contraindications (4.1)*)

Rash was reported in 5% of 3 (3/57) pediatric patients (40%) treated with efavirenz capsules, patient experienced Grade 3 rash (confluent rash with fever), and two patients had Grade 4 rash (erythema multiforme). The median time to onset of rash in pediatric patients was 8 days. Prophylaxis with appropriate antihistamines before initiating therapy with

DESTAQUES DAS INFORMAÇÕES DE PRESCRIÇÃO

Estes destaques não incluem todas as informações necessárias para usar o EFAVIRENZ, LAMIVUDINE E TENOFOVIR DISOPROIL FUMARATO em comprimidos de forma segura e eficaz. Veja bulas de informações completas de prescrição para comprimidos de EFAVIRENZ, LAMIVUDINA E TENOFOVIR DISOPROIL FUMARATO.

Comprimidos de EFAVIRENZ, LAMIVUDINA e TENOFOVIR DISOPROIL FUMARATO para uso oral

Aprovação inicial: 2007

ATENÇÃO: ACIDOSE LÁTICA/HEPATOMEGALIA GRAVE COM ESTEATOSE e EXACERBAÇÕES AGUDAS DA HEPATITE B PÓS-TRATAMENTO

Veja bula e informação completa em seu.

- **Acidose lática e hepatomegalia com esteatose, incluindo casos fatais, foram relatados com o uso de análogos de nucleósidos, incluindo lamivudina e tenofovir disoproil fumarato. (5.1)**
- **Foram identificadas exacerbações graves agudas da hepatite B em pacientes coinfectados com vírus da hepatite B (HBV) e vírus da imunodeficiência humana (HIV) após descontinuação do tratamento com tenofovir disoproil. Monitorar a função hepática em pelo menos pacientes, e, se apropriado, iniciar o tratamento anti-hepatite B. (5.2)**

INDICAÇÕES E USO

Os comprimidos de efavirenz, lamivudina e tenofovir disoproil fumarato são indicados como terapia antiretroviral combinada para o tratamento de infecção pelo vírus da imunodeficiência humana (HIV-1) em adultos e em pacientes pediátricos que pesam pelo menos 35 kg. (1)

DOSAGEM E ADMINISTRAÇÃO

- Coadministração com outros produtos: não use com outros produtos que contenham efavirenz, lamivudina, tenofovir ou produtos que contenham entricitabina. Não administrar juntamente com adefovir dipivoxil. (2.3)
- Recomenda-se que a depuração estimada de creatinina seja avaliada antes de iniciar a terapia, que pacientes que recebem drogas de eliminação renal concorrentes e regimes baseados em interferão e lamivudina sejam monitorados. Descontinue os comprimidos de efavirenz, lamivudina e tenofovir disoproil fumarato em pacientes que desenvolvem sinais ou sintomas de nefrotoxicidade relacionada à interferão alfa, efavirenz ou lamivudina. (2.4)
- Rancovirib, use com cautela em pacientes pediátricos com histórico de pancreatite ou outros fatores de risco para pancreatite. Descontinue em pacientes que desenvolvem pancreatite. Não é recomendado o tratamento de pacientes pediátricos com clearance estimado apropriado. (2.5)
- Nunca inicie os Pivots da lamivudina Renal: pode ocorrer insuficiência renal aguda e síndrome de Fanconi. Em pacientes com depuração de creatinina estimada abaixo de 50 mL/min, descontinue o tratamento com os comprimidos dos componentes de efavirenz, lamivudina e tenofovir disoproil fumarato. Em pacientes com risco de disfunção renal, calcular a depuração estimada de creatinina. Nunca administrar a pacientes com depuração estimada de creatinina abaixo de 50 mL/min, a menos que o benefício justifique o risco. Monitorar a função renal em todos os pacientes. (2.6)

DOSAGEM

Comprimidos: 600 mg de efavirenz, 300 mg de lamivudina e 300 mg de fumarato de tenofovir disoproil. (2)

CONTRAINDICAÇÕES

- Os componentes de efavirenz, lamivudina e tenofovir disoproil fumarato estão contraindicados a pacientes com hipersensibilidade a qualquer componente. Reações graves, incluindo anafilaxia, angioedema e síndrome de Stevens-Johnson e qualquer das categorias de seus produtos. (4)

ANÁLISES E PRECAUÇÕES

- Coadministração com outros produtos: não use com outros produtos que contenham efavirenz, lamivudina, tenofovir ou produtos que contenham entricitabina. Não administrar juntamente com adefovir dipivoxil. (2.3)
- Recomenda-se que a depuração estimada de creatinina seja avaliada antes de iniciar a terapia, que pacientes que recebem drogas de eliminação renal concorrentes e regimes baseados em interferão e lamivudina sejam monitorados. Descontinue os comprimidos de efavirenz, lamivudina e tenofovir disoproil fumarato em pacientes que desenvolvem sinais ou sintomas de nefrotoxicidade relacionada à interferão alfa, efavirenz ou lamivudina. (2.4)
- Rancovirib, use com cautela em pacientes pediátricos com histórico de pancreatite ou outros fatores de risco para pancreatite. Descontinue em pacientes que desenvolvem pancreatite. Não é recomendado o tratamento de pacientes pediátricos com clearance estimado apropriado. (2.5)
- Nunca inicie os Pivots da lamivudina Renal: pode ocorrer insuficiência renal aguda e síndrome de Fanconi. Em pacientes com depuração de creatinina estimada abaixo de 50 mL/min, descontinue o tratamento com os componentes dos comprimidos de efavirenz, lamivudina e tenofovir disoproil fumarato. Em pacientes com risco de disfunção renal, calcular a depuração estimada de creatinina. Nunca administrar a pacientes com depuração estimada de creatinina abaixo de 50 mL/min, a menos que o benefício justifique o risco. Monitorar a função renal em todos os pacientes. (2.6)

INFORMAÇÕES COMPLETAS DE PRESCRIÇÃO: CONTEÚDO*

ATENÇÃO: ACIDOSE LÁTICA/HEPATOMEGALIA GRAVE COM ESTEATOSE e EXACERBAÇÕES AGUDAS DA HEPATITE B PÓS-TRATAMENTO

Acidose lática e hepatomegalia com esteatose, incluindo casos fatais, foram relatados com o uso de análogos de nucleósidos, incluindo lamivudina e tenofovir de tenofovir disoproil. Suspender o tratamento com efavirenz, lamivudina e tenofovir disoproil fumarato em pacientes que desenvolvem sinais ou sintomas de acidose lática ou hepatotoxicidade pronunciada (ver *Advertências e Precauções* (5.1)).

Exacerbações agudas graves da hepatite B foram relatadas em pacientes coinfectados com vírus da hepatite B (HBV) e vírus da imunodeficiência humana (HIV) após descontinuação do tratamento com tenofovir disoproil. Monitorar a função hepática em pelo menos pacientes e, se apropriado, iniciar o tratamento anti-hepatite B (ver *Advertências e Precauções* (5.2)).

1 INDICAÇÕES E USO

Os comprimidos de efavirenz, lamivudina e tenofovir disoproil fumarato são indicados para uso isolado como terapia antiretroviral combinada para o tratamento de infecção pelo vírus da imunodeficiência humana (HIV-1) em adultos e em pacientes pediátricos que pesam pelo menos 35 kg.

2 DOSAGEM E ADMINISTRAÇÃO

2.1 Dosagem recomendada para pacientes adultos e pediátricos pesando pelo menos 40 kg

A dose recomendada de comprimidos de efavirenz, lamivudina e tenofovir disoproil fumarato é de 600 mg/300 mg/300 mg administrada por via oral em dose única diária. Os comprimidos de efavirenz, lamivudina e tenofovir disoproil fumarato devem ser tomados em jejum, de preferência na hora de dormir. A dose nunca deve ser alterada ou reduzida (interrupção) dos sistemas de ingesta ou terapia (ver *Advertências e Precauções* (5.2)). Regular Adverse Reactions (ver *Advertências e Precauções* (5.1)). Se for necessário descontinuar um dos componentes de efavirenz, lamivudina e tenofovir disoproil fumarato devem ser avaliados ritmicos com rigidez.

2.2 Ajuste da Dose para Insuficiência Renal

Como o efavirenz, lamivudina e tenofovir disoproil fumarato é um componente de combinação de dose fixa, não é recomendado para pacientes com insuficiência renal (depuração de creatinina inferior a 50 mL/min) os pacientes que necessitam de ajuste de dose (ver *Farmacologia Clínica* (12.3)).

3 FORMAS DE DOSAGEM

Os Comprimidos de Fumarato, Disoproil, Efavirenz, Lamivudina e Tenofovir estão disponíveis contendo 600 mg de efavirenz, 300 mg de lamivudina, USP e 300 mg de fumarato de tenofovir disoproil equivalente a 245 mg de tenofovir disoproil.

Os comprimidos de 600 mg/300 mg/300 mg são cápsulas compridas, revestidas, no formato de cápsula, por película bicôncava de cor amarela, com 1 de um lado e "321" no outro.

4 CONTRAINDICAÇÕES

4.1 Hipersensibilidade

Os componentes de efavirenz, lamivudina e tenofovir disoproil fumarato são contraindicados a diretrizes com uma reação de hipersensibilidade a qualquer dos componentes destes comprimidos.

5 AVISOS E PRECAUÇÕES

5.1 Acidose Lática/Hepatomegalia Grave com Esteatose

Acidose lática e hepatomegalia grave com esteatose, incluindo casos fatais, foram relatados com o uso de análogos de nucleósidos e outros antirretrovirais. A maioria destes casos ocorreram em mulheres. A obesidade e a exposição por curto tempo aos nucleosídeos podem ser fatores de risco. Deve-se ter cuidado ao administrar análogos de nucleósidos a qualquer paciente com fatores de risco conhecidos para doença hepática. No entanto, também foram relatados casos em pacientes sem fatores de risco conhecidos. Deve-se suspender o tratamento com efavirenz, lamivudina e tenofovir em qualquer paciente que desenvolva achados clínicos ou laboratoriais sugestivos de acidose lática ou hepatotoxicidade pronunciada (que pode incluir hepatomegalia e esteatose, mesmo na ausência de elevações marcadas de aminotransferases).

5.2 Pacientes Coinfectados com HIV-1 e HBV

A descontinuação por todos os pacientes com HIV-1 agem tentados quanto à presença de vírus crônicos da hepatite B (VHB) antes de iniciar a terapia antirretroviral. A interrupção da terapia anti-VHB, incluindo lamivudina e tenofovir disoproil fumarato, pode estar associada a exacerbações graves da hepatite. Os pacientes coinfectados com VHB e HIV-1 que param de tomar os comprimidos de efavirenz, lamivudina e tenofovir disoproil fumarato devem ser cuidadosamente monitorizados com acompanhamento clínico e laboratorial durante, pelo menos, vários meses após a interrupção do tratamento. Se apropriado, a retomada da terapia antihepatite B pode ser justificada. *Diferenças importantes entre os Pivotos por ambos Lamivudina:* os comprimidos de lamivudina contêm para o tratamento da infecção crônica pelo vírus da hepatite B (VHB) e não foram estudados em pacientes com VHB coinfectados para com hepatite B crônica. A dosagem e a dosagem de lamivudina em 300 mg/dia não são adequadas para pacientes coinfectados com HIV-1 e VHB. A segurança e a eficácia de lamivudina não foram estabelecidas para o tratamento de VHB descompensada. Os dados disponíveis são insuficientes para que a terapia com EFAVIRENZ 600 mg produza contenda tenofovir disoproil fumarato, como o VIREAD, for prescrito para hepatite B crônica para um paciente com infecção não reconhecida ou não tratada com HIV-1, a emergência de resistência ao HIV-1 pode ser desenvolvida. Não é recomendado o tratamento de pacientes com infecção não reconhecida ou não tratada com HIV-1 com a dosagem de lamivudina com 300 mg/dia.

5.3 Coadministração com Outros Produtos

Os comprimidos de efavirenz, lamivudina e tenofovir disoproil fumarato não um produto contendo de dose fixa de efavirenz, lamivudina e tenofovir de tenofovir disoproil e não devem ser coadministrados concomitantemente com outros produtos que contenham efavirenz, lamivudina, tenofovir ou produtos que contenham entricitabina. EMTRIVA ou EFAVIR-VIAIRO (lamivudina), ATRIZIVIR (sulfato de abacavir e lamivudina), TRIZIVIR (sulfato de abacavir, lamivudina e zidovudina), ATRIPLA (efavirenz, fumarato de tenofovir disoproil e entricitabina), TRUVIVIRD (sulfato de abacavir e lamivudina), COMPLERA (rilpivirina/emtricitabina/fumarato de tenofovir disoproil), STRIBILD (elvitegravir/cobicistate/emtricitabina/fumarato de tenofovir disoproil).

Os comprimidos de efavirenz, lamivudina e tenofovir disoproil fumarato não devem ser administrados de forma associada ao HEPSERA® (adefovir dipivoxil) (ver *Interações Medicamentosas* (7.2)).

5.4 Uso com Regimes Baseados em Interferão e Ribavirina

Estudos in vitro mostraram que a ribavirina pode reduzir a fosforilação de análogos de nucleósidos de primidina, como a lamivudina, um componente dos comprimidos de efavirenz, lamivudina e fumarato de tenofovir disoproil. Esta observação foi feita em células Molt-4 clone 8 e em células CEM, mas não em células Vero. Não foi estabelecida a relevância clínica desta observação. Foram relatados casos de acidose lática e disfunção hepática grave em pacientes coinfectados com HIV-1/HIV-2/HIV-Farmacologia Clínica (12.3)) descompensação hepática (ligados ou não a acidose lática) em pacientes coinfectados com HIV-1/HIV-2 e hepatite B ou hepatite C tratados com HIV-1 e interferão alfa com ou sem ribavirina. Os pacientes que recebem interferão alfa com ou sem ribavirina devem ser cuidadosamente monitorizados quanto à toxicidade associada ao tratamento, especialmente descompensação hepática. Também deve ser considerada a interrupção do tratamento se houver evidência de piora clínica significativa, particularmente se houver aumento da hepatotoxicidade em qualquer um dos medicamentos usados na terapia combinada. Não foram estabelecidas as dosagens de lamivudina para uso em

Size : 500 x 950 mm
Book Folding : 45 x 60 mm
Color : Black
Spec: Printed on 40 GSM Bible paper, front & back side printing.

Pharma Code : 4646 4647

Note: Pharma code position, Orientation & Product Name Size are tentative, will be change based on folding size

Efavirenz, Lamivudine and Tenofovir Disoproxil Fumarate Tablets 200/300/300 mg

Rx Only

Revised: 06/2017

HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all the information needed to use EFAVIRENZ, LAMIVUDINE AND TENOFOVIR DISOPROXIL FUMARATE TABLETS safely and effectively. See full prescribing information for EFAVIRENZ, LAMIVUDINE AND TENOFOVIR DISOPROXIL FUMARATE TABLETS.

EFAVIRENZ, LAMIVUDINE and TENOFOVIR DISOPROXIL FUMARATE tablets, for oral use
Initial U.S. Approval: 2006

WARNING: LACTIC ACIDOSIS/SEVERE HEPATOMEGALY WITH STEATOSIS and POST TREATMENT ACUTE EXACERBATIONS OF HEPATITIS B
See full prescribing information for complete boxed warning.
- **Lactic acidosis and hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogues including lamivudine and tenofovir disoproxil fumarate. (5.1)**
- **Severe acute exacerbations of hepatitis B have been reported in patients who are co-infected with HIV-1 and hepatitis B virus and have discontinued FTC, 3TC or TDF, components of EFAVIRENZ, LAMIVUDINE and tenofovir disoproxil fumarate. Monitor hepatic function closely in these patients and, if appropriate, initiate anti-hepatitis B treatment. (5.2)**

INDICATIONS AND USAGE
Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, a combination of one non-nucleoside reverse transcriptase inhibitor (efavirenz) and two nucleoside reverse transcriptase inhibitors (lamivudine and tenofovir disoproxil fumarate), are indicated for use alone as a complete regimen for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in adult patients with no prior antiretroviral use. (1)

DOSAGE AND ADMINISTRATION
- Recommended dose: One tablet (containing 600 mg of efavirenz, 300 mg of lamivudine and 300 mg of tenofovir disoproxil fumarate) taken once daily orally on an empty stomach, preferably at bedtime. (2.1)

DOSAGE FORMS AND STRENGTHS
Tablets: 600 mg of efavirenz, 300 mg lamivudine and 300 mg tenofovir disoproxil fumarate. (3)

CONTRAINDICATIONS
- Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are contraindicated in patients with previous hypersensitivity (e.g., Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to any of the components of this product. (4.1)

WARNINGS AND PRECAUTIONS
- Coadministration with Other Products: Do not use with other efavirenz, lamivudine or tenofovir-containing products. Do not administer with lamivudine as the only component of a combination product. (5.3)
- Hepatic decompensation, some fatal, has occurred in HIV-1/HCV co-infected patients receiving combination antiretroviral therapy and may occur in patients without pre-existing liver disease or other identifiable risk factors. Discontinue efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, as medically appropriate and consider dose reduction or discontinuation of interferon alfa, ribavirin, or both. (5.4)
- Mitochondrial Dysfunction after Exposure in Utero: Nucleoside analogs may cause delayed mitochondrial dysfunction syndromes. Monitor pediatric patients exposed in utero to nucleoside analogs and consider appropriate diagnostic evaluation. (5.5)
- New Onset or Worsening Renal Impairment: Can include acute renal failure and Fanconi syndrome. Assess estimated creatinine clearance before initiating treatment with tenofovir disoproxil fumarate, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Avoid use of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets with concurrent or recent use of nephrotoxic drugs. (5.6)

FULL PRESCRIBING INFORMATION: CONTENTS*
WARNING: LACTIC ACIDOSIS/SEVERE HEPATOMEGALY WITH STEATOSIS and POST TREATMENT ACUTE EXACERBATIONS OF HEPATITIS B
1 INDICATIONS AND USAGE
2 DOSAGE AND ADMINISTRATION
2.1 Recommended Dosage for Adult Patients and Pediatric Patients (40 kg or More)
2.2 Dose Adjustment for Renal Impairment
3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
4.1 Hypersensitivity
5 WARNINGS AND PRECAUTIONS
5.1 Lactic Acidosis/Severe Hepatomegaly With Steatosis
5.2 Patients Coinfected with HIV-1 and HBV
5.3 Coadministration with Other Products
5.4 Use with Interferon- and Ribavirin-Based Regimens
5.5 Pancreatitis
5.6 New Onset or Worsening Renal Impairment
5.7 Psychiatric Symptoms
5.8 Nervous System Symptoms
5.9 Embryo-fetal Toxicity
5.10 Rash
5.11 Hepatotoxicity
5.12 Convulsions
5.13 Lipid Elevations
5.14 Bone Effects
5.15 Immune Reconstitution Syndrome
5.16 Fat Redistribution
6 ADVERSE REACTIONS
6.1 Clinical Trials Experience
6.2 Postmarketing Experience

FULL PRESCRIBING INFORMATION

WARNING: LACTIC ACIDOSIS/SEVERE HEPATOMEGALY WITH STEATOSIS and POST TREATMENT ACUTE EXACERBATIONS OF HEPATITIS B

Lactic acidosis and hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogues including lamivudine and tenofovir disoproxil fumarate. Suspend treatment if clinical or laboratory findings suggestive of lactic acidosis or pronounced hepatotoxicity occur *(see Warnings and Precautions (5.1)]*.

Severe acute exacerbations of hepatitis B have been reported in patients who are co-infected with hepatitis B virus (HBV) and HIV-1 and have discontinued FTC, 3TC or TDF, components of efavirenz, lamivudine and tenofovir disoproxil fumarate. Monitor hepatic function closely in these patients and, if appropriate, initiate anti-hepatitis B treatment *(see Warnings and Precautions (5.2)]*.

1 INDICATIONS AND USAGE

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are indicated for use alone as a complete regimen for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in adult patients weighing at least 40 kg.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage for Adult and Pediatric Patients weighing at least 40 kg

The recommended dosage of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets is 600 mg/300 mg/300 mg orally, once daily. It is recommended that efavirenz, lamivudine and tenofovir disoproxil fumarate tablets be taken on an empty stomach, preferably at bedtime. Administration with food may result in increased efavirenz exposure and increased adverse reactions *(see Clinical Pharmacology and Pharmacokinetics (12.3), Adverse Reactions (6.1) and Patient Counseling Information (17)]*. Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets should be swallowed intact with liquid.

2.2 Dose Adjustment for Renal Impairment

Because efavirenz, lamivudine and tenofovir disoproxil fumarate is a fixed-dose combination tablet, it is not recommended for patients with estimated creatinine clearance less than 50 mL/min or pediatric patients and end-stage renal disease (ESRD) requiring hemodialysis.

3 DOSAGE FORMS AND STRENGTHS

Efavirenz, Lamivudine and Tenofovir Disoproxil Fumarate Tablets are available containing 600 mg of efavirenz, 300 mg of lamivudine, USP and 300 mg tenofovir disoproxil fumarate equivalent to 245 mg of tenofovir disoproxil.

- The 600 mg/300 mg/300 mg tablets are yellow colored, capsule-shaped, bevel edged biconvex film coated tablets debossed with "1" on one side and "227" on the other side.

4 CONTRAINDICATIONS

4.1 Hypersensitivity

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are contraindicated in patients with a previous hypersensitivity reaction to any of the components contained in the formulation.

5 WARNINGS AND PRECAUTIONS

5.1 Lactic Acidosis/Severe Hepatomegaly With Steatosis

Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogues including lamivudine and tenofovir. A majority of these cases have been in women. Obesity and prolonged nucleoside exposure may be risk factors. Caution should be exercised when administering any nucleoside analogs to any patient with known risk factors for liver disease; however, cases have also been reported in patients with no known risk factors. Treatment should be suspended in any patient who develops clinical or laboratory findings suggestive of lactic acidosis or pronounced hepatotoxicity (which may include hepatomegaly and steatosis even in the absence of marked transaminase increases).

5.2 Patients Coinfected with HIV-1 and HBV

It is recommended that all patients with HIV-1 be tested for the presence of chronic hepatitis B virus (HBV) before initiating antiretroviral therapy. Discontinuation of anti-HBV therapy, including lamivudine and tenofovir disoproxil fumarate, may be associated with severe acute exacerbations of hepatitis. Patients infected with HBV who discontinue efavirenz, lamivudine and tenofovir disoproxil fumarate tablets should be closely monitored with both clinical and laboratory follow-up for at least several months after stopping treatment. If appropriate, resumption of anti-hepatitis B therapy may be warranted.

Reported Differences Among Lamivudine-Containing Products: Lamivudine contains a higher dose of the same active ingredient (lamivudine) than EPIVIR-HBV tablets. EPIVIR-HBV was developed for patients with chronic hepatitis B. The formulation and dosage of lamivudine tablets in EPIVIR-HBV are not appropriate for co-infected patients with HIV-1 and HBV. Safety and effectiveness of lamivudine have not been established for treatment of chronic hepatitis B in patients who are co-infected with HIV-1.

If treatment with EPIVIR-HBV or tenofovir disoproxil fumarate-containing product such as VIREAD is prescribed for chronic hepatitis B for a patient with unrecognized or untreated HIV-1 infection, rapid emergence of HIV-1 resistance is likely to result because of the subtherapeutic dose and the inappropriateness of monotherapy HIV-1 treatment.

5.3 Coadministration with Other Products

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are a fixed-dose combination product of efavirenz, lamivudine and tenofovir disoproxil fumarate. Do not administer efavirenz, lamivudine and tenofovir disoproxil fumarate (including lamivudine-containing, or emtricitabine-containing drugs), emtricitabine (Emtriva®, Coviracil®), EPIVIR® or EPIVIR-HBV® (lamivudine), EPZICOM® (abacavir sulfate/lamivudine), TRIZIVIR® (abacavir sulfate/lamivudine/zidovudine), METHADOSE® (methadone), SUSTIVA® (efavirenz) or delavirdine mesylate, VIREAD® (tenofovir disoproxil fumarate), ATRIPLA® (emtricitabine/efavirenz/tenofovir disoproxil fumarate), COMBIVIR® (lamivudine/zidovudine), or EMTRIVA® (emtricitabine) with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets.

Tenofovir disoproxil fumarate and lamivudine should not be administered in combination with HEPSERA® (adefovir dipivoxil) *(see Drug Interactions (7.2)]*.

5.4 Use With Interferon- and Ribavirin-Based Regimens

In vitro studies have shown ribavirin can reduce the phosphorylation of pyrimidine nucleoside analogues such as lamivudine, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Although no evidence of an interaction between ribavirin and efavirenz, lamivudine or tenofovir has been observed, caution should be exercised when ribavirin was coadministered with lamivudine or HIV-1/HCV co-infected patients *[see Clinical Pharmacology (12.3)]* hepatic decompensation (some fatal) has occurred in HIV-1/HCV co-infected patients receiving combination antiretroviral therapy and may occur in patients without pre-existing liver disease or other identifiable risk factors. Discontinuation of efavirenz with ribavirin, or both should also be considered if worsening clinical hepatic decompensation. Discontinuation of lamivudine should be considered as medically appropriate. Dose reduction or discontinuation of interferon alfa, ribavirin, or both should also be considered if worsening clinical toxicities are observed, including hepatic decompensation (e.g., Child-Pugh-c). See the full prescribing information for interferon and ribavirin.

5.5 Pancreatitis

In pediatric patients with a history of prior antiretroviral nucleoside exposure, a history of pancreatitis, or other significant risk factors for the development of pancreatitis, lamivudine should be used with caution. Treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets should be stopped immediately if clinical signs, symptoms, or laboratory abnormalities suggestive of pancreatitis occur.

5.6 New Onset or Worsening Renal Impairment

Tenofovir, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets is principally eliminated by the kidney. Renal impairment, including cases of acute renal failure and Fanconi syndrome (renal tubular injury with severe hypophosphatemia), has been reported with the use of tenofovir disoproxil fumarate *[see Clinical Pharmacology (12.3)]*.

It is recommended that estimated creatinine clearance be assessed in all patients prior to initiating therapy and as clinically appropriate during therapy with tenofovir disoproxil fumarate. In patients at risk of renal dysfunction, including patients who have previously experienced renal events while receiving HEPSERA® (adefovir dipivoxil), it is also recommended that the estimated creatinine clearance be assessed, and serum phosphorus be monitored, prior to initiation of efavirenz, lamivudine and tenofovir disoproxil fumarate, and periodically during tenofovir disoproxil fumarate therapy.

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets should be avoided with concurrent or recent use of a nephrotoxic agent (e.g., high-dose or multiple non-steroidal anti-inflammatory drugs (NSAIDs)) *[see Drug Interactions (7.2)]*. Cases of acute renal failure after initiation of high-dose or multiple NSAIDs have been reported in HIV-infected patients with risk factors for renal dysfunction who appeared stable on tenofovir disoproxil fumarate. Some patients required hospitalization and renal replacement therapy. Alternatives to NSAIDs should be considered, if needed, in patients at risk for renal dysfunction.

Persistent or worsening bone pain, pain in extremities, fractures and/or muscular pain or weakness may be manifestations of proximal renal tubulopathy and may prompt an evaluation of renal function in at-risk patients.

5.7 Psychiatric Symptoms

Serious psychiatric adverse experiences have been reported in patients treated with efavirenz, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. In controlled trials of 1008 patients treated with regimens containing efavirenz for a mean of 2.1 years and 635 patients treated with control regimens for a mean of 1.5 years, the frequency (regardless of causality) of specific serious psychiatric events among patients exposed efavirenz or control regimens, respectively, were severe depression (2.4%, 0.9%), suicidal ideation (0.7%, 0.3%), nonaggressive hostility directed at others (0.4%, 0.3%), severe manic reactions (0.4%, 0.3%), and severe psychotic reactions (0.8%, 0.3%). When psychiatric symptoms similar to those noted above were considered, or evaluated as a group in a multifactorial analysis of data from Study 006, treatment with efavirenz was associated with a 2- to 3-fold higher risk of these symptoms relative to control. Some of these symptoms appeared to contribute to the occurrence of these psychiatric symptoms were history of injection drug use, psychiatric history, and receipt of psychiatric medication at study entry; similar associations were observed in both the efavirenz and control treatment groups. In Study 006, short of new serious psychiatric symptoms occurred throughout the study for both efavirenz-treated and control-treated patients. One percent of efavirenz-treated patients discontinued therapy.

Interrupted treatment because of one or more of these selected psychiatric symptoms.

There have also been occasional postmarketing reports of death by suicide, delusions, psychosis-like behavior, and catatonia, although a causal relationship to the use of efavirenz cannot be determined from these reports. Patients with serious psychiatric adverse experiences should seek immediate medical evaluation to assess the possibility of suicidality, to establish the diagnosis and identify the role, if any, of efavirenz, and to institute appropriate treatment. Discontinuation of treatment should be considered.

5.8 Nervous System Symptoms

Fifty-three percent (531/1008) of patients receiving efavirenz, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, in controlled trials reported central nervous system symptoms (any grade, regardless of causality) compared to 26% (135/510) of patients receiving control regimens. These symptoms included, but were not limited to, dizziness (9.1% of the 1008 patients), insomnia (8.3%), impaired concentration (6.2%), somnolence (5.2%), and abnormal dreams or vivid dreams (4.2%). Moderate to severe dizziness was seen in 2.7% of patients and 0.4% of patients discontinued therapy as a result. These symptoms usually began during the first or second day of therapy and generally resolve after the first 2 to 4 weeks of therapy. After 4 weeks of therapy, the frequency of these symptoms was similar to that seen in control-treated patients. In Study 006, 19% of patients treated with efavirenz-containing regimens and 2% to 5% in patients treated with control regimens reported these symptoms. Patients should be informed that these common symptoms were likely to improve with continued therapy and were not predictive of subsequent onset of the less frequent psychiatric symptoms *(see Warnings and Precautions (5.7)]*. Dosing of bedtime may improve the tolerability of these nervous system symptoms *(see Dosage and Administration (2.1)]*.

Analysis of long-term data from Study 006 (median follow-up 180 weeks, 100 weeks, and 36 weeks for patients treated with efavirenz + lamivudine + stavudine, efavirenz + indinavir, and indinavir + zidovudine + lamivudine, respectively) showed that, beyond the first 4 weeks of therapy, the incidences of new nervous system symptoms among efavirenz-treated patients were generally similar to those in the indinavir-containing control arm.

Efavirenz has a prolonged half-life (40 to 55 hours). Patients should be informed that dizziness or impaired concentration may occur for an unpredictable period of time. Caution should be exercised when administering efavirenz to patients who operate potentially hazardous machinery, including motor vehicles or heavy equipment.

Patients who experience central nervous system symptoms such as dizziness, impaired concentration, and/or drowsiness should avoid performing such hazardous tasks as driving or operating machinery.

5.9 Embryo-Fetal Toxicity

Efavirenz, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, may cause fetal harm when administered during the first trimester to a pregnant woman. Advise females of reproductive potential who are receiving efavirenz to avoid pregnancy. *(See Use in Specific Populations (8.1).)*

5.10 Rash

In controlled clinical trials, 26% (266/1008) of patients treated with 600 mg efavirenz experienced new-onset skin rash compared with 17% (117/682) of patients treated in control groups. Rash associated with blistering, moist desquamation, or ulceration occurred in 0.9% (9/1008) of patients treated with efavirenz. The majority of cases of rash (a majority of mild to moderate macular papular rash with pruritus) occurred during the first 2 weeks of therapy and resolved spontaneously with continued dosing; the median time to resolution was 29 days. Rashes are usually mild-to-moderate maculopapular skin eruptions that occur within the first 2 weeks of therapy. Efavirenz may be restarted following mild to moderate rash. In clinical trials, in most patients continuing therapy with efavirenz, rash resolves within 1 month (median 16 days). The discontinuation rate for rash in clinical trials was 1.7% (17/1008).

Efavirenz can generally be reinitiated in patients interrupting therapy because of rash. Efavirenz should be discontinued in patients developing severe rash associated with blistering, desquamation, mucosal involvement, or fever. Appropriate antihistamines and/or corticosteroids may improve the tolerability and hasten the resolution of rash. For patients who have had a life-threatening cutaneous reaction (e.g., Stevens-Johnson syndrome), alternative therapy should be considered. *(see Contraindications (4.1))*.

Rash was reported in 5% of 3 hundred and seventy (48/1) treated with efavirenz regimens; patients experienced Grade 3 rash (confluent rash with fever), and two patients had Grade 4 rash (erythema multiforme). The median time to onset of rash in pediatric patients was 8 days. Prophylaxis with appropriate antihistamine before initiating therapy with efavirenz in pediatric patients should be considered.

5.11 Hepatotoxicity

Monitoring of liver enzymes before and during treatment is recommended for patients with underlying hepatic disease, including chronic hepatitis B or C infection, or in patients who have marked elevations of liver enzymes or with other medications associated with liver toxicity *(see Use in Specific Populations (8.6))*. A few of these postmarketing reports of hepatic failure occurred in patients with no pre-existing hepatic disease or other identifiable risk factors. Fatal cases have been reported in patients with pre-existing liver disease or cirrhosis. Patients with pre-existing hepatic dysfunction or chronic liver infections, or patients with persistent elevations of serum transaminases to greater than five times the upper limit of the normal range, the benefit of continued therapy with efavirenz needs to be weighed against the potential risks of significant liver toxicity.

5.12 Convulsions

Convulsions have been observed in patients receiving efavirenz, generally in the presence of known medical history of seizures *[see Nonclinical Toxicology (13.2)]*. Caution should be taken in any patient with a history of seizures. These patients should be monitored during treatment with efavirenz. Concomitant medications may also alter the plasma levels of phenytoin and phenobarbital. If possible, these agents should be avoided during treatment with efavirenz.

5.13 Lipid Elevations

Treatment with efavirenz has resulted in increases in the concentration of total cholesterol and triglycerides. Cholesterol and triglyceride testing should be performed before initiating efavirenz therapy and at periodic intervals during therapy.

5.14 Bone Effects

Bone Mineral Density: In clinical trials in HIV-1 infected adults, tenofovir disoproxil fumarate was associated with slightly greater decreases in bone mineral density (BMD) and increases in biochemical markers of bone metabolism, indicative of bone turnover, than comparators. These changes in bone mineral density were not associated with increased bone pain or fractures in clinical trials. Decreases in BMD and increases in biochemical markers of bone turnover were similar in patients receiving tenofovir disoproxil fumarate or stavudine.

Clinical trials evaluating tenofovir disoproxil fumarate in pediatric and adolescent subjects were conducted. Under normal circumstances, BMD increases rapidly in pediatric patients. In HIV-1 infected subjects aged 2 years to less than 18 years, bone effects were similar to those observed in adult subjects and suggest increased bone turnover. In this population, there was a greater effect on height z-score and greater increases in BMD in the tenofovir disoproxil fumarate group compared to the control group. Similar trends were observed in chronic hepatitis B infected subjects between the aged 12 years to less than 18 years. In all pediatric trials, skeletal growth (defined as height and weight) was evaluated. There was no evidence of clinically significant abnormalities of skeletal growth.

Effects of tenofovir disoproxil fumarate-associated changes in BMD and biochemical markers on long-term bone health are unknown. Assessment of BMD should be considered for adults and pediatric patients who have a history of pathologic bone fracture or other risk factors for osteoporosis or bone loss. Although the effect of supplementation with calcium and vitamin D was not studied, such supplementation may be beneficial for all patients. If bone abnormalities are suspected then appropriate consultation should be obtained.

Osteonecrosis/Avascular Necrosis: Cases of osteonecrosis associated with proximal renal tubulopathy, manifested as bone pain or pain in extremities and which may contribute to fracture, have been reported in association with the use of tenofovir disoproxil fumarate. Osteonecrosis is multifactorial in etiology, and the relative contribution of tenofovir disoproxil fumarate is unknown. Appropriate diagnostic and therapeutic measures should be instituted as clinically indicated. Patients should be considered at risk of renal dysfunction who present with persistent or worsening bone pain, pain in extremities, fractures and/or muscular pain or weakness *(see Warnings and Precautions (5.6))*.

5.15 Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in HIV-infected patients treated with coordination antiretroviral therapy, including efavirenz, lamivudine and tenofovir disoproxil fumarate. During the initial phase of combination antiretroviral treatment, patients whose immune systems respond may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium* infection, cytomegalovirus, *Pneumocystis jirovecii* pneumonia [PCP], or tuberculosis), which may necessitate further evaluation and treatment. Autoimmune disorders (such as Graves' disease, polymyositis, Guillain-Barre syndrome) have also been reported to occur in the setting of immune reconstitution, however, the time to onset is more variable, and can occur many months after initiation of treatment.

5.16 Fat Redistribution

In HIV-infected patients, redistribution/accumulation of body fat including central obesity, dorsocervical fat enlargement (buffalo hump), peripheral wasting, facial wasting, breast enlargement, and "cushingoid appearance" have been observed in patients receiving antiretroviral therapy. The mechanism and long-term health effects of these consequences of these events are currently unknown. A causal relationship has not been established.

5.17 GI Prokinetics

GI prokinetics has been observed with the use of efavirenz *[see Drug Interactions (7.2, 7.6) and Clinical Pharmacology (12.3)]*. Consider alternatives to efavirenz when coadministered with a drug with a known risk of Torsade de Pointes or when administered to patients at higher risk of Torsade de Pointes.

6 ADVERSE REACTIONS

The following adverse reactions are discussed in other sections of the labeling:

- Lactic Acidosis/Severe Hepatomegaly with Steatosis *[see Boxed Warning, Warnings and Precautions (5.1)]*.
- Exacerbations of Hepatitis B *[see Boxed Warning, Warnings and Precautions (5.2)]*.
- Hepatic Decompensation in Patients Co-infected with HIV-1 and Hepatitis C *[see Warnings and Precautions (5.4)]*.
- Pancreatitis *[see Warnings and Precautions (5.5)]*.
- New Onset or Worsening Renal Impairment *[see Warnings and Precautions (5.6)]*.
- Psychiatric Symptoms *[see Warnings and Precautions (5.7)]*.
- Nervous System Symptoms *[see Warnings and Precautions (5.8)]*.
- Rash *[see Warnings and Precautions (5.10)]*.
- Hepatotoxicity *[see Warnings and Precautions (5.11)]*.
- Convulsions *[see Warnings and Precautions (5.12)]*.
- Lipid Elevations *[see Warnings and Precautions (5.13)]*.
- Bone Effects *[see Warnings and Precautions (5.14)]*.
- Immune Reconstitution Syndrome *[see Warnings and Precautions (5.15)]*.
- Fat Redistribution *[see Warnings and Precautions (5.16)]*.

6.1 Clinical Trials Experience

Clinical Trials Experience in Adult Patients with HIV-1 Infection

- Lactic Acidosis/Severe Hepatomegaly with Steatosis *[see Boxed Warning, Warnings and Precautions (5.1)]*.
- Exacerbations of Hepatitis B *[see Boxed Warning, Warnings and Precautions (5.2)]*.
- Hepatic Decompensation in Patients Co-infected with HIV-1 and Hepatitis C *[see Warnings and Precautions (5.4)]*.
- Pancreatitis *[see Warnings and Precautions (5.5)]*.
- New Onset or Worsening Renal Impairment *[see Warnings and Precautions (5.6)]*.
- Psychiatric Symptoms *[see Warnings and Precautions (5.7)]*.
- Nervous System Symptoms *[see Warnings and Precautions (5.8)]*.
- Rash *[see Warnings and Precautions (5.10)]*.
- Hepatotoxicity *[see Warnings and Precautions (5.11)]*.
- Convulsions *[see Warnings and Precautions (5.12)]*.
- Lipid Elevations *[see Warnings and Precautions (5.13)]*.
- Bone Effects *[see Warnings and Precautions (5.14)]*.
- Immune Reconstitution Syndrome *[see Warnings and Precautions (5.15)]*.
- Fat Redistribution *[see Warnings and Precautions (5.16)]*.

6.1 Clinical Trials Experience

Clinical Trials Experience in Adult Patients with HIV-1 Infection

- Lactic Acidosis/Severe Hepatomegaly with Steatosis *[see Boxed Warning, Warnings and Precautions (5.1)]*.
- Exacerbations of Hepatitis B *[see Boxed Warning, Warnings and Precautions (5.2)]*.
- Hepatic Decompensation in Patients Co-infected with HIV-1 and Hepatitis C *[see Warnings and Precautions (5.4)]*.
- Pancreatitis *[see Warnings and Precautions (5.5)]*.
- New Onset or Worsening Renal Impairment *[see Warnings and Precautions (5.6)]*.
- Psychiatric Symptoms *[see Warnings and Precautions (5.7)]*.
- Nervous System Symptoms *[see Warnings and Precautions (5.8)]*.
- Rash *[see Warnings and Precautions (5.10)]*.
- Hepatotoxicity *[see Warnings and Precautions (5.11)]*.
- Convulsions *[see Warnings and Precautions (5.12)]*.
- Lipid Elevations *[see Warnings and Precautions (5.13)]*.
- Bone Effects *[see Warnings and Precautions (5.14)]*.
- Immune Reconstitution Syndrome *[see Warnings and Precautions (5.15)]*.
- Fat Redistribution *[see Warnings and Precautions (5.16)]*.

- The interaction between efavirenz and the drug was evaluated in a clinical study. All other drug interactions observed are class interactions.

7 DRUG INTERACTIONS

7.6 Drugs without Clinically Significant Interactions

No clinically significant drug interactions have been observed when lamivudine, tenofovir disoproxil fumarate tablets are coadministered with the following: aluminum/magnesium hydroxide antacids, azithromycin, cetirizine, famotidine.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Efavirenz:

Risk Summary. There are retrospective case reports of neural tube defects in infants whose mothers were exposed to efavirenz-containing regimens in the first trimester of pregnancy. Prospective pregnancy data from the Antiretroviral Pregnancy Registry (APR) are insufficient to evaluate the risk of neural tube defects following exposure to efavirenz during the first trimester of human pregnancy. Efavirenz should not be used in the first trimester of pregnancy. Advise pregnant women of the potential risk to the fetus.

Data: Human Data: There are retrospective postmarketing reports of findings consistent with neural tube defects, including myelomeningocele, in infants whose mothers were exposed to efavirenz during the first trimester of pregnancy. Based on prospective reports to the APR of 766 exposed pregnancies (live births, stillbirths, or elective terminations following exposure to efavirenz-containing regimens (including over 600 live births exposed in the first trimester), there was no difference between the overall birth defect rate for efavirenz compared with the background birth defect rate of 2.7% in the U.S. reference population of the Metropolitan Atlanta Congenital Defects Program. As of the reporting date of the APR, there were 19 prospectively reported cases of neural tube defects following first trimester exposure to efavirenz (2.5% [19/766]). The observed prevalence of neural tube defects in these cases is higher than the background rate of neural tube defect. A single case of anophthalmia with first-trimester exposure to efavirenz has also been prospectively reported to the APR. Neural tube defects, including anencephaly and myelomeningocele, have been observed in animal studies *[see Nonclinical Toxicology (13.1)]*.

Animal Data: Effects of efavirenz on embryo-fetal development have been studied in three nonclinical species (mouse, rat, and rabbit). Efavirenz was teratogenic in mice at exposures 0.2 times the exposure in humans at the recommended dose of 600 mg. Lamivudine was not teratogenic in mice at clinically relevant exposures, but *in vivo* effects on embryo-fetal development were observed only at exposures sufficiently in excess of human exposures to be of little clinical significance. Efavirenz was not teratogenic in rats at exposures 17 times the exposure in humans at the recommended dose of 600 mg, but was embryocidal at exposures 0.9 times the exposure in humans at the recommended dose. Efavirenz was not teratogenic in rabbits at exposures 0.3 times the exposure in humans at the recommended dose of 600 mg. Lamivudine was not teratogenic in rabbits at clinically relevant exposures, but was embryocidal at exposures sufficiently in excess of human exposures to be of little clinical significance. Tenofovir disoproxil fumarate was not teratogenic in mice and rats at exposures 2 and 6 times, respectively, the human exposure at the recommended dose of 300 mg, but was embryocidal at these exposures in rats. Tenofovir disoproxil fumarate was not teratogenic in rabbits at exposures 0.3 times the human exposure at the recommended dose of 300 mg, but was embryocidal at this exposure in rabbits.

Lamivudine:

Risk Summary. Available data from an ERS, a synthetic nucleoside analogue with activity against HIV-1 and HBV. The molecular formula is $C_8H_{11}N_3O_3S$ and the molecular weight of 229.26. It is a white to off-white crystalline powder.

Lamivudine



Lamivudine USP is an off – white solid. It is soluble in water.

Tenofovir disoproxil fumarate

Tenofovir disoproxil fumarate is a nucleotide analog reverse transcriptase inhibitor and an HBV reverse transcriptase inhibitor. Tenofovir disoproxil fumarate is a fumaric acid salt of a diester prodrug of tenofovir, which is converted to tenofovir by cellular and viral esterases. The chemical name of tenofovir disoproxil fumarate is [(*R*)-9- [(*R*)- 2- [bis [(isopropoxy-carbonyloxy) methoxy] phosphinyl] propyl] adenine fumarate (1:1) salt. The molecular formula of tenofovir disoproxil fumarate is $C_{19}H_{30}N_5O_{10}P \cdot C_4H_4O_4$ and a molecular weight of 635.52. It is a white to off-white powder.



Tenofovir disoproxil fumarate is a white crystalline powder which is sparingly soluble in methanol and in ethanol.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are a fixed-dose combination of antiviral drugs with activity against HIV-1 *[see Microbiology (12.4)]*.

12.2 Pharmacodynamics

Cardiac Electrophysiology: The effect of efavirenz on the QTc interval was evaluated in an open-label, positive- and placebo-controlled, crossover study in 39 healthy adult subjects who received a single dose of efavirenz 400 mg, moxifloxacin 400 mg (a positive control), and placebo. The mean (90% CI) placebo-adjusted change from baseline in QTcF (Fridericia correction) following the administration of 400 mg daily dose for 7 days was 3.26-fold the mean C_{max} observed in subjects receiving 600 mg daily dose of efavirenz. No subject experienced a change from baseline QTcF > 500 msec. The effect of efavirenz on the mean corrected QT interval was similar on Day 1 and Day 7, indicating that no accumulation occurred.

Viral Dynamics: Following oral administration of a single 600 mg dose of efavirenz to HIV-1 infected subjects, efavirenz plasma concentrations were characterized by a median T_{max} of 5 hours and a mean terminal half-life of 52 to 76 hours. Steady-state plasma concentrations were reached within 6 to 10 days. In 35 HIV-1 infected subjects receiving efavirenz 600 mg once daily for 14 days, the mean (± SD) C_{max} was 4.3 (± 1.1) µM and AUC was 58.1 (± 17.3) µM • hr. Efavirenz is highly bound (approximately 99.5 to 99.75%) to human plasma proteins, predominantly albumin. Efavirenz is metabolized by CYP450 enzymes to primarily hydroxylated metabolites which undergo subsequent glucuronidation before excretion. The primary enzyme responsible for efavirenz metabolism is CYP2B6. CYP3A4 and CYP2A6 are the major enzymes responsible for efavirenz metabolism. Efavirenz has been shown to induce CYP3A4 and CYP2B6 enzymes, which may result in increased metabolism of these substrates. In 47 HIV-1 infected subjects given efavirenz 600 mg once daily for 10 days, approximately 14 to 69% of the intravenous dose of tenofovir is recovered as unchanged drug in the urine. Tenofovir is eliminated by a combination of glomerular filtration and active tubular secretion with a mean renal clearance of 210 to 307 mL/min (more than creatinine clearance) following a single oral dose, the terminal elimination half-life of tenofovir is approximately 17 hours.

Special Populations: Hepatic Impairment: There was no significant or clinically relevant racial difference in efavirenz and tenofovir plasma pharmacokinetics.

Renal Impairment: Efavirenz pharmacokinetics have not been studied in patients with renal impairment.

Gender: There are no significant or clinically relevant gender differences in the pharmacokinetics of efavirenz, lamivudine, and tenofovir disoproxil fumarate.

Geriatric Patients: The pharmacokinetics of lamivudine and tenofovir disoproxil fumarate have not been studied in patients over the age of 65 years.

Pediatric: The combination tablet should not be administered to patients weighing less than 40 kg.

Patients with Hepatitis B and/or Hepatitis C Virus Co-infection:

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are not recommended for patients with impaired renal function (i.e., creatinine clearance less than 50 mL/min) or patients with end-stage renal disease (ESRD) requiring hemodialysis because it is a fixed-dose combination formulation that cannot be adjusted.

Lamivudine: After oral administration of 2 mg/kg of lamivudine twice a day to 8 adults with HIV-1, the peak serum concentration (C_{max}) was 2.2 ± 0.8 µg/mL (mean ± SD), and the area under the plasma concentration-time curve (AUC) and C_{max} increased in proportion to oral dose over the range from 0.5 to 10 mg/kg (the clinical dose is 4 mg/kg). The absolute bioavailability of oral dose was 86 ± 16% (mean ± SD). Less than 10% to the oral solution. Binding of lamivudine to human plasma proteins is low (< 36%). Within 12 hours after a single oral dose of lamivudine to 6 HIV-1 infected adults, 5.2% ± 1.4% (mean ± SD) of the dose was excreted as the trans-sulfoxide metabolite in the urine. The majority of lamivudine is eliminated unchanged in urine by organic cationic secretion and the observed mean elimination half-life ($t_{1/2}$) ranged from 5 to 7 hours in most clinical studies of patients with normal renal function.

Tenofovir Disoproxil Fumarate: Following oral administration of a single 300 mg dose of tenofovir disoproxil fumarate to HIV-1 infected subjects, tenofovir dis

POINTS SAILLANTS DES RENSEIGNEMENTS THÉRAPEUTIQUES

Ce tableau présente les renseignements importants figurant ailleurs dans la présente monographie. Il ne contient pas tous les renseignements pertinents au sujet de TRUVADA. Pour de plus amples renseignements, veuillez lire la présente monographie de produit au complet.

Pour patients avec VHB

Voir Section 10.2 (VHB)

RISQUE DE GRAVE EXACERBATION DE L'HÉPATITE B APRÈS L'ARRÊT DU TRAITEMENT PAR TRUVADA

CHEZ LES PERSONNES ATTEINTES DU VIH-1 ET DU VHB

Chez les personnes atteintes du VIH-1 et du VHB qui cessent de prendre TRUVADA, il faut surveiller étroitement la fonction hépatique pendant au moins quelques mois. Si cela est indiqué sur le plan clinique, il faut envisager la reprise du traitement contre l'hépatite B.

Voir Mises en garde et Précautions (5.2)

Pour patients avec VIH

Voir Mises en garde et Précautions (5.1)

MISES EN GARDE – ACIDOSE LACTIQUE / HÉPATOMÉGALIE SÉVÈRE AVEC STÉATOSE AU TRAITEMENT ULTÉRIEUR DES EXACERBATIONS AIGUËS DE L'HÉPATITE B

Voir toutes les mises en garde et précautions pour de plus amples renseignements sur cette mise en garde.

- **Une acidose lactique et une hépatomégalie avec stéatose, y compris des cas mortels, ont été rapportés avec l'utilisation d'analogues nucléotidiques, y compris la ténofovir et la lamivudine de thérapie antirétrovirale.**
- **Des exacerbations aiguës sévères de l'hépatite B ont été rapportées chez des patients qui ont cessé la prise de TRUVADA (l'hépatite B [VHB] et le virus de l'immunodéficience humaine [VIH-1]) et qui ont également été infectés par le VHB. Ces exacerbations ont été rapportées principalement chez des patients qui ne sont pas traités et, ou non éclairés, selon un traitement contre l'hépatite B. (5.2).**

INDICATIONS ET UTILISATIONS :

Efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés, une combinaison d'un inhibiteur non nucléotidique de la transcriptase inverse (efavirenz) et de deux inhibiteurs nucléotidiques dispose de la transcriptase inverse (lamivudine et fumarate de ténofovir disoproxil), est indiqué pour le traitement de l'infection par le virus de l'immunodéficience humaine de type 1 (VIH-1) chez les patients adultes et pédiatriques pesant au moins 40 kg (1).

POSOLOGIE ET ADMINISTRATION

- Dose recommandée : un comprimé pelliculé 600 mg d'efavirenz, 300 mg de lamivudine et 300 mg de fumarate de ténofovir disoproxil, administré une fois par jour sur un estomac vide, de préférence au coucher (2.1).

FORMES POSOLOGIQUES ET CONCENTRATIONS

Comprimés : 600 mg d'efavirenz, 300 mg de lamivudine et 300 mg de fumarate de ténofovir disoproxil. (3)

CONTRE-INDICATIONS

- Efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés est contre-indiqué chez les patients présentant une hypersensibilité aux substances actives ou à tout composant de ce produit. Pour obtenir une liste complète, veuillez consulter la section des formes posologiques, concentrations, compositions et conditionnement (4).
- Ces administrations avec d'autres produits : Ne pas utiliser avec d'autres produits contenant de l'efavirenz, de la lamivudine ou du fémitivir ou avec des produits contenant de l'emtricitabine. Ne pas administrer avec le dipivoxil d'adéfovir (4, 7.3).
- Des administrations fébrifuge, perfect, folate, c'est produits chez des patients ou n'existe pas de l'infection par le VIH-1 (4); associé une interaction médicamenteuse connue et des données thérapeutiques à base d'interféron et de ribavirine (4, 7.3). Ne pas administrer avec des produits contenant de l'emtricitabine ou de la lamivudine ou avec d'autres produits contenant de fémitivir, comme médicament agonyste, et envisager une réduction posologique en cas de l'interféron alfa, de la ribavirine ou des médicaments. (4, 7.3)
- Ne pas administrer avec des médicaments dont la clairance dépend fortement des antidémétrie de pancréatite ou d'autres facteurs de risque significatifs de pancréatite. Arrêter efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés chez tout patient développant de la pancréatite (4).

MISES EN GARDE ET PRÉCAUTIONS

- Ne pas administrer efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés à une insuffisance rénale légère et la syndrome de rénale. Assurer la clairance adéquate du ténofovir avant d'initier le traitement par le fumarate de ténofovir disoproxil, en surveillant d'albumine), lamivudine et fumarate de ténofovir disoproxil comprimés, chez les patients présentant une insuffisance rénale légère à modérée. Ajuster l'intervalle posologique de ténofovir chez les patients présentant une insuffisance rénale modérée. Ne pas administrer efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés à des patients présentant une insuffisance rénale sévère (4).
- Des cas d'insuffisance rénale aiguë et de syndrome de Fanconi associés à l'utilisation du fumarate de ténofovir disoproxil ont été rapportés. Il faut envisager la possibilité d'une atteinte rénale chez les patients qui présentent ou développent des anomalies des paramètres biologiques rénaux. Il faut envisager l'interruption de l'administration de fumarate de ténofovir disoproxil chez les patients présentant des anomalies des paramètres biologiques rénaux (5.4).

RENSEIGNEMENTS THÉRAPEUTIQUES COMPLETS - CONTENU*

MISES EN GARDE – ACIDOSE LACTIQUE / HÉPATOMÉGALIE SÉVÈRE AVEC STÉATOSE AU TRAITEMENT ULTÉRIEUR DES EXACERBATIONS AIGUËS DE L'HÉPATITE B

Une acidose lactique et une hépatomégalie avec stéatose, y compris des cas mortels, ont été rapportés avec l'utilisation d'analogues nucléotidiques, y compris la lamivudine et le fumarate de ténofovir disoproxil de thérapie antirétrovirale. Des exacerbations aiguës sévères de l'hépatite B ont été rapportées chez des patients qui ont cessé la prise de TRUVADA (l'hépatite B [VHB] et le virus de l'immunodéficience humaine [VIH-1]) et qui ont également été infectés par le VHB. Ces exacerbations ont été rapportées principalement chez des patients qui ne sont pas traités et, ou non éclairés, selon un traitement contre l'hépatite B (5.2).

1 INDICATIONS ET UTILISATIONS

Efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés est destiné à être utilisé aux fins de traiter l'infection par le virus de l'immunodéficience humaine de type 1 (VIH-1) chez les patients adultes et pédiatriques pesant au moins 40 kg.

2 POSOLOGIE ET ADMINISTRATION

2.1 Posologie recommandée chez les patients adultes et pédiatriques pesant au moins 40 kg

La posologie recommandée d'efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés est de 600 mg d'efavirenz, 300 mg de lamivudine et 300 mg de fumarate de ténofovir disoproxil, administrée une fois par jour. Efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés doit administrée sur un estomac vide, préférablement au coucher, afin d'améliorer la tolérance au médicament (*voir Effets indésirables (6.1), Effets indésirables du système nerveux (6.1), Pharmacologie clinique (12.3), Effets indésirables (6.1), et Renseignements sur les conseils à donner aux patients (17.2)*). Efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés doit être avalé entier avec de l'eau.

2.2 Ajustement posologique pour l'insuffisance rénale

Étant donné que l'efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés est contre-indiqué chez les patients présentant une insuffisance rénale modérée ou sévère (clairance de la créatinine estimée ≤ 50 ml/min) ou chez les patients atteints d'une insuffisance rénale au stade terminal (IRAT) nécessitant une hémodialyse.

3 FORMES POSOLOGIQUES ET CONCENTRATIONS

Efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés est disponible sous forme de comprimés pelliculés contenant 600 mg d'efavirenz, 300 mg de lamivudine, et 300 mg de fumarate de ténofovir disoproxil, équivalent à 245 mg de ténofovir disoproxil.

- Les comprimés de 600 mg / 300 mg / 300 mg sont des comprimés pelliculés biconvexes jaunes en forme de capsule, gravés de « 30 » sur une face et de « 030 » sur l'autre face.

4 CONTRE-INDICATIONS

4.1 Hypersensibilité

Efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés sont contre-indiqués chez les patients présentant une hypersensibilité aux substances actives ou à tout composant contenus dans la formulation.

5 MISES EN GARDE ET PRÉCAUTIONS

5.1 Acidose lactique / hépatomégalie sévère avec stéatose

Une acidose lactique et une hépatomégalie sévère avec stéatose, y compris des cas mortels, ont été rapportés avec l'utilisation d'analogues nucléotidiques, y compris la lamivudine et le fumarate de ténofovir disoproxil de thérapie antirétrovirale. La plupart de ces cas ont été rapportés chez des femmes. L'obésité et l'exposition prolongée à un traitement peuvent être des facteurs de risque. Il faut faire preuve de prudence lors de l'administration d'analogues nucléotidiques chez un patient présentant des facteurs de risque connus de maladie hépatique, mais cette acidose lactique peut survenir en l'absence de facteurs de risque connus reconnus. Il faut interrompre le traitement par efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés chez tout patient présentant une hépatotoxicité symptomatique, une acidose métabolique ou lactique ou une stéatose hépatique.

5.2 Patients co-infectés par le VHB et le VHD

Efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés n'est pas indiqué pour le traitement de l'infection par le VHB ou le VHD. Des exacerbations aiguës sévères de l'hépatite B, y compris la décompensation hépatique et l'insuffisance hépatique, ont été rapportées chez des patients qui ont cessé la prise de TRUVADA, un produit contenant de la lamivudine et du fumarate de ténofovir disoproxil, qui est aussi indiqué pour le traitement de l'infection par le VIH-1. Il faut surveiller étroitement la fonction hépatique (tests cliniques et de laboratoire) pendant au moins quelques mois chez les patients co-infectés par le VIH-1 et le VHB qui cessent la prise de TRUVADA. Si nécessaire, une reprise du traitement contre l'hépatite B pourrait être justifiée.

Différences importantes pour les produits contenant de la lamivudine : Lamivudine comprimés contient moins de deux fois plus de dose orale équivalent (en) lamivudine que EPIVIR-HBV comprimés. EPIVIR-HBV a été développé pour le traitement de l'infection chronique par le virus de l'hépatite B (VHB) et n'est pas approuvé pour le traitement de l'infection par le VIH-1. Lamivudine comprimés et lamivudine solution orale ne sont pas indiqués pour le traitement de l'hépatite B chronique chez les patients co-infectés par le VIH-1 et par le VHB.

Le traitement par EPIVIR-HBV et un produit contenant du fumarate de ténofovir disoproxil n'est pas recommandé pour l'hépatite B chronique et n'est pas étudié chez des patients par le VIH-1 non traités ou aux traits. Le traitement par EPIVIR-HBV et un produit contenant du fumarate de ténofovir disoproxil est associé à une réduction de la susceptibilité à la lamivudine à la suite d'une

5.3 Co-administration avec d'autres produits

Ne pas administrer avec d'autres produits contenant comprimés est un produit de combinaison à dose fixe contenant de l'efavirenz, de la lamivudine et du fumarate de ténofovir disoproxil et il ne devrait pas être administré en concomitance avec d'autres médicaments contenant de l'efavirenz, de la lamivudine, du fémitivir ou de l'emtricitabine. Ne pas administrer avec des produits contenant de la lamivudine, comme EPIVIR-HBV, EPIVIR-HBV solution d'association/lamivudine, ZIAGEN® (sulfate d'abacavir/lamivudine/zidovudine), COMBIVIR® (zidovudine/lamivudine), TRIZIVIR® (sulfate d'abacavir/lamivudine/zidovudine), KIVEXA® (abacavir/lamivudine), TRIUMEQ® (dolutégravir/abacavir/lamivudine), VIREAD® (fumarate de ténofovir disoproxil), ATRIPLA® (fumarate de ténofovir disoproxil/emtricitabine/efavirenz), STRIBILD® (fumarate de ténofovir disoproxil/emtricitabine/cobicistat/elvitégravir), TRUVADA® (fumarate de ténofovir disoproxil/emtricitabine), ou ETRAVIRINE (étravirine) ou avec d'autres médicaments contenant de la lamivudine ou de l'emtricitabine.

5.4 Utilisation avec des schémas posologiques à base d'interféron et de ribavirine

Des études *in vitro* ont montré que la ribavirine peut réduire la phosphorylation des analogues nucléotidiques de la cytidine. Il a été démontré que la lamivudine inhibe la phosphorylation de la cytidine endogène. Cependant, l'importance clinique de ces observations *in vitro* est inconnue. Il faut faire preuve de prudence lorsque l'on administre de la ribavirine à des co-administration avec de la lamivudine chez les patients co-infectés par le VIH-1 et l'hépatite C (*voir Interactions médicamenteuses (7.3)*). Les médecins qui envisagent l'utilisation de schémas thérapeutiques à base d'interféron chez des patients co-infectés par le VIH-1 et l'hépatite C doivent consulter la monographie d'interféron pour obtenir des renseignements sur la surveillance et les contre-indications.

5.5 Pancréatite

Des cas de pancréatite ont été rapportés chez des patients traités par la lamivudine. Il faut arrêter efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés chez tout patient développant de la pancréatite (*voir Effets indésirables (6.2)*).

5.6 Troubles hépatiques

Il faut faire preuve de prudence lors de l'administration d'efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés à des patients présentant une insuffisance hépatique modérée ou sévère (cirrhose). Il n'est pas nécessaire d'ajuster la dose de ce produit chez les patients présentant une insuffisance hépatique légère (score de Child-Pugh de 5 à 6). Il faut surveiller étroitement les patients présentant une insuffisance hépatique modérée ou sévère (cirrhose) pendant les premiers mois de traitement par efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés. Il faut envisager l'interruption de l'administration de efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés chez les patients présentant des anomalies persistantes ou aggravées des tests de la fonction hépatique (ALAT, ASAT, bilirubine, albumine, temps de prothrombine) (*voir Pharmacologie clinique (12.3)*).

Les interactions médicamenteuses avec l'étravirine sont résumées dans le Tableau 3 (pour les données pharmacocinétiques, voir Pharmacologie Clinique (12.3). Tableau 4 et 5). Ce tableau comprend les interactions potentiellement significatives, mais n'est pas exhaustif.

Tableau 3 : Interactions médicamenteuses établies et autres interactions potentiellement significatives : Une modification de la dose de ce médicament peut être recommandée sur la base des études sur les interactions médicamenteuses ou prévues

Médicament concomitant par classe : Dénominations du médicament	Effet	Commentaire clinique
Antiacides : Maalox, Mylanta	↓ de 1 à 2 h	Surveiller la fréquence d'apparition de survenue si nécessaire.
Antiarrhythmiques : Carbamazépine	↓ carbamazépine* ↓ étravirine*	Il n'y a pas assez de données pour donner une recommandation posologique pour l'étravirine. Un autre médicament antiviral devrait être envisagé.
Flécaïnine, Lidocaïnine, Mexilétine, Propafénone, Quinidine	↓ antiarrhythmiques*	Surveiller les concentrations plasmatiques des antiarrhythmiques et envisager une adaptation potentielle de la posologie. Des concentrations plasmatiques des antiarrhythmiques et, ou de l'étravirine peuvent être nécessaires.
Antihistaminiques : Bupropion	↓ bupropion*	Les augmentations de la dose de sertraline devront être envisagées en fonction de la réponse clinique.
Sertraline	↓ sertraline*	
Antimycobactériens : Rifampicine, Rifapentine, Rifabutine	↓ étravirine* ↓ diarylrhénylpyrimidine* ↓ rifamycine*	Les augmentations de la dose de sertraline devront être envisagées en fonction de la réponse clinique.
Posaconazole	↑ posaconazole*	Il n'est pas recommandé d'administrer du posaconazole avec de l'étravirine.
Anti-infectieux : Clarithromycine, Itraconazole, Kétoconazole, Voriconazole	↑ clarithromycine* ↑ itraconazole* ↑ kétoconazole* ↑ voriconazole*	Il faudrait envisager des alternatives aux antifongiques azolés ou aux antibiotiques macrolides. Surveiller les concentrations plasmatiques de l'étravirine.
Anticoagulants : Warfarine	↓ étravirine*	Augmenter la dose quotidienne de warfarine de 2,5 mg. Surveiller l'INR et ajuster la dose de warfarine si nécessaire.
Antiépileptiques : Carbamazépine, Phénobarbital, Phénytoïne	↓ antiépileptiques* ↓ étravirine*	Il n'est pas recommandé d'administrer de la carbamazépine avec de l'étravirine. Il faudrait envisager des alternatives aux antiépileptiques. Surveiller les concentrations plasmatiques de l'étravirine.
Antipsychotiques : Rispéridone	↓ rispéridone*	Il faudrait envisager des alternatives à la rispéridone. L'ajustement de la dose de rispéridone en fonction de la réponse clinique peut être nécessaire.
Atorvastatine*	↓ atorvastatine*	Il faut commencer par la dose la plus faible d'atorvastatine et ajuster la dose en fonction de la réponse clinique.
Lovastatine*	↓ lovastatine*	Il faut envisager une augmentation de la dose de lovastatine en fonction de la réponse clinique.
Pravastatine*	↓ pravastatine*	Il faut envisager une augmentation de la dose de pravastatine en fonction de la réponse clinique.
Rosuvastatine*	↓ rosuvastatine*	Il faut commencer par la dose la plus faible de rosuvastatine et ajuster la dose en fonction de la réponse clinique.
Simvastatine*	↓ simvastatine*	Il faut envisager une augmentation de la dose de simvastatine en fonction de la réponse clinique.
Bêta-bloquants : Métoprolol, Timolol	↓ bêta-bloquants*	Il faut envisager une augmentation de la dose de bêta-bloquants en fonction de la réponse clinique.
Bloqueurs des canaux calciques : Diltiazem, Félodipine, Nicardipine, Nifédipine, Vérapamil	↓ bloqueurs des canaux calciques*	Il faut envisager une augmentation de la dose de bloqueurs des canaux calciques en fonction de la réponse clinique.
Contraceptifs hormonaux : Éthinylestradiol / Norgestimate	↓ éthinylestradiol ↓ norgestimate	Il faut envisager d'autres méthodes contraceptives.
Implant Etonogestrel	↓ étonogestrel	Les données cliniques démontrent une diminution de l'exposition à l'étonogestrel lors de l'administration concomitante d'étravirine. Il faut envisager d'autres méthodes contraceptives.
Immunosuppresseurs : Cyclosporine, sirolimus, tacrolimus	↓ immunosuppresseurs*	Il faut envisager une augmentation de la dose de l'immunosuppresseur nécessaire pour les immunosuppresseurs. Une surveillance étroite des concentrations thérapeutiques est recommandée pendant la co-administration et après l'arrêt de l'étravirine. Il faut envisager une augmentation de la dose de l'immunosuppresseur nécessaire pour les immunosuppresseurs. Une surveillance étroite des concentrations thérapeutiques est recommandée pendant la co-administration et après l'arrêt de l'étravirine.
Narcotique analgésique : Méthadone	↓ méthadone*	Il faut envisager une augmentation de la dose de méthadone en fonction de la réponse clinique.

* = Interaction établie ou potentielle à cause d'une étude d'interaction médicamenteuse ou d'une analyse d'interaction médicamenteuse prévue sur la base des voies métaboliques connues.

↑ = Augmentation potentielle ou observée ; ↓ = diminution potentielle ou observée ; ↔ = aucun changement observé ou prévu

6 EFFETS INDÉSIRABLES

6.1 Médicaments associés couramment significatifs

Aucun questionnaire prospectif n'est recommandé lorsque efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés est administré à des patients présentant une insuffisance rénale modérée ou sévère (clairance de la créatinine estimée ≤ 50 ml/min) ou chez des patients atteints d'une insuffisance rénale au stade terminal (IRAT) nécessitant une hémodialyse.

7 INTERACTIONS MÉDICAMENTEUSES

7.1 Médicaments non affectés par l'étravirine

Les études *in vitro* ont montré que la ribavirine peut réduire la phosphorylation des analogues nucléotidiques de la cytidine. Il a été démontré que la lamivudine inhibe la phosphorylation de la cytidine endogène. Cependant, l'importance clinique de ces observations *in vitro* est inconnue. Il faut faire preuve de prudence lorsque l'on administre de la ribavirine à des co-administration avec de la lamivudine chez les patients co-infectés par le VIH-1 et l'hépatite C (*voir Effets indésirables (6.2)*). Les médecins qui envisagent l'utilisation de schémas thérapeutiques à base d'interféron chez des patients co-infectés par le VIH-1 et l'hépatite C doivent consulter la monographie d'interféron pour obtenir des renseignements sur la surveillance et les contre-indications.

5.7 Syndrome immunitaire de reconstitution immunitaire

Un syndrome de reconstitution immunitaire a été rapporté chez des patients infectés par le VIH-1 traités avec des associations d'antirétroviraux, y compris avec efavirenz, lamivudine et fumarate de ténofovir disoproxil. Au cours de la phase initiale du traitement par association d'antirétroviraux, les patients dont le système immunitaire répond peuvent développer une réponse inflammatoire aux infections asymptomatiques latentes ou résiduelles (comme les infections mycobactériennes atypiques, cytomégalovirus, pneumonie à *Pneumocystis jirovecii* [PCP] ou tuberculose), qui peut nécessiter une évaluation et un traitement.

5.8 Ostéonécrose

Des cas d'ostéonécrose ont été rapportés, particulièrement chez des patients présentant des facteurs de risque connus de maladie avancée liée au VIH-1 ou d'exposition prolongée à un traitement antirétroviral combiné. La fréquence de ces événements est inconnue. Les patients doivent être informés qu'il faut consulter un médecin s'ils ressentent des douleurs articulaires, des raideurs ou des difficultés à bouger.

5.9 Exacerbations aiguës de l'hépatite B après l'arrêt du traitement

Des exacerbations aiguës sévères de l'hépatite B ont été rapportées chez des patients qui ont cessé la prise de TRUVADA, un produit contenant de la lamivudine et du fumarate de ténofovir disoproxil, qui est aussi indiqué pour le traitement de l'infection par le VIH-1. Il faut surveiller étroitement la fonction hépatique (tests cliniques et de laboratoire) pendant au moins quelques mois chez les patients co-infectés par le VIH-1 et le VHB qui cessent la prise de TRUVADA. Si nécessaire, une reprise du traitement contre l'hépatite B pourrait être justifiée.

5.10 Ostéoporose

Des cas d'anomalies de la densité minérale osseuse ont été rapportés. Il faut envisager une évaluation de la densité minérale osseuse chez les adultes et les enfants si cela est cliniquement indiqué.

5.11 Hypercalciurie

La surveillance des enzymes hépatiques avant et pendant le traitement est recommandée chez les patients présentant une maladie hépatique sous-jacente, y compris une hépatite chronique active B ou C, ou chez les patients présentant des élévations marquées des transaminases. Il faut envisager l'interruption du traitement par efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés chez les patients présentant des signes de détérioration de la maladie hépatique (*voir Effets indésirables (6.2)*), y compris l'apparition d'ictère, de prurit, de coloration foncée de l'urine, de selles décolorées, d'hépatite cutanée ou digestive ou d'asthénie. Il faut également considérer la réduction ou l'interruption de la prise de TRUVADA chez les patients présentant des élévations persistantes des tests de la fonction hépatique ou une hépatotoxicité symptomatique.

5.12 Cancérogenèse

Des convulsions ont été observées chez des adultes (7 %) et des enfants (10 %) recevant efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés lors des études cliniques. Les patients présentant des antécédents de convulsions peuvent présenter un risque accru de convulsions après l'administration de efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés. Il faut surveiller les patients traités par efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés pour détecter l'apparition de convulsions. Les patients doivent être informés que des convulsions peuvent survenir pendant le traitement par efavirenz, lamivudine et fumarate de ténofovir disoproxil comprimés.

5.13 Troubles psychiatriques

Des événements psychiatriques graves et indésirables ont été rapportés chez des patients traités par efavirenz. Dans une étude contrôlée, des événements psychiatriques graves de tout grade clinique (psychose, dépression) ont été observés chez 3 % des patients traités par des schémas thérapeutiques contenant de l'efavirenz pendant quatre ans par rapport à 2 % des patients traités par des schémas thérapeutiques ne contenant pas d'efavirenz. Des idées suicidaires graves ont été rapportées chez des patients traités par efavirenz (1,9 %) ou par des schémas thérapeutiques ne contenant pas d'efavirenz (0,8 %). Des événements psychiatriques graves ont été rapportés chez 2,4 % des patients recevant de l'efavirenz pendant 48 semaines dans une étude contrôlée, contre 0,9 % des patients ne recevant pas d'efavirenz. Des événements psychiatriques graves ont été rapportés chez 1,6 % des patients recevant de l'efavirenz pendant 48 semaines dans une étude contrôlée, contre 0,4 % des patients ne recevant pas d'efavirenz. Des événements psychiatriques graves, tels qu'une dépression sévère, des idées suicidaires, ou des comportements suicidaires, ont été observés chez