

Guidance on Linezolid

This guidance represents the Food and Drug Administration's (FDA's) current thinking on this topic. It does not create or confer any rights for or on any person and does not operate to bind FDA or the public. You can use an alternative approach if the approach satisfies the requirements of the applicable statutes and regulations. If you want to discuss an alternative approach, contact the Office of Generic Drugs.

Active ingredient: Linezolid

Form/Route: For Suspension/Oral

Recommended studies: **2 Options: BCS or In-Vivo Studies**

I. BCS Waiver option:

It may be possible to request a waiver of in-vivo testing of this product provided that the appropriate documentation regarding high solubility, high permeability and rapid dissolution as detailed in the Guidance for Industry: *Waiver of In Vivo Bioavailability and Bioequivalence for Immediate – Release Solid Oral Dosage Forms Based on the Biopharmaceutics Classification System* is submitted in the application. You may use information contained in the approved labeling of the reference product. Peer reviewed articles may not contain the necessary details of the testing for the Agency to make a judgment regarding the quality of the studies. A decision regarding the acceptability of the waiver request can only be made upon review of the data submitted in the application.

II. In-Vivo option:

1. Type of study: Fasting
Design: Single-dose, two-way, crossover *in-vivo*
Strength: 100 mg/5ml
Subjects: Healthy males and nonpregnant females, general population
Additional Comments:
2. Type of study: Fed
Design: Single-dose, two-way, crossover *in-vivo*
Strength: 100 mg/5ml
Subjects: Healthy males and nonpregnant females, general population
Additional comments:

Analytes to measure: Linezolid in plasma

Bioequivalence based on (90% CI): Linezolid

Waiver request of in-vivo testing: Not Applicable

Dissolution test method and sampling times:

Please note that a **Dissolution Methods Database** is available to the public at the OGD website at <http://www.accessdata.fda.gov/scripts/cder/dissolution/>. Please find the dissolution information for this product at this website. Please note that a dosage unit is based on the labeled concentration of the suspension product. Please use the dosage unit (5 ml). A total of 12 units from 12 different bottles should be used. Specifications will be determined upon review of the application.