

Contains Nonbinding Recommendations

Draft – Not for Implementation

Draft Guidance on Hydroxychloroquine Sulfate

October 2024

This draft guidance, when finalized, will represent the current thinking of the Food and Drug Administration (FDA, or the Agency) on this topic. It does not establish any rights for any person and is not binding on FDA or the public. You can use an alternative approach if it satisfies the requirements of the applicable statutes and regulations. To discuss an alternative approach, contact the Office of Generic Drugs.

In general, FDA's guidance documents do not establish legally enforceable responsibilities. Instead, guidances describe the Agency's current thinking on a topic and should be viewed only as recommendations, unless specific regulatory or statutory requirements are cited. The use of the word *should* in Agency guidances means that something is suggested or recommended, but not required.

Active Ingredient:	Hydroxychloroquine sulfate
Dosage Form:	Tablet
Route:	Oral
Strength:	200 mg
Recommended Studies:	Two options: (1) Biopharmaceutics Classification System (BCS)-based biowaiver or (2) one in vivo bioequivalence study with pharmacokinetic endpoints

I. Option 1: BCS Class III-based biowaiver

A waiver request of in vivo testing for this product may be considered provided that the appropriate documentation regarding high solubility, very rapid dissolution, and the test product formulation is qualitatively the same and quantitatively similar as detailed in the most recent version of the FDA guidance for industry on *M9 Biopharmaceutics Classification System-Based Biowaivers^a* is submitted in the application. Applicants may use the information contained in the approved labeling of the reference listed drug (RLD). Peer reviewed articles may not contain the necessary details of the testing for the Agency to make a judgment regarding the quality of the studies. A decision regarding the acceptability of the waiver request can only be made upon assessment of the data submitted in the application.

II. Option 2: One in vivo bioequivalence study with pharmacokinetic endpoints

1. Type of study: Fed
Design: Single-dose, two- treatment, randomized, parallel in vivo
Strength: 200 mg
Subjects: Healthy males and non-pregnant, non-lactating females
Additional comments: None

Analyte to measure: Hydroxychloroquine in whole blood

Bioequivalence based on (90% CI): Hydroxychloroquine

Waiver request of in vivo testing: Not applicable

Dissolution test method and sampling times: The dissolution information for this drug product can be found in the FDA's Dissolution Methods database, <http://www.accessdata.fda.gov/scripts/cder/dissolution>. Conduct comparative dissolution testing on 12 dosage units for each of the test product and RLD.¹ Specifications will be determined upon review of the abbreviated new drug application.

Document History: Recommended April 2011; Revised April 2020, October 2024

Unique Agency Identifier: PSG_009768

^a For the most recent version of a guidance, check the FDA guidance website at <https://www.fda.gov/regulatory-information/search-fda-guidance-documents>.

¹ If the RLD is not available, refer to the most recent version of the FDA guidance for industry on *Referencing Approved Drug Products in ANDA Submissions*.