

Draft Guidance on Amphetamine Aspartate; Amphetamine Sulfate; Dextroamphetamine Saccharate; Dextroamphetamine Sulfate

October 2024

This draft guidance, when finalized, will represent the current thinking of the Food and Drug Administration (FDA, or the Agency) on this topic. It does not establish any rights for any person and is not binding on FDA or the public. You can use an alternative approach if it satisfies the requirements of the applicable statutes and regulations. To discuss an alternative approach, contact the Office of Generic Drugs.

In general, FDA's guidance documents do not establish legally enforceable responsibilities. Instead, guidances describe the Agency's current thinking on a topic and should be viewed only as recommendations, unless specific regulatory or statutory requirements are cited. The use of the word *should* in Agency guidances means that something is suggested or recommended, but not required.

Active Ingredients:	Amphetamine aspartate; Amphetamine sulfate; Dextroamphetamine saccharate; Dextroamphetamine sulfate
Dosage Form:	Tablet
Route:	Oral
Strengths:	1.25 mg; 1.25 mg; 1.25 mg; 1.25 mg, 1.875 mg; 1.875 mg; 1.875 mg; 1.875 mg, 2.5 mg; 2.5 mg; 2.5 mg; 2.5 mg, 3.125 mg; 3.125 mg; 3.125 mg; 3.125 mg, 3.75 mg; 3.75 mg; 3.75 mg; 3.75 mg, 5 mg; 5 mg; 5 mg; 5 mg, 7.5 mg; 7.5 mg; 7.5 mg; 7.5 mg
Recommended Study:	One in vivo bioequivalence study with pharmacokinetic endpoints
1.	Type of study: Fasting Design: Single-dose, two-treatment, two-period crossover in vivo Strength: 7.5 mg; 7.5 mg; 7.5 mg; 7.5 mg Subjects: Healthy males and non-pregnant, non-lactating females Additional comments: None
Analytes to measure:	D-amphetamine and l-amphetamine, measured separately in plasma
Bioequivalence based on (90% CI):	D-amphetamine and l-amphetamine

Waiver request of in vivo testing: 1.25 mg; 1.25 mg; 1.25mg; 1.25 mg, 1.875 mg; 1.875 mg; 1.875 mg; 1.875 mg, 2.5 mg; 2.5 mg; 2.5 mg; 2.5 mg, 3.125 mg; 3.125 mg; 3.125 mg; 3.125 mg, 3.75 mg; 3.75 mg; 3.75 mg; 3.75 mg, 5 mg; 5 mg; 5 mg; 5 mg strengths based on (i) acceptable bioequivalence study on the 7.5 mg; 7.5 mg; 7.5 mg; 7.5 mg strength, (ii) acceptable in vitro dissolution testing of all strengths, and (iii) proportional similarity of the formulations across all strengths

Dissolution test method and sampling times: The dissolution information for this drug product can be found in the FDA's Dissolution Methods database, <http://www.accessdata.fda.gov/scripts/cder/dissolution/>. Conduct comparative dissolution testing on 12 dosage units for each of all strengths of the test product and reference listed drug (RLD).¹ Specifications will be determined upon review of the abbreviated new drug application.

Document History: Recommended December 2008; Revised November 2013, October 2024

Unique Agency Identifier: PSG_011522

¹ If the RLD is not available, refer to the most recent version of the FDA guidance for industry on *Referencing Approved Drug Products in ANDA Submissions*.