

Draft Guidance on Cephalexin

May 2026

This draft guidance, when finalized, will represent the current thinking of the Food and Drug Administration (FDA, or the Agency) on this topic. It does not establish any rights for any person and is not binding on FDA or the public. You can use an alternative approach if it satisfies the requirements of the applicable statutes and regulations. To discuss an alternative approach, contact the Office of Generic Drugs.

In general, FDA’s guidance documents do not establish legally enforceable responsibilities. Instead, guidances describe the Agency’s current thinking on a topic and should be viewed only as recommendations, unless specific regulatory or statutory requirements are cited. The use of the word *should* in Agency guidances means that something is suggested or recommended, but not required.

Active Ingredient: Cephalexin

Dosage Form: Suspension

Route: Oral

Strengths: EQ 100 mg Base/mL | EQ 125 mg Base/5 mL |
EQ 250 mg Base/5 mL

Reference Listed Drug: NDA 050406

Recommended Study: One in vivo bioequivalence study with pharmacokinetic endpoints

1. Class of study: Bioequivalence
Type of study: Fasting
Design: Single-dose, two-treatment, two-period crossover in vivo
Strength: EQ 250 mg Base/5 mL
Dose: EQ 250 mg Base/5 mL administered as 5 mL
Subjects: Healthy males and non-pregnant, non-lactating females

Analyte to measure: Cephalexin in plasma

Bioequivalence based on (90% CI): Cephalexin

Waiver request of in vivo testing of additional strengths: Justification based on (i) an acceptable bioequivalence study on the EQ 250 mg Base/5 mL strength, (ii) acceptable comparative in vitro dissolution studies between the additional strengths and the EQ 250 mg Base/5 mL strength using 12 units per strength, and (iii) proportional similarity of the formulations across all strengths

Dissolution: Dissolution test(s) should be included for quality control and to support a waiver request of in vivo testing of additional strengths. For the quality control dissolution method, provide a dissolution method development report for the test product containing information and data that demonstrate appropriateness of the selected dissolution method¹ and sampling times, such as the discriminating ability to detect changes in critical quality attributes that could potentially impact drug product performance.

For drug products containing high solubility drug substances that meet the rapidly dissolving criteria, demonstration of discriminating ability may not be needed. For additional information, refer to the guidance for industry *Dissolution Testing and Acceptance Criteria for Immediate-Release Solid Oral Dosage Form Drug Products Containing High Solubility Drug Substances*.^a

Document History: Recommended December 2016; Revised October 2024, May 2026

^a We update guidances periodically. For the most recent version of a guidance, refer to the FDA guidance website at <https://www.fda.gov/regulatory-information/search-fda-guidance-documents>.

¹ Applicant-developed, United States Pharmacopeia drug product monograph or Dissolution Methods database, <https://www.accessdata.fda.gov/scripts/cder/dissolution/index.cfm>