

Draft Guidance on Momelotinib Dihydrochloride

May 2025

This draft guidance, when finalized, will represent the current thinking of the Food and Drug Administration (FDA, or the Agency) on this topic. It does not establish any rights for any person and is not binding on FDA or the public. You can use an alternative approach if it satisfies the requirements of the applicable statutes and regulations. To discuss an alternative approach, contact the Office of Generic Drugs.

In general, FDA's guidance documents do not establish legally enforceable responsibilities. Instead, guidances describe the Agency's current thinking on a topic and should be viewed only as recommendations, unless specific regulatory or statutory requirements are cited. The use of the word *should* in Agency guidances means that something is suggested or recommended, but not required.

Active Ingredient:	Momelotinib dihydrochloride
Dosage Form:	Tablet
Route:	Oral
Strengths:	EQ 100 mg Base, EQ 150 mg Base, EQ 200 mg Base
Recommended Study:	One in vivo bioequivalence study with pharmacokinetic endpoints

1. Type of study: Fasting
Design: Single-dose, two-treatment, two-period crossover in vivo
Strength: EQ 200 mg Base
Subjects: Healthy males and non-pregnant, non-lactating females
Additional comments: Exclude subjects with abnormal complete blood counts. Monitor for signs and symptoms of infection and bleeding during the study. Females of reproductive potential should use non-hormonal contraception during the study and continue to use effective contraception for one week after the last dose of the study.

Analyte to measure: Momelotinib in plasma

Bioequivalence based on (90% CI): Momelotinib

Waiver request of in vivo testing: EQ 100 mg Base and EQ 150 mg Base strengths based on (i) an acceptable bioequivalence study on the 200 mg strength, (ii) acceptable in vitro dissolution testing of all strengths, and (iii) proportional similarity of the formulations across all strengths

Dissolution test method and sampling times: The dissolution information for this drug product can be found in the FDA's Dissolution Methods database, <http://www.accessdata.fda.gov/scripts/cder/dissolution/>. Conduct comparative dissolution testing on 12 dosage units for each of all strengths of the test product and reference listed drug (RLD)¹. Specifications will be determined upon review of the abbreviated new drug application.

Document History: Recommended May 2025

Unique Agency Identifier: PSG_216873

¹ If the RLD is not available, refer to the most recent version of the FDA guidance for industry on *Referencing Approved Drug Products in ANDA Submissions*.