

ORDER SET/PROTOCOL TEMPLATE



This template is to be used as a guideline for developing procedures, protocols, and/or order sets around the use of ADASUVE® to ensure that REMS requirements are being met.

INDICATION

- Agitation associated with bipolar I disorder
- Agitation associated with schizophrenia

PATIENT SCREENING

PRIOR TO DOSING:

Assess for acute respiratory symptoms

- wheezing cough
- dyspnea other: _____

Assess for respiratory signs (include chest auscultation)

- wheezing other: _____

Inquire and/or check medical records for:

- current diagnosis or history of asthma, COPD, or other lung disease associated with bronchospasm
- current use of medications to treat airways disease, such as asthma or COPD
- history of bronchospasm following ADASUVE treatment
- known hypersensitivity (e.g., serious skin reaction) to loxapine or amoxapine

If any of the above screening is positive, DO NOT USE ADASUVE. ADASUVE is contraindicated in these patients.

DOSE ADMINISTERED

TIME: _____

Limit ADASUVE use to a single dose per patient within a 24-hour period.

POST-TREATMENT OBSERVATION/MONITORING

Patient must be monitored every 15 minutes for at least 1 hour after treatment for signs/symptoms of bronchospasm.

MONITORING

TIME: _____

- Heart rate (rate per minute): _____
- Respiratory rate (rate per minute): _____
- Chest auscultation (clear/other) _____
- Check for respiratory signs/symptoms
 - wheezing cough
 - dyspnea chest tightness
 - other: _____

TIME: _____

- Heart rate (rate per minute): _____
- Respiratory rate (rate per minute): _____
- Chest auscultation (clear/other) _____
- Check for respiratory signs/symptoms
 - wheezing cough
 - dyspnea chest tightness
 - other: _____

TIME: _____

- Heart rate (rate per minute): _____
- Respiratory rate (rate per minute): _____
- Chest auscultation (clear/other) _____
- Check for respiratory signs/symptoms
 - wheezing cough
 - dyspnea chest tightness
 - other: _____

TIME: _____

- Heart rate (rate per minute): _____
- Respiratory rate (rate per minute): _____
- Chest auscultation (clear/other) _____
- Check for respiratory signs/symptoms
 - wheezing cough
 - dyspnea chest tightness
 - other: _____

Note: Treat bronchospasm with inhaled short-acting beta-agonist bronchodilator (e.g., albuterol) and other measures as clinically indicated.